



光華劍橋
Guanghua Cambridge School

Comparison of Two Models to Calculate the Period of Compound Pendulum with Liquid Damping

Jintian Wang AS8

上海光華學院劍橋國際中心
Guanghua Cambridge International School



CONTENTS

- **General review**
- **Experiment & data**
- **Analysis & conclusion**
- **Reflection & summarize**
- **Acknowledgements**



GENERAL REVIEW




Fig. 1. For an extended object of moment of inertia $I = M(a^2 + k^2)$ about the pivot O, the period of small oscillation depends only on the distance a from the pivot to the center of mass, and the gyradius k .

Fig 1.




Fig 2.

← **Compound Pendulum**

GENERAL REVIEW



光華劍橋
Guanghua Cambridge School

Methods:

- $f \propto v$ $f = kR\omega = kR\dot{\theta}$ Proportional model
- $f \propto v^2$ $f = kR^2\omega^2 = kR^2\dot{\theta}^2$ Square model

THEORY DERIVATION



光華劍橋
Guanghua Cambridge School

$$T = 2\pi \sqrt{\frac{I^2}{IR(m - \rho V)g - 1/4(k^2 R^4)}}$$

$$f = kv$$

$$T = 2\pi \sqrt{\frac{I^2 - 2kR^3\theta(I - kR^3)}{gR(m - \rho V)(I - 2kR^3\theta)}}$$

$$f = kv^2$$

ROUGH EXPERIMENT



光華劍橋
Guanghua Cambridge School

$$I\ddot{\theta} = WR - fR - FR$$


$$1. f = kR\omega = kR\dot{\theta}$$

$$I\ddot{\theta} = (m - \rho V)gR\theta - kR^2\dot{\theta}$$

$$\text{For } \theta \ll 5, \sin \theta \approx \theta_{[3]}$$

里, $\omega_0 = 0.7$ 。显然, 当幅角较小时, 两者的振动周期、振幅都比较接近, 当幅角更小时, 复摆的振动可看作简谐振动。而实际的复摆, 幅角小于 5° 的

Fig 6.



ROUGH EXPERIMENT



光華劍橋
Guanghua Cambridge School



Fig 7.

Nov. 19th




Fig 8.

Water tank



THEORY DERIVATION

2023.09-10:

Collect references,
work out the first
version of formula
about the pendulum

2023.12-24.01:

Design experiment,
make theoretical
expectations, calculate
various answers

2024.03-04:

Analyze data, finish
the essay, submit to
tutor for final revision

2023.07-08:

Determine research
topic, make the first
step to the question,
have a basic
understanding

2023.11-12:

Study how to plot graph
and how to meet the
needs of experiments(e.g.
python+3d design)

2024.01-02:

Complete experiment,
check if any mistakes,
Draw graphs and
compare the
expectations with reality



ROUGH EXPERIMENT DATA

Table 2. Data of rough experiments

No.	Frame/f	10T/s	T/s
1	740	6.167	0.614
2	736	6.133	
3	732	6.100	
4	738.5	6.154	

Variable	Value	Unit
k	0.84 ± 0.02	$kg\ s^{-1}$
I	0.60 ± 0.003	$g\ m^2$
m	36.0 ± 0.5	g
v	5.00 ± 0.05	ml
R	25.7 ± 0.1	mm

$$\ddot{\theta} + r_1 \dot{\theta} + r_2 \theta = 0$$

Let $\delta = \frac{c}{2m}$ for $m\ddot{x} + C\dot{x} + kx = 0$, and $\delta > 1$ is overdamping

$$\delta = \frac{r_1}{2} = \frac{kR^2}{2I} = 0.15 < 1 \text{ not over damped}$$

ROUGH EXPERIMENT DATA



光華劍橋
Guanghua Cambridge School

```
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 # 给定的参数
5 I = 0.000655
6 F = 0.7246667
7 a = 5
8
9 # 不同的 k 值
10 k_values = [0, 0.05, 0.5, 1, 2, 5, 10, 20, 50, 100]
11
12 # 定义理论拟合曲线
13 def function(x,k):
14     return 2 * np.pi / k * np.sin(k * x)
15
16 # 绘制不同 k 值下的曲线
17 x_values = np.linspace(0, 35, 1000)
18 colors = ['black', 'red', 'blue', 'green', 'purple', 'brown', 'pink', 'gray', 'olive', 'cyan']
19
20 for i, k_value in enumerate(k_values):
21     y_values = [function(x, k_value) for x in x_values]
22     plt.plot(x_values, y_values, color=colors[i])
23
24 # 设置图形的标题和坐标轴标签
25 plt.title('Effect of Diffusion Coefficient on the Period of Oscillation')
26 plt.xlabel('R (cm)')
27 plt.ylabel('T (s)')
28
29 # 显示图形
30 plt.show()
```




Fig 7. Curves of θ about t [python]


EXPERIMENT



光華劍橋
Guanghua Cambridge School

No.	Equipment Name	Device Model	Number
1	Mass block	2cm * 1cm in cross-section	9
2	Iron stand	100cm in length	4
3	Clamps	Iron with rubber	2
4	Rod	M8, 10cm in length	2
5	Bearing	10mm in inner diameter	2
6	Silicon gaskets	M8, 1mm in thickness	2
7	Nut	M8, hexagon	2
8	Rubber band	8cm in original length	1
9	Water bucket	20cm in diameter	1
10	Camera	iPhone 11 *0.125 in slow-motion	1
11	Vanier caliper	15cm max.	1

Table 2. Equipment list



Container overview


EXPERIMENT



光華劍橋
Guanghua Cambridge School



b. Design display [blender]



c. Container ready for experiments

Fig 3. Container and design diagram [blender]

EXPERIMENT



光華劍橋
Guanghua Cambridge School





Fig 5. Clamps, water bucket, and equipment

EXPERIMENT



光華劍橋
Guanghua Cambridge School



Fig 6. An overview

DATA



光華劍橋
Guanghua Cambridge School

No.	material	A/cm ²	h/cm	V/ml	m1/g	m2/g	m3/g	m/g	density
1	plastic	2.00	2.82	5.64	7.78	7.76	7.76	7.77	1.38
2		2.00	4.00	8.00	10.97	10.98	11.02	10.99	1.37
3		2.00	4.68	9.36	11.47	11.48	11.47	11.47	1.23
4	Al	2.00	3.34	6.68	16.63	16.63	16.63	16.63	2.49
5		2.00	4.32	8.64	22.10	22.08	22.08	22.09	2.56
6		2.00	5.34	10.68	27.64	27.64	27.62	27.63	2.59
7	Fe	2.00	3.14	6.28	46.26	46.25	46.27	46.26	7.37
8		2.00	4.48	8.96	61.89	61.90	61.90	61.90	6.91
9		2.00	5.28	10.56	77.92	77.95	77.95	77.94	7.38

Table 4. Table of masses

DATA



光華劍橋
Guanghua Cambridge School

No.	R/cm	F1/f	F2/f	F3/f	T1/s	T2/s	T3/s	T/s
1	7.96	290	292	284	1.208	1.217	1.183	1.203
2	7.67	226	232	244	0.941	0.965	1.017	0.974
3	5.76	192	209	199	0.802	0.872	0.828	0.834
4	5.00	465	425	448	1.939	1.772	1.866	1.859
5	4.28	227	223	223	0.946	0.929	0.929	0.935
6	3.92	222	219	222	0.925	0.913	0.925	0.921
7	3.50	107	107	105	0.448	0.446	0.438	0.444
8	3.13	258	252	246	1.075	1.050	1.025	1.050
9	2.41	318	295	301	1.325	1.229	1.254	1.269
10	1.50	361	368	378	1.503	1.533	1.575	1.537
11	1.05	439	451	451	1.831	1.879	1.879	1.863

Table 5. Table of Recordings

DATA



光華劍橋
Guanghua Cambridge School

- Useful Data
- Wrong Data
- Squared Model
- Proportional Model

$$k = 0.5 \text{ kg s}^{-1}$$
$$V = 185.0 \text{ ml}$$
$$L = 20.55 \text{ cm}$$
$$\theta = 5^\circ$$




Fig 8. Curves of t about R [python]



ANALYSIS


```
Variance of squared model: 0.0007568985124692573  
Variance of proportional model: 0.0007271175137054576
```

Fig 9. Log of the program using the first four sets of data [python]

```
Variance of squared model: 0.0003286635921692313  
Variance of proportional model: 0.013896866472079114
```

Fig 10. Log of the program using the last four sets of data [python]

- **Small** radius of rotation
 - **Proportional** model fits better
- **Large** radius of rotation
 - **Square** model fits better



THEORATICAL ANALYSIS



光華劍橋
Guanghua Cambridge School

A screenshot of a program log showing two lines of output. The first line is '0.0007568985124692573' and the second line is '0.0007271175137054576'. A red rectangular box highlights the decimal parts of both numbers, specifically the digits '568985124692573' and '271175137054576'.

```
0.0007568985124692573  
0.0007271175137054576
```

Fig 9. Log of the program using the first four sets of data [python]

Small radius of rotation

- Almost no difference in variance - 0.0003%

Does not **strongly support** the proportional model

- Further stimulations needed

THEORATICAL ANALYSIS



光華劍橋
Guanghua Cambridge School

Fig 11. Effects of k on the period about radius [python]

$$k = 0 \text{ kg s}^{-1}$$

$$k = 0.05 \text{ kg s}^{-1}$$

$$k = 0.5 \text{ kg s}^{-1}$$

$$k = 1.0 \text{ kg s}^{-1}$$


$$k = 2.0 \text{ kg s}^{-1}$$

$$k = 2.5 \text{ kg s}^{-1}$$

$$V = 184.95 \text{ ml}$$

$$m = 258.82 \text{ g}$$

$$\theta = 5^\circ$$




Square
model
+
Proportional
model

k changing, other variables constant

THEORATICAL ANALYSIS



光華劍橋
Guanghua Cambridge School



Small radius

- Both - proportional easier


Large radius

- Squared more accurate

REFLECTION



光華劍橋
Guanghua Cambridge School



- Two of the eleven sets of experiments are **useless**

REFLECTION





Fig 12. Buoyance higher than weight

- Two of the ten sets of experiments are useless
- Easily making mistakes **affects the efficiency**

REFLECTION



光華劍橋
Guanghua Cambridge School



- Two of the ten sets of experiments are useless
- Easily making mistakes affects the efficiency
- M8 screw causing **fluid resistance**



SUMMARIZE

2023.09-10:

Collect references,
work out the first
version of formula
about the pendulum

2023.12-24.01:

Design experiment,
make theoretical
expectations, calculate
various answers

2024.03-04:

Analyze data, finish
the essay, submit to
tutor for final revision

2023.07-08:

Determine research
topic, make the first
step to the question,
have a basic
understanding

2023.11-12:

Study how to plot graph
and how to meet the
needs of experiments(e.g.
python+3d design)


2024.01-02:

Complete experiment,
check if any mistakes,
Draw graphs and
compare the
expectations with reality

SUMMARIZE



光華劍橋
Guanghua Cambridge School



0s to 5s.png



0s to 30s.png




3d.PNG




all.jpg




clamps.jpg




desmos-graph
(1).png




desmos-graph.
png



effect 1.png



effect2.png



Figure_1.png



hang in air.jpg




hang in
water.jpg



hang.jpg



model
nothing.jpg



model on
groundf.jpg



model.jpg



pendulum.jpg



support all.jpg



support.jpg



water rank
all.jpg



water tank.jpg

SUMMARIZE



光華劍橋
Guanghua Cambridge School

This represents a second-order non-linear ordinary differential equation, distinct from our previous solution. A new mathematical approach is required due to the presence of a square on the first derivative. Assume $y = \left(\frac{d\theta}{dt}\right)^2$, where θ is a function about θ . According to

$$\frac{d}{d\theta} y = \frac{d}{d\theta} \left(\frac{d\theta}{dt}\right)^2 = 2 \frac{d^2\theta}{dt^2}$$

[11] can be shifted into another differential equation about θ .

$$\frac{dy}{d\theta} + 2r_1 y + 2r_2 \theta = 0 \quad [12]$$

Here [12] is a linear differential equation. By solving y we can get the function θ through some extent. However, details still need to be listed due to this is not an easy job. Convert [12] into

$$dy + 2(r_1 y + r_2 \theta) d\theta = 0 \quad [13]$$

Let $m = 2(r_1 y + r_2 \theta)$, where $dm = 2r_1 dy + 2r_2 d\theta$, and [13] can be transformed into $y + m d\theta = 0$. According to $m = \frac{m}{2r_1} - \frac{r_2}{r_1} \theta$. Differentiate y , $dy = \frac{dm}{2r_1} - \frac{r_2}{r_1} d\theta$. Use instead into [13].

$$\frac{dm}{2r_1} - \frac{r_2 d\theta}{r_1} + m d\theta = 0$$

Simplify the equation and we get an integrable expression of dm and $d\theta$.

$$\frac{dm}{2r_1 - 2r_1 m} = d\theta$$

Upon integration, the function m is determined as:

$$m = \frac{A}{e^{2r_1 \theta}} - \frac{r_2}{r_1}$$

Take into $y = \frac{m}{2r_1} - \frac{r_2}{r_1} \theta$.

$$y = \frac{A}{e^{2r_1 \theta}} - \frac{r_2 \theta}{r_1} + \frac{r_2}{2(r_1)^2}$$

Where A is integral constant. To calculate A , consider when $\theta = \theta_0$ i.e., maximum amplitude, $\frac{d\theta}{dt} \Big|_{\theta=\theta_0} = 0$, along with $y = \left(\frac{d\theta}{dt}\right)^2 = 0$. Through substitution,

$$A = \frac{r_1}{2e^{2r_1 \theta_0}} (2r_1 \theta_0 - 1) e^{2r_1 \theta_0} \quad [14]$$

Now y and A have been worked out, and it should be followed by deriving θ from y , which

$$\frac{A}{e^{2r_1 \theta}} - \frac{r_2 \theta}{r_1} + \frac{r_2}{2(r_1)^2} = \left(\frac{d\theta}{dt}\right)^2$$

Take the square root of both LHS and RHS sides. Separate the variables.

$$\sqrt{\frac{A}{e^{2r_1 \theta}} - \frac{r_2 \theta}{r_1} + \frac{r_2}{2(r_1)^2}} = \frac{d\theta}{dt}$$

Expand and integrate both sides.

$$\int dt = \int \left(\frac{A}{e^{2r_1 \theta}} - \frac{r_2 \theta}{r_1} + \frac{r_2}{2(r_1)^2} \right)^{\frac{1}{2}} d\theta \quad [15]$$

From using Maclaurin series $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$, the approximate value of e can be simply calculated by the first three terms. And

$$e^{-2r_1 \theta} \approx 1 + \frac{-2r_1 \theta}{1!} + \frac{(-2r_1 \theta)^2}{2!} \quad [16]$$

Take into [15], RHS can be exchanged into

$$\int \left(\left(A + \frac{r_2}{2(r_1)^2} \right) - \left(2Ar_1 + \frac{r_2^2}{r_1} \right) \theta + 2Ar_1^2 \theta^2 \right)^{\frac{1}{2}} d\theta \quad [17]$$

By completing the square method, notice that

$$\left(A + \frac{r_2}{2(r_1)^2} \right) - \left(2Ar_1 + \frac{r_2^2}{r_1} \right) \theta + 2Ar_1^2 \theta^2 = \left(\frac{A}{2} - \frac{r_2^2}{4(r_1)^2} \right) + (2A) \left(\left(\frac{1}{2} + \frac{r_2}{4(r_1)^2 2A} \right) - r_1 \theta \right)^2$$

And [17] is equivalent to

$$\int \frac{\sqrt{-3A}}{\sqrt{\frac{r_2^2}{4(r_1)^2} - A^2}} \sqrt{1 - \frac{(2Ar_1 \theta - (A + \frac{r_2}{2(r_1)^2}))^2}{\frac{r_2^2}{4(r_1)^2} - A^2}} d\theta \quad [18]$$

From $\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1}(x)$, take [18] into [17].

$$\Delta t = -\frac{\sqrt{-3A}}{r_1} \sin^{-1} \left[\frac{2Ar_1 \theta - \left(A + \frac{r_2}{2(r_1)^2} \right)}{\sqrt{\frac{r_2^2}{4(r_1)^2} - A^2}} \right] \quad [19]$$

ACKNOWLEDGEMENTS



光華劍橋
Guanghua Cambridge School

- My tutor –Dr. Hou
- Teachers from physics group
- My parents
- My classmates –Tony Zhao
- Other students and friends who support me



光華劍橋
Guanghua Cambridge School

Thank you for your listening

上海光華學院劍橋國際中心
Guanghua Cambridge International School