

Basics

Notations

- $\binom{V}{k} := \{A : A \subseteq V \wedge |A| = k\}$
- $[n] := \{1, \dots, n\} \subset \mathbb{N}$
- Power set** $2^X := \{A : A \subseteq X\}$

Graphs

- Definition:** $G = (V, E)$ with $E \subseteq V^2$, $V \cap E = \emptyset$
- Vertex:** $v \in V$ for graph $G = (V, E)$
 - v incident with $e \Leftrightarrow v \in e$
 - v_1, v_2 ends of $e \Leftrightarrow e = v_1 v_2$
 - v_1, v_2 adjacent/neighbors $\Leftrightarrow v_1 v_2 \in E$
- Edge:** $e = \{x, y\} \in E$ for graph $G = (V, E)$ (short $e = xy$)
 - e edge at $v \Leftrightarrow v$ incident with e
 - e joins $v_1, v_2 \Leftrightarrow e = v_1 v_2$
 - xy is X - Y -edge $\Leftrightarrow x \in X \wedge y \in Y$
 - e_1, e_2 adjacent/neighbors $\Leftrightarrow \exists v : v \in e_1 \wedge v \in e_2$
- Vertex sets:**
 - $V(G) = V$ for graph $G = (V, E)$
 - $X \subset V(G)$ independent \Leftrightarrow no $x_1, x_2 \in X$ are adjacent
 - neighborhood of $v \in V(G)$: $N(v) = \{u \in V(G) : uv \in E(G)\}$
- Edge sets:**
 - $E(G) = E$ for graph $G = (V, E)$
 - $E(X, Y)$: set of edges between $X \subset V(G)$ and $Y \subset V(G)$
 - $E(x, Y)$: set of edges between vertex $x \in V(G)$ and $Y \subset V(G)$
 - $E(v)$: set of edges at $v \in V(G)$
- Order:** $= |V(G)|$, short $|G|$
- Size:** $= |E(G)|$, short $\|G\|$
- Trivial graph:** graph of order 0 or 1
- Isomorphic** (G_1 to another graph G_2 , write $G_1 \cong G_2$ or even $G_1 = G_2$):
 - \exists bijection $f : V_1 \rightarrow V_2 : \{u, v\} \in E_1 \Leftrightarrow \{f(u), f(v)\} \in E_2$
- Graph union:** $G \cup G' = (V(G) \cup V(G'), E(G) \cup E(G'))$
- Graph intersection:** $G \cap G' = (V(G) \cap V(G'), E(G) \cap E(G'))$
- Graph multiplication:** $G * G'$: join all $v \in G$ with all $v' \in G'$ (with $V(G) \cap V(G') = \emptyset$)
- Subgraph** G' of G (write $G' \subseteq G$): if $V(G') \subseteq V(G)$ and $E(G') \subseteq E(G)$
 - G contains G'
 - G' proper subgraph of G : if $G' \subseteq G$ and $G' \neq G$
 - G' induced subgraph of G : $G' \subseteq G$ and $E(G')$ contains all edges of G with both ends in $V(G')$, $V(G')$ induces G' , write $G' = G[X]$ (with $X = V(G')$)
 - Edge-induced subgraph: subgraph induced by $X \subseteq E(G)$, note $G[X]$
 - G' spanning subgraph of G : $V(G') = V(G)$
- Supergraph:** G of G' (write $G \supseteq G'$): as above.
- Vertex cover:** $V' \subseteq V(G)$ s.t. any $e \in E(G)$ is incident to a vertex in V'
- Graph subtraction:**
 - $G - U = G[V(G) \setminus U]$ for some vertex set U
 - $G - v = G[V(G) \setminus \{v\}]$ for some vertex v
 - $G - G' = G[V(G) \setminus V(G')]$ for some graph G'
- Edge addition:** $G + F = (V(G), V(E) \cup F)$ for some $F \subseteq V(G)^2$
- Complement:** $\overline{G} = (V(G), V^2 \setminus E(G))$
- Line graph** of G : $L(G) = (E(G), \{xy \in E(G)^2 : x, y \text{ adjacent in } G\})$
- Complete graph:** (X, X^2) with vertex set X
 - K_n : complete graph on n vertices

Vertex degrees

- Degree** of $v \in V$: $d(v) = \deg(v) = |N(v)|$
 - $v \in V(G)$ isolated: $d(v) = 0$
 - $v \in V(G)$ leaf: $d(v) = 1$
 - number of vertices of odd degree is even
- Minimum degree** of graph G : $\delta(G) = \min\{d(v) : v \in V(G)\}$
- Maximum degree** of graph G : $\Delta(G) = \max\{d(v) : v \in V(G)\}$
- Degree sum:** $\sum_{v \in V(G)} \deg(v) = 2|E(G)|$
- Average degree** of graph G : $d(G) = \frac{1}{|G|} \sum_{v \in V} d(v)$
 - $\delta(G) \leq d(G) \leq \Delta(G)$
- k-regular graph:** $\forall v \in V(G) : d(v) = k$
 - cubic graph: 3-regular graph
- Vertex-Edge-ratio** of graph G : $\varepsilon(G) = \frac{\|G\|}{|G|}$
 - $\varepsilon(G) = \frac{1}{2} d(G)$
 - every graph with $\|G\| \geq 1$ has $H \subseteq G$ with $\delta(H) > \varepsilon(H) \geq \varepsilon(G)$

Paths and Cycles

- Path:** $(\{v_1, \dots, v_n\}, \{\{v_1, v_2\}, \dots, \{v_{n-1}, v_n\}\})$ (read: $v_0 v_n$ -path)
 - shorthand: (v_1, \dots, v_n)
 - v_0, v_n linked by path
 - v_0, v_n end-vertices/ends of path
 - v_1, \dots, v_{n-1} inner vertices of path
- Length** (of path): $= |E(P)| \neq |V(P)|$
- Shorthands** ($0 \leq i \leq j \leq k$):
 - $P = (x_0, \dots, x_k)$, $\hat{P} = (x_1, \dots, x_{k-1})$
 - $Px_i = (x_0, \dots, x_i)$, $P\hat{x}_i = (x_0, \dots, x_{i-1})$
 - $x_i P = (x_i, \dots, x_k)$, $\hat{x}_i P = (x_{i+1}, \dots, x_k)$
 - $x_i P x_j = (x_i, \dots, x_j)$, $\hat{x}_i P \hat{x}_j = (x_{i+1}, \dots, x_{j-1})$
- Path concatenation:** $Px \cap xQy \cap yR = PxQyR$
- A-B-path:** $V(P) \cap A = \{x_0\} \wedge V(P) \cap B = \{x_n\}$
- H-path:** graph H , P meets H exactly in its ends
- Independent:** two ab -paths are independent \Leftrightarrow they only share a and b
- Subgraph separation:** $X \subset V(G)$ separates $A, B \subset V(G) \Leftrightarrow$ any A - B -path has vertex in X

Rest

- Degree sequence:** multiset of degrees of vertices in $V(G)$
 - graphic: deg. seq. (d_1, \dots, d_n) , iff
 - $d_1 + \dots + d_n$ even
 - $\sum_{i=1}^k d_i \leq k(k-1) + \sum i = k + 1^n \min(d_i, k) \quad (\forall 1 \leq k \leq n)$
- Adjacency matrix:** $A(G) = \mathbb{R}^{n \times n} \ni A_{i,j} = \begin{cases} 1, & ij \in E \\ 0, & \text{else} \end{cases}$
- Incidence graph** of G : $IG = (V \cup E, \{\{v, e\} : v \in e, e \in E\})$
- Eulerian:** if it contains an Eulerian tour
- Connected:** for any two vertices there is a link between them
 - spanning tree: if G is connected, then it has a spanning tree
 - peeling leaves: vertices can be ordered v_1, \dots, v_n s.t. $G[\{v_1, \dots, v_i\}]$ is connected for $i \in \{1, \dots, n\}$

Digraph

- Definition:** $G = (V, E)$ with vertex set V and edge set $E \subseteq \{(u, v) : u, v \in V, u \neq v\}$

Multigraph

- Definition:** $G = (V, E)$ with vertex set V and multiset E of V -pairs

Hypergraph

- Definition:** $G = (V, E)$ with vertex set V and edge set $E \subseteq 2^V = \{A : A \subseteq V\}$

Cycle

- Definition:** $C_n := (\{v_1, \dots, v_n\}, \{\{v_1, v_2\}, \dots, \{v_{n-1}, v_n\}, \{v_n, v_1\}\})$
- Shorthand:** (v_1, \dots, v_n, v_1)
- Length** (of cycle): $= |V| \equiv |E|$
- Cyclic subgraph:** If $\delta(G) \geq 2$, then G has cycle with length $\geq \delta + 1$

Walk

- Definition:** non-empty alternating sequence of vertices and edges
$$v_0 e_0 \dots e_{k-1} v_k$$
with $e_i = v_i v_{i+1}$, length $k \in \mathbb{N}$
 - closed: if $v_0 = v_k$
 - even: if k is even
 - odd: if k is odd
- Eulerian tour:**
 - Definition: closed walk with
 - no edges of G are repeatedly used
 - all edges of G are used
 - Even degrees: G connected has Euler tour $\Leftrightarrow \forall v \in V(G) : \deg(v)$ even
- $v_0 v_k$ -walk: walk starting at v_0 and ending at v_k
- Induces path:** $\exists uv$ -walk $\Rightarrow \exists uv$ -path
- Odd closed walk, odd cycle:** G has odd closed walk $\Rightarrow G$ has odd cycle

Connected component

- Definition:** maximal connected subgraph (connected, but any supergraph isn't)

Block

- **Block:** maximal 2-connected subgraph or bridge
 - share ≤ 1 vertices with one another
- **Block-cut-vertex graph**
 - V = set of blocks \cup set of vertices
 - $E = \{\{v, B\} : v \in V(B), \text{ cut-vertex } v, \text{ block } B\}$
 - block-cut-vertex graph of connected graph is tree

Acyclic graph, Forest

- **Definition:** Graph with no cycle as subgraph

Tree

- **Definition:** Graph that is connected and acyclic
 - $\Leftrightarrow G$ is connected and $\forall e \in E(G) : G - e$ is disconnected (*minimal-connected*)
 - $\Leftrightarrow G$ is acyclic and $\forall xy \notin E(G) : G \cup xy$ has cycle (*maximal-acyclic*)
 - $\Leftrightarrow G$ is connected and 1-degenerate ($\forall G' \subseteq G : \delta(G') \leq 1$)
 - $\Leftrightarrow G$ is connected and $\|G\| = |G| - 1$
 - $\Leftrightarrow G$ is acyclic and $\|G\| = |G| - 1$
 - $\Leftrightarrow \forall u, v \in V(G) \exists$ unique uv -path
- **Special trees:** path, star, spider, caterpillar, broom
- **Leaf existence:** Tree $T, |T| \geq 2 \Rightarrow T$ has leaf
- **Edge count:** Tree $T, |T| = n \Rightarrow \|T\| = n - 1$

Bipartite graph

- **Definition:** G is bipartite $\Leftrightarrow G$ contains no cycles of odd length
 - *complete bipartite:* $K_{m,n} = (A \cup B, \{a, b\} : a \in A, b \in B)$
- **Matchings:**
 - *saturating:* $G = (A \cup B, E)$ has matching saturating A
 $\Leftrightarrow \forall S \subseteq A : N(S) \geq |S| \quad (N(S) := \{b \in B : ab \in E, a \in S\})$
 - *nearly:* $G = (A \cup B, E), \forall S \subseteq A : |N(S)| \geq |S| - d \quad (d \geq 1).$
 $\Rightarrow \exists$ matching M saturating all but at most d vertices of A
- **Matching vs vertex cover:** size of largest matching = size of smallest vertex cover

Matching

- **Definition:** graph with $\delta(G) = \Delta(G) = 1$
- **Perfect matching:** spanning + matching subgraph of G (aka 1-factor)
 - *existence:* G has perfect matching $\Leftrightarrow \forall S \subseteq V(G) : q(G - S) \leq S$
($q(G)$ = number of components in G with odd order)

Factors

- **k-factor:** spanning k -regular subgraph (easy to find)
- **f-factor:** spanning subgraph $H \subseteq G$ with $\deg_H(v) = f(v),$
 $f : V(G) \rightarrow \{0, 1, \dots\}$ with $f(v) \leq \deg(v) \quad (\forall v \in V)$
- **H-factor** (aka *perfect H-packing*): spanning subgraph s.t. each component is $\cong H$
 - *existence:* if $\delta(G) \geq (1 - \frac{1}{k}|V(G)|)$ and k divides $|G|$, then G has K_k -factor

Connectivity

- **k-connected:** if $|G| > k$ and deleting $< k$ vertices does not disconnect G
- **k-linked:** if for any $2k$ vertices $(s_1, \dots, s_k, t_1, \dots, t_k) \exists$ pairwise disjoint $s_i t_i$ -paths (*note:* k -connected $\nRightarrow k$ -linked)
- **Vertex-connectivity:** $\kappa(G) = \max\{k : G \text{ is } k\text{-connected}\}$
- **l-edge-connected:** if deleting $< l$ edges does not disconnect G
- **Edge-connectivity:** $\kappa'(G) = \max\{l : G \text{ is } l\text{-edge-connected}\}$
- **Vertex- vs Edge-connectivity:** $\kappa(G) \leq \kappa'(G) \leq \delta(G)$
- **Three-connected + contraction:** 3-connected $\Leftrightarrow \exists$ separate G_0, \dots, G_k with
 $G_0 = K_4, G_k = G, G_i = G_{i+1} \circ xy$
with $\deg(x), \deg(y) \geq 3$
- **Three-connected + decontraction:** all 3-connected graphs can be built by iteratively de-contracting vertices of K_4
- **Average degree ≥ 4 :** has k -connected subgraph ($k \geq 2$)

Cuts

- **Cut-Set:** $X \subseteq V(G) \cup E(G)$ s.t. #components in $(G - X)$ greater than in G
- **Cut-Vertex:** Cut-Set consisting of single vertex
- **Cut-Edge** (or *bridge*): Cut-Set consisting of single edge
- **Menger's theorem:** for $A, B \subseteq V(G)$:
min # of vertices separating A and B = max # of disjoint A - B -paths
- **Menger global:**
 1. k -connected $\Leftrightarrow \forall a, b \in V(G) \exists k$ pairwise independent ab -paths

2. k -edge-connected $\Leftrightarrow \forall a, b \in V(G) \exists k$ pairwise edge-disjoint ab -paths

Ear-decomposition

- **Definition:** G has *ear-decomposition* $\Leftrightarrow \exists$ sequence of graphs G_0, \dots, G_k with $G_k = G, G_0 = \text{cycle}, G_{i+1}$ obtained from G_i by attaching "ear" (path that shares only endpoints with G_i)
- **2-connected** $\Leftrightarrow \forall$ cycles C in G there is ear-decomposition starting at C

Edge contraction

- **Contraction:**

$$G \circ xy = ((V \setminus \{x, y\}) \cup v_{xy},$$

$$(E \setminus \{e : x \in E \vee y \in e\}) \cup \{v_{xy}z : z \in (N_G(x) \cup N_G(y)) \setminus \{x, y\}\})$$

with $xy \in E(G)$

- **De-contraction:** if $\exists xy \in E(G) : \kappa(G \circ xy) \geq 3$
(for G with $\kappa(G) \geq 3, |G| \geq 5$)

Planar graph tools

- **Homeomorphism:** $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ continuous s.t. f^{-1} is also continuous
- **Arc:** homeomorphic image of $[0, 1]$ in \mathbb{R}^2 under f
 - *endpoints:* $f(0)$ and $f(1) \leadsto$ arc "joins" endpoints
 - *polynomial arc:* arc that is union of finitely many straight line segments
- **Region** $Y \subseteq \mathbb{R}^2 \setminus X$: any two points $\in Y$ could be joined by arc and Y is maximal ($X \subseteq \mathbb{R}^2$)
- **Boundary** of $X \subseteq \mathbb{R}^2$:
 $\partial X = \{y \mid \forall \varepsilon > 0 : B(y, \varepsilon) \text{ contains points of } X \text{ and not of } X\}$
- **Jordan curve theorem:** If $X \subseteq \mathbb{R}^2$ and homeomorphic to $\{\bar{x} : \text{dist}(\bar{x}, 0) = 1\}$ (*unit circle*), then $\mathbb{R}^2 \setminus X$ has two regions R_1, R_2 and $\delta R_1 = X = \delta R_2$.

Plane graph

- **Definition:** graph such that $E(G)$ is set of arcs in \mathbb{R}^2 and endpoints of arcs in $E(G)$ are vertices and:
 - $\forall e, e' \in E, e \neq e' : e$ and e' have distinct sets of edge sets
 - $\forall e \in E, \hat{e} = e \setminus \{\text{endpoints}\}$ doesn't contain any vertices and points from other arcs
- **Faces:** regions of $\mathbb{R}^2 \setminus (\bigcup_{e \in E} e \cup V)$
- **Maximally plane:** no edges can be added without breaking planarity
 - *plane triangulation:* every face is bounded by triangle \Leftrightarrow graph is maximally plane
- **Edge limitation 1:** Plane graph: $|G| \geq 3 \Rightarrow \|G\| \leq 3n - 6$
- **Edge limitation 2:** Plane graph with no Δ : $\|G\| \leq 2|G| - 4$
- **Properties:** Let G be plane graph and $H \subseteq G$.
 - *face inheritance:* $\forall f \in F(G) \exists f' \in F(H) : f' \supseteq f$
 - *border inheritance:* $\delta f \subseteq H \Rightarrow f' = f$
 - *edge-border relations:* $e \in E(G), f \in F(G) \Rightarrow e \subseteq \delta f \vee \delta f \cap \hat{e} = \emptyset$
 - *edges in circles:*
 $e \in E(G)$ is edge of a cycle $\Rightarrow e$ is on boundary of exactly 2 faces
not edge of a cycle $\Rightarrow e$ is on boundary of exactly 1 face
 - *faces in cycles:* $f_1, f_2 \in F(G). f_1 \neq f_2 \wedge \delta f_1 = \delta f_2 \Rightarrow G$ is cycle
 - *cyclic boundaries:* $\kappa(G) \geq 2 \Rightarrow$ each face is bounded by cycle
 - *plane forests:* plane forests have exactly 1 face
- **Dual multigraph:** Given plane G :
 1. Insert vertex in each face
 2. Put edge \tilde{e} between vertices if respective faces share e (s.t. \tilde{e} and e cross once)
 3. *Result:* Dual graph G' of G (plane multigraph)
 \leadsto faces of G properly k -colored $\Leftrightarrow \exists$ proper k -coloring of vertices of G'

Planar graph

- **Definition:** graph s.t. \exists plane graph G' and bijection $f : V(G) \rightarrow V(G')$ s.t. $\forall u, v \in V(G), uv \in E(G) : f(u), f(v)$ are endpoints of arc in G'
- **Planar embedding** of G : f from the definition
- **Planar because of minors:** The following statements are equivalent:
 - G is planar
 - $G \not\preceq MK_5 \wedge G \not\preceq MK_{3,3}$
 - $G \not\preceq TK_5 \wedge G \not\preceq TK_{3,3}$
- **Euler's formula:** If G is connected plane graph with f faces, then
 $|G| - \|G\| + f = 2$
- $\delta(G)$ **limitation:** Planar graph $\delta(G) \leq 5$
- **Non-planar graphs:** K_5 and $K_{3,3}$ are not planar
- **Kuratowski's lemmas:**
 1. $(TK_5 \subseteq G \vee TK_{3,3} \subseteq G) \Leftrightarrow MK_5 \subseteq G \vee MK_{3,3} \subseteq G$
 2. $\kappa(G) \geq 3 \wedge MK_5 \not\subseteq G \wedge MK_{3,3} \not\subseteq G \Rightarrow G$ is planar

3. $\kappa(G) \geq 3$, G edge-maximal wrt not containing TX . If S is vertex-cut of G , $|S| \leq 2 \wedge G = G_1 \cup G_2$, $S = V(G_1) \cap V(G_2)$, then G_i is edge-maximal with no TX and S induces an edge
 4. $|G| \geq 3$, G edge-maximal wrt not containing TK_5 and $TK_{3,3} \Rightarrow \kappa(G) \geq 3$
- **2-cell** (embedding of G on surface S): any closed simple curve in any region of $S - G$ is continuously contractible into a point
 - **Euler characteristic**: G embedded on surface $S \Rightarrow n - e + f = \text{Euler characteristic}$ is invariant
 - **Euler genus**: $n - e + f = 2 - 2\gamma \rightsquigarrow \text{Euler genus } 2\gamma \text{ of } S$
 - **Heawood's formula**: $\chi(G) \leq \left\lfloor \frac{7 + \sqrt{1 + 48\gamma}}{2} \right\rfloor$
 $f(\gamma)$, Heawoods number
(for G embedded on S with Euler char $2 - 2\gamma$)
 - **Klein bottle**: $K_{f(\gamma)}$ is embeddable on S , unless S is *klein bottle*

Minors

- **MH**: $G \stackrel{(\star)}{=} MH$ is *minor* of H if
 - $V(G) = V_1 \cup \dots \cup V_n$ with $n = |H|$
 - $G[V_i]$ connected ($\forall i = 1, \dots, n$)
 - If $V(H) = \{v_1, \dots, v_n\}$ and $v_i v_j \in E(H)$, then \exists edge between V_i and V_j (\star) : *Notation abuse*: MH is class of graphs
- **Branch sets**: V_i 's from above
- **Extended branch graph**: Branch set together with incident edges
- **Minor** (H of G , noted $H \preceq G$): $\Leftrightarrow MH \subseteq G$
 $\sim H \preceq G \Leftrightarrow H$ can be obtained by edge/vertex deletions + contractions.
- **Topological minor**: H is topological minor if $TH \subseteq G$ where TH is built from H by subdividing edges
- **Note**: $TH \subseteq MH$

Coloring

- **Co-clique number**: $\alpha(G)$ = size of largest independent set
- **Clique number**: $\omega(G)$ = size of largest clique
- **Proper coloring**: $c : V(G) \rightarrow [k]$ with $c(u) \neq c(v)$ ($\forall uv \in E(G)$)
- **Equitable coloring**: proper coloring + color classes have almost (± 1) equal size
 - *existence*: any graph has equitable coloring in $(\Delta(G) + 1)$ colors
- **4-color-theorem**: G planar $\Rightarrow \chi(G) \leq 4$
- **ij-flip**: $c' : V(G) \rightarrow [k]$ is ij -flip at $v \in V(G)$
 $\Leftrightarrow c'$ obtained by flipping colors i and j in max. conn. component containing v
-

Chromatic number

- **Definition**: $\chi(G) = \min\{k : G \text{ has proper coloring with } k \text{ colors}\}$
- **Examples**: $\chi(C_{2n}) = 2$, $\chi(C_{2n+1}) = 3$
- **Properties**:
 - $\chi(G) \geq \omega(G)$
 - $\chi(G) \geq \frac{|G|}{\alpha(G)}$
 - $\chi(G) \leq \Delta(G) + 1$ (*greedy coloring*)
 - G connected, not complete, no odd cycles $\Rightarrow \chi(G) \leq \Delta(G)$

Perfect graph

- **Definition**: $\forall H \subseteq_{\text{ind}} G : \chi(H) = \omega(H)$
- **Perfect complement**: G is perfect $\Leftrightarrow \overline{G}$ is perfect
- **Perfect graph conjecture**: G is perfect $\Leftrightarrow C_{2k+1} \not\subseteq G$ for $k \geq 2 \wedge \overline{C_{2k+1}} \not\subseteq G$

Posets

- **Definition**: antisymmetric, reflexive, transitive relation on X
(write $x \leq y$ instead of (x, y))
- **Incidence poset** of G : poset whose cover diagram is represented by IG with vertices all below the edges
- **Poset dimension**: $\dim(R)$ = smallest $k \in \mathbb{N} : R$ is intersection of k total orders
- **Poset dimension in planar graphs**: G planar $\Leftrightarrow \dim(\text{incidence poset}) \leq 3$

List-colorings

- **L-list-colorable**: if $\exists c : V \rightarrow \mathbb{N} \forall v \in V : c(v) \in L(v)$
(for list of colors $L(v) \subseteq \mathbb{N}$ for each vertex, adjacent vertices receive different colors)
- **k-list-colorable/-choosable**: if G is L -list-colorable for each list L
- **List chromatic number**: $\chi_l(G) = \text{ch}(G)$
 $= \min\{k : G \text{ is } L\text{-colorable } \forall L : V \rightarrow 2^{\mathbb{N}} : |L(v)| = k \forall v \in V(G)\}$
 - $\chi_l(G) \geq \chi(G)$ because we can choose $L(v) = \{1, \dots, k\}$ ($\forall v \in V(G)$)
 - often $\chi_l(G) \gg \chi(G)$ (see $K_{m,n} : \chi = 2, \chi_l \approx \log n$)