

# Thesis title: Quantum-mechanical systems in traps and density functional theory

Jørgen Høgberget

## Aims

The aim of this thesis is to study the structure of fermionic systems using density functional theory, parametrizing the functionals using *ab initio* variational and diffusion function Monte Carlo techniques. The thesis will explore various Monte Carlo optimization strategies and use these to define the best possible density functional using the so-called adiabatic connection method. Furthermore, if possible, the developed density functional will be applied in so-called time-dependent density functional approaches to quantum mechanical systems with external time-dependent fields.

## General introduction to possible physical systems

What follows here is a general introduction to systems of confined electrons in two or three dimensions. However, although the thesis will focus on such systems, the codes will be written so that other systems of trapped fermions or eventually bosons can be handled. Examples could be neutrons in trap as done in Ref. [1] or ions in various traps [2].

Strongly confined electrons offer a wide variety of complex and subtle phenomena which pose severe challenges to existing many-body methods. Quantum dots in particular, that is, electrons confined in semiconducting heterostructures, exhibit, due to their small size, discrete quantum levels. The ground states of, for example, circular dots show similar shell structures and magic numbers as seen for atoms and nuclei. These structures are particularly evident in measurements of the change in electrochemical potential due to the addition of one extra electron,  $\Delta_N = \mu(N+1) - \mu(N)$ . Here  $N$  is the number of electrons in the quantum dot, and  $\mu(N) = E(N) - E(N-1)$  is the electrochemical potential of the system. Theoretical predictions of  $\Delta_N$  and the excitation energy spectrum require accurate calculations of ground-state and of excited-state energies. Small confined systems, such as quantum dots (QD), have become very popular for experimental study. Beyond their possible relevance for nanotechnology, they are highly tunable in experiments and introduce level quantization and quantum interference in a controlled way. In a finite system, there can not, of course, be a true phase transition, but a cross-over between weakly and strongly correlated regimes is still expected. There are several other fundamental differences between quantum dots and bulk systems: (a) Broken translational symmetry in a QD reduces the ability of the electrons to delocalize. As a result, a Wigner-type cross-over is expected for a smaller value of

$r_s$ . (b) Mesoscopic fluctuations, inherent in any confined system, lead to a rich interplay with the correlation effects. These two added features make strong correlation physics particularly interesting in a QD. As clean 2D bulk samples with large  $r_s$  are regularly fabricated these days in semiconductor heterostructures, it seems to be just a matter of time before these systems are patterned into a QD, thus providing an excellent probe of correlation effects.

The above-mentioned quantum mechanical levels can, in turn, be tuned by means of, for example, the application of various external fields. The spins of the electrons in quantum dots provide a natural basis for representing so-called qubits [3]. The capability to manipulate and study such states is evidenced by several recent experiments [4, 5]. Coupled quantum dots are particularly interesting since so-called two-qubit quantum gates can be realized by manipulating the exchange coupling which originates from the repulsive Coulomb interaction and the underlying Pauli principle. For such states, the exchange coupling splits singlet and triplet states, and depending on the shape of the confining potential and the applied magnetic field, one can allow for electrical or magnetic control of the exchange coupling. In particular, several recent experiments and theoretical investigations have analyzed the role of effective spin-orbit interactions in quantum dots [6–9] and their influence on the exchange coupling.

A proper theoretical understanding of the exchange coupling, correlation energies, ground state energies of quantum dots, the role of spin-orbit interactions and other properties of quantum dots as well, requires the development of appropriate and reliable theoretical few- and many-body methods. Furthermore, for quantum dots with more than two electrons and/or specific values of the external fields, this implies the development of few- and many-body methods where uncertainty quantifications are provided. For most methods, this means providing an estimate of the error due to the truncation made in the single-particle basis and the truncation made in limiting the number of possible excitations. For systems with more than three or four electrons, *ab initio* methods that have been employed in studies of quantum dots are variational and diffusion Monte Carlo [10, 12, 13], path integral approaches [14], large-scale diagonalization (full configuration interaction) [15–17, 19], and to a very limited extent coupled-cluster theory [20–24]. Exact diagonalization studies are accurate for a very small number of electrons, but the number of basis functions needed to obtain a given accuracy and the computational cost grow very rapidly with electron number. In practice they have been used for up to eight electrons [15, 16, 19], but the accuracy is very limited for all except  $N \leq 3$ .

Monte Carlo methods have been applied up to  $N = 24$  electrons [12, 13]. Diffusion Monte Carlo, with statistical and systematic errors, provide, in principle, exact benchmark solutions to various properties of quantum dots. However, the computations start becoming rather time-consuming for larger systems. Hartree[25], restricted Hartree-Fock, spin- and/or space-unrestricted Hartree-Fock[26–28] and local spin-density, and current density functional methods[29–32] give results that are satisfactory for a qualitative understanding of some systematic properties. However, comparisons with exact results show discrepancies in the energies that are substantial on the scale of energy differences.

For systems with many electrons many, the dimensionalities of the systems render the usage of many of the above methods almost impossible. Density functional theory is now the computational paradigm for quantum mechanical calculations of energies and other properties of molecules and condensed matter. Furthermore, the time-dependent theory applies particularly to electronic excitations and interactions with the electromagnetic field, see for example Lecture Notes in Physics, volume 837.

### Specific tasks

The specific aim of this thesis is, based on an already developed Variational Monte Carlo code by the candidate, to develop a diffusion Monte Carlo program for quantum dots (a program which could be extended to other systems as well and to electrons confined in three dimensions) which will allow for exact benchmarks and reference for closed-shell systems with up to approximately  $\sim 50 - 70$  electrons.

Based on the exact diffusion Monte Carlo calculations, one can in turn extract a single-particle density. This single-particle density will in turn be used to determine a

density functional for the same closed-shell systems. The method to be developed will be based on the so-called adiabatic connection, see Refs. [33–35]. These functionals will in turn be compared with existing functionals for quantum dots under various trapping frequencies. If possible, the application of a real-time formalism for time-dependent density functional theory can be developed, see Ref. [36].

### Progress plan and milestones

The aims and progress plan of this thesis are as follows

- Fall 2012: Develop and finalize a diffusion Monte Carlo code for electrons trapped in a single harmonic oscillator in two dimensions. The code should be made flexible enough to be extended to trapped electrons or other trapped fermionic systems in two and three dimensions. An optimized and parallel version of the code should be tested for systems up to or larger than  $\sim 70$  electrons.
- Fall 2012 and first part of spring 2013: The obtained ground states (from the diffusion Monte Carlo code) will in turn be used to define a as exact as possible density functional for quantum dots using the adiabatic-connection method. The density functional can in turn be used to model systems with a large number of electrons in quantum dots. It will also be compared to standard functionals used for studies of quantum dots as done in Refs. [31, 32].
- Last part of spring 2013: write up of thesis and discussion of results.

The thesis is expected to be handed in June 1 2013.

- 
- [1] S. K. Bogner, R. J. Furnstahl, H. Hergert, M. Kortelainen, P. Maris, M. Stoitsov, and J. P. Vary, *Phys. Rev. C* **84**, 044306 (2011).
  - [2] Y. Alhassid, G. F. Bertsch, and L. Fang, *Phys. Rev. Lett.* **100**, 230401 (2008).
  - [3] D. Loss and D. P. DiVincenzo, *Phys. Rev. A* **57**, 120 (1998).
  - [4] R. Hanson, L. H. Willems van Beveren, I. T. Vink, J. M. Elzerman, W. J. M. Naber, F. H. L. Koppens, L. P. Kouwenhoven, and L. M. K. Vandersypen, *Phys. Rev. Lett.* **94**, 196802 (2005).
  - [5] H.-A. Engel, V. N. Golovach, D. Loss, L. M. K. Vandersypen, J. M. Elzerman, R. Hanson, and L. P. Kouwenhoven, *Phys. Rev. Lett.* **93**, 106804 (2004).
  - [6] C. Fasth, A. Fuhrer, L. Samuelson, V. N. Golovach, and D. Loss, *Phys. Rev. Lett.* **98**, 266801 (2007).
  - [7] S. Roddaro *et al.*, *Phys. Rev. Lett.* **101**, 18682 (2008).
  - [8] A. Ambrosetti, J. M. Escartin, E. Lipparini, and F. Ped-  
eriva, arXiv:1003.2433.
  - [9] F. Baruffa, P. Stano, and J. Fabian, *Phys. Rev. Lett.* **104**, 126401 (2010).
  - [10] A. Harju, *J. Low Temp. Phys.* **140**, 181 (2005).
  - [11] F. Bolton, *Phys. Rev. B* **54**, 4780 (1996).
  - [12] F. Pederiva, C. J. Umrigar, and E. Lipparini, *Phys. Rev. B* **62**, 8120 (2000).
  - [13] L. Colletti, F. Pederiva, and E. Lipparini, *Eur. Phys. J. B* **27**, 385 (2002).
  - [14] M. Harowitz, D. Shin, and J. Shumway, *J. Low. Temp. Phys.* **140**, 211 (2005).
  - [15] M. Eto, *Jpn. J. Appl. Phys.* **36**, 3924 (1997).
  - [16] P. A. Maksym and T. Chakraborty, *Phys. Rev. Lett.* **65**, 108 (1990); D. Pfannkuche, V. Gudmundsson, and P. A. Maksym, *Phys. Rev. B* **47**, 2244 (1993); P. Hawrylak and D. Pfannkuche, *Phys. Rev. Lett.* **70**, 485 (1993); J.J. Palacios, L. Moreno, G. Chiappe, E. Louis, and C. Tejedor, *Phys. Rev. B* **50**, 5760 (1994); T. Ezaki,

- N. Mori, and C. Hamaguchi, Phys. Rev. B **56**, 6428 (1997).
- [17] S. Kvaal, Phys. Rev. C **78**, 044330 (2008).
- [18] S. Kvaal, arXiv:0810.2644, unpublished.
- [19] M. Rontani, C. Cavazzoni, D. Belucci, and G. Goldoni, J. Chem. Phys. **124**, 124102 (2006).
- [20] I. Shavitt and R. J. Bartlett, *Many-body Methods in Chemistry and Physics*, (Cambridge University Press, Cambridge UK, 2009).
- [21] R. J. Bartlett and M. Musiał, Rev. Mod. Phys. **79**, 291 (2007).
- [22] T. M. Henderson, K. Runge, and R. J. Bartlett, Chem. Phys. Lett. **337**, 138 (2001); Phys. Rev. B **67**, 045320 (2003).
- [23] I. Heidari, S. Pal, B. S. Pujari, and D. G. Kanhere, J. Chem. Phys. **127**, 114708 (2007).
- [24] M. Pedersen Lohne, G. Hagen, M. Hjorth-Jensen, S. Kvaal, and F. Pederiva, Phys. Rev. B **84**, 115302 (2011).
- [25] A. Kumar, S. E. Laux, and F. Stern, Phys. Rev. B **42**, 5166 (1990).
- [26] M. Fujito, A. Natori, and H. Yasunaga, Phys. Rev. B **53**, 9952 (1996).
- [27] H. M. Muller and S. Koonin, Phys. Rev. B **54**, 14532 (1996).
- [28] C. Yannouleas and U. Landman, Phys. Rev. Lett. **82**, 5325 (1999).
- [29] M. Koskinen, M. Manninen, and S.M. Reimann, Phys. Rev. Lett. **79**, 1389 (1997).
- [30] K. Hirose and N. S. Wingreen, Phys. Rev. B **59**, 4604 (1999).
- [31] P. Gori-Giorgi, M. Seidl, and G. Vignale, Phys. Rev. Lett. **103**, 166402 (2009).
- [32] E. Räsänen, S. Pittalis, J. G. Vilhena, M. A. L. Marques, Int. J. Quantum Chem. **110**, 2308 (2010).
- [33] F. Colonna and A. Savin, J. Chem. Phys. **110**, 2828 (1999).
- [34] A. M. Teale, S. Coriani, and T. Helgaker, J. Chem. Phys. **130**, 104111 (2009).
- [35] Q. Wu and W. Yang, J. Chem. Phys. **118**, 2498 (2003).
- [36] K.-L. Lee, K. Yabana, G.F. Bertsch, J. Chem. Phys. **134**, 144106 (2011).