Name: Jisoo Hwang
GitHub Username: Jisoo2073
Purdue Username: hwang332
Instructor: Gregory David Sturm
Section: 005

Problem 1

1.

Given the probability density function:

$$f_X(x) = \begin{cases} \frac{3}{4}(1-x^2) \; when \; x \in [-1,1] \\ 0 \qquad\quad otherwise \end{cases}$$

If you happen to plot this distribution, then by looking at the plot, determine by inspection, what is the mean of the distribution.

Mean of the distribution:

0

From this distribution we grab 100 i.i.d samples $X_1, X_2, \ldots X_{100}$.

The sample mean random variable is defined as

$$\bar{X}_n = \frac{1}{n}\sum_{i=1}^{n} X_i$$

What is the shape of the probability density function of the sample mean?

Gaussian

What is the mean of the probability density function of the sample mean?

0

Explain how you know the shape and mean.

We grab 100 samples, this means, according to Central Limit Theorem, the distribution of the samples will be approximately normal distributed. That is why the shape will be Gaussian. And the mean will be 0 since over 100 sample's mean will be collected.

Part B

Let X be a normally distributed random variable with mean = 0 and variance = 1. What is the distribution of the random variable Z = 3X+1 (Give your answer with the shape, mean and variance).

What is the shape of the distribution?

Normal Distribution

What is the mean?

1

What is the variance?

1.732

X ~ N(0,1)

Z = 3(N(0,1)) + 1 -> the shape will not change.

Mean: u_f = A*u_i + B = 3 * 0 + 1

Deviation: sigma_f =sqrt(sigma^2 * A^2)

Problem 2

You are given data in *city_vehicle_survey.txt* representing the average age of vehicles across various counties. The transportation department claims the average age of vehicles is 5 years. You are tasked with testing this claim.

1. Formulate null and alternative hypotheses for a statistical test that seeks to challenge this belief. What are the null and alternative hypotheses?

   1. Null Hypothesis:

$\mu = 5$ years

   2. Alternative Hypothesis:

$\mu \neq 5$ years

   3. What type of test should be used and why?

t-test should be used because the population standard deviation is unknown.

2. Carry out this statistical test using the *city_vehicle_survey.txt* sample. Report the sample size, the sample mean, the standard error, the standard score (z or t, depending on what was used), and the p-value.

   ********ROUND ALL DECIMAL VALUES TO 4 DECIMAL PLACES******

Sample size	1024
Sample mean	5.1141
Standard error	0.0614
Standard score	1.8585
p – value (if less than 0.01 use scientific notation)	0.0634

   Are the results statistically significant at a level of 0.05? 0.0634 > 0.05

   Yes **No**

   What (if anything) can we conclude about the hypothesis at the confidence level of 0.05?

Because the p-value is greater than 0.05, we cannot reject the null hypothesis.

Are the results statistically significant at a level of 0.10? 0.0634 < 0.10

Yes **No**

What (if anything) can we conclude about the hypothesis at the confidence level of 0.10?

Because 0.0634 is smaller than 0.10, we can reject the null hypothesis.

3. What is the largest standard error for which the test will be significant at a level of 0.05? What is the corresponding minimum sample size? (You may assume that the population variance and mean does not change.)

********ROUND ALL DECIMAL VALUES TO 4 DECIMAL PLACES******

Largest standard error	0.0581
Corresponding minimum sample size	1141

4. Suppose the transportation department believes the mean vehicle age is the same in counties with and without emission control programs. Two datasets, *vehicle_data_1.txt* (with emission programs) and *vehicle_data_2.txt* (without emission programs), are used to test this assumption.

   1. Null Hypothesis:

mu_1 = mu_2

   2. Alternative Hypothesis:

mu_1 is not equal to mu_2

   3. What type of test should be used and why?

A two-sample z-test should be used because we are comparing the means of two samples.

5. Carry out this statistical test using the *vehicle_data_1.txt* population and *vehicle_data_2.txt* population samples. Report the sample sizes, the sample means, the standard error, the z-score, and the p-value. Are the results significant at levels 0.05 or 0.10? What (if anything) can we conclude about the hypothesis at the two different confidence levels?

********ROUND ALL DECIMAL VALUES TO 4 DECIMAL PLACES**** ******

Sample size of *vehicle_data_1* (Emission)	512
Sample size of *vehicle_data_2* (Without Emission)	868
Sample mean of *vehicle_data_1* (Emission)	5.5415
Sample mean of *vehicle_data_2* (Without Emission)	6.2482
Standard error	0.1043
Standard score	-6.7770
p – value (if less than 0.01 use scientific notation)	1.2269* 10 ^-11

1. Are the results statistically significant at a level of 0.05?

Yes **No**

2. Are the results statistically significant at a level of 0.10?

Yes **No**

3. What (if anything) can we conclude (i.e., what is the interpretation of the result)?

We reject the null hypothesis at both the 0.05 and 0.10 significance levels.

Problem 3

1. Use the sample to construct a 90% confidence interval for the average sodium of snacks. Report whether you will use a z-test or t-test and report the sample mean, the standard error, the standard statistic (t or z value), and the interval. (Think, which distribution should you use here if very few data points are available?)

t-distribution should be used because the sample size is small n = 20.

********ROUND ALL DECIMAL VALUES TO 4 DECIMAL PLACES******

Sample mean	140.45
Standard error	1.3426
Standard score (t or z value)	1.7291
90% confidence interval	

2. Repeat Q1 for a 95% confidence interval.

********ROUND ALL DECIMAL VALUES TO 4 DECIMAL PLACES******

Standard error	
Standard score (t or z value)	2.0930
95% confidence interval	(137.6400, 143.2600)

Is your interval wider or narrower compared to using the 90% confidence interval in Q1?

Wider **Narrower**

3. Repeat Q2 if you are told that the population standard deviation is 5.
Will you use a t-test or z-test (Hint: Think which distribution should you use here now that you have the true population standard deviation)? Justify your answer.

I should use z distribution because I have the population standard deviation.

********ROUND ALL DECIMAL VALUES TO 4 DECIMAL PLACES******

Standard error	1.1180
Standard score (t or z value)	1.9600
95% confidence interval	(138.2587, 142.6413)

Is your interval wider or narrower than the interval computed in Q2?

Wider **Narrower**