

Binomial Identity in Arbitrary Bases

Lin Jiu



Research Institute for Symbolic Computation
Johannes Kepler University Linz
@ Beijing Institute of Technology



JKU
JOHANNES KEPLER
UNIVERSITÄT LINZ

July 26th 2016

Binomial Identity

$$(X + Y)^n = \sum_{k=0}^n \binom{n}{k} X^k Y^{n-k}.$$

Generalization.

Multi-nomial Identity

$$(X_1 + \cdots + X_m)^n = \sum_{k_1 + \cdots + k_m = n} \binom{n}{k_1, \dots, k_m} X_1^{k_1} \cdots X_m^{k_m}.$$

Binomial Identity

$$(X + Y)^n = \sum_{k=0}^n \binom{n}{k} X^k Y^{n-k}.$$

Generalization.

Multi-nomial Identity

$$(X_1 + \cdots + X_m)^n = \sum_{k_1 + \cdots + k_m = n} \binom{n}{k_1, \dots, k_m} X_1^{k_1} \cdots X_m^{k_m}.$$

Binomial Identity

$$(X + Y)^n = \sum_{k=0}^n \binom{n}{k} X^k Y^{n-k}.$$

Generalization.

Multi-nomial Identity

$$(X_1 + \cdots + X_m)^n = \sum_{k_1 + \cdots + k_m = n} \binom{n}{k_1, \dots, k_m} X_1^{k_1} \cdots X_m^{k_m}.$$

Binary

| | | | | | | | | | |
|---------|---|---|----|----|-----|-----|-----|-----|-----|
| n | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | ... |
| $(n)_2$ | 0 | 1 | 10 | 11 | 100 | 101 | 110 | 111 | ... |

DEF.

$S_2(n) = \#$ of 1's in the binary expansion of n .

Example

$$\begin{cases} S_2(3) = 2 \\ S_2(4) = 1 \\ S_2(7) = 3 \end{cases}$$

| | | | | | | | | | |
|---------|---|---|----|----|-----|-----|-----|-----|-----|
| n | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | ... |
| $(n)_2$ | 0 | 1 | 10 | 11 | 100 | 101 | 110 | 111 | ... |

DEF.

$S_2(n) = \# \text{ of } 1\text{'s in the binary expansion of } n.$

Example

$$\begin{cases} S_2(3) = 2 \\ S_2(4) = 1 \\ S_2(7) = 3 \end{cases}$$

Binary

| | | | | | | | | | |
|---------|---|---|----|----|-----|-----|-----|-----|-----|
| n | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | ... |
| $(n)_2$ | 0 | 1 | 10 | 11 | 100 | 101 | 110 | 111 | ... |

DEF.

$S_2(n) = \#$ of 1's in the binary expansion of n .

Example

$$\begin{cases} S_2(3) = 2 \\ S_2(4) = 1 \\ S_2(7) = 3 \end{cases}$$

Binary

| | | | | | | | | | |
|---------|---|---|----|----|-----|-----|-----|-----|-----|
| n | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | ... |
| $(n)_2$ | 0 | 1 | 10 | 11 | 100 | 101 | 110 | 111 | ... |

DEF.

$S_2(n) = \#$ of 1's in the binary expansion of n .

Example

$$\begin{cases} S_2(3) = 2 \\ S_2(4) = 1 \\ S_2(7) = 3 \end{cases}$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{0 \leq k \leq n \text{ \& \& some condition}} X^{S_2(k)} Y^{S_2(n-k)},$$

some condition = there is no carry when adding $k + (n - k) = n$.

In fact,

$$k + (n - k) = n \text{ is carry free} \Leftrightarrow S_2(k) + S_2(n - k) = S_2(n).$$

Thus,

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{0 \leq k \leq n \text{ \& some condition}} X^{S_2(k)} Y^{S_2(n-k)},$$

some condition = there is no carry when adding $k + (n - k) = n$.

In fact,

$$k + (n - k) = n \text{ is carry free} \Leftrightarrow S_2(k) + S_2(n - k) = S_2(n).$$

Thus,

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{0 \leq k \leq n \text{ \& some condition}} X^{S_2(k)} Y^{S_2(n-k)},$$

some condition = there is no carry when adding $k + (n - k) = n$.

In fact,

$$k + (n - k) = n \text{ is carry free} \Leftrightarrow S_2(k) + S_2(n - k) = S_2(n).$$

Thus,

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{0 \leq k \leq n \text{ \& some condition}} X^{S_2(k)} Y^{S_2(n-k)},$$

some condition = there is no carry when adding $k + (n - k) = n$.

In fact,

$$k + (n - k) = n \text{ is carry free} \Leftrightarrow S_2(k) + S_2(n - k) = S_2(n).$$

Thus,

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{0 \leq k \leq n \text{ \& some condition}} X^{S_2(k)} Y^{S_2(n-k)},$$

some condition = there is no carry when adding $k + (n - k) = n$.

In fact,

$$k + (n - k) = n \text{ is carry free} \Leftrightarrow S_2(k) + S_2(n - k) = S_2(n).$$

Thus,

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

Example

Example

$n = 6 = (110)_2$. Therefore,

$$LHS = (X + Y)^{S_2(2)} = (X + Y)^2.$$

On the other hand,

| $k + (n - k)$ | $0 + 6$ | $1 + 5$ | $2 + 4$ | $3 + 3$ | $4 + 2$ | $5 + 1$ | $6 + 0$ |
|---------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | 000 | 001 | 010 | 011 | 100 | 101 | 110 |
| Binary | $\frac{110}{110}$ | $\frac{101}{110}$ | $\frac{100}{110}$ | $\frac{011}{110}$ | $\frac{010}{110}$ | $\frac{001}{110}$ | $\frac{000}{110}$ |
| Carry-free | ✓ | × | ✓ | × | ✓ | × | ✓ |

$$\begin{aligned} RHS &= X^{S_2(0)} Y^{S_2(6)} + X^{S_2(2)} Y^{S_2(4)} + X^{S_2(4)} Y^{S_2(2)} + X^{S_2(6)} Y^{S_2(0)} \\ &= Y^2 + XY + XY + X^2. \end{aligned}$$

Remark

Carry Free $\Leftrightarrow 1 + 1$ does not appear.

Example

Example

$n = 6 = (110)_2$. Therefore,

$$LHS = (X + Y)^{S_2(2)} = (X + Y)^2.$$

On the other hand,

| $k + (n - k)$ | 0 + 6 | 1 + 5 | 2 + 4 | 3 + 3 | 4 + 2 | 5 + 1 | 6 + 0 |
|---------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | 000 | 001 | 010 | 011 | 100 | 101 | 110 |
| Binary | $\frac{110}{110}$ | $\frac{101}{110}$ | $\frac{100}{110}$ | $\frac{011}{110}$ | $\frac{010}{110}$ | $\frac{001}{110}$ | $\frac{000}{110}$ |
| Carry-free | ✓ | × | ✓ | × | ✓ | × | ✓ |

$$\begin{aligned} RHS &= X^{S_2(0)} Y^{S_2(6)} + X^{S_2(2)} Y^{S_2(4)} + X^{S_2(4)} Y^{S_2(2)} + X^{S_2(6)} Y^{S_2(0)} \\ &= Y^2 + XY + XY + X^2. \end{aligned}$$

Remark

Carry Free $\Leftrightarrow 1 + 1$ does not appear.

Example

Example

$n = 6 = (110)_2$. Therefore,

$$LHS = (X + Y)^{S_2(2)} = (X + Y)^2.$$

On the other hand,

| $k + (n - k)$ | $0 + 6$ | $1 + 5$ | $2 + 4$ | $3 + 3$ | $4 + 2$ | $5 + 1$ | $6 + 0$ |
|---------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | 000 | 001 | 010 | 011 | 100 | 101 | 110 |
| Binary | $\frac{110}{110}$ | $\frac{101}{110}$ | $\frac{100}{110}$ | $\frac{011}{110}$ | $\frac{010}{110}$ | $\frac{001}{110}$ | $\frac{000}{110}$ |
| Carry-free | ✓ | × | ✓ | × | ✓ | × | ✓ |

$$\begin{aligned} RHS &= X^{S_2(0)} Y^{S_2(6)} + X^{S_2(2)} Y^{S_2(4)} + X^{S_2(4)} Y^{S_2(2)} + X^{S_2(6)} Y^{S_2(0)} \\ &= Y^2 + XY + XY + X^2. \end{aligned}$$

Remark

Carry Free $\Leftrightarrow 1 + 1$ does not appear.

Example

Example

$n = 6 = (110)_2$. Therefore,

$$LHS = (X + Y)^{S_2(2)} = (X + Y)^2.$$

On the other hand,

| $k + (n - k)$ | 0 + 6 | 1 + 5 | 2 + 4 | 3 + 3 | 4 + 2 | 5 + 1 | 6 + 0 |
|---------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| Binary | 000 | 001 | 010 | 011 | 100 | 101 | 110 |
| | $\frac{110}{110}$ | $\frac{101}{110}$ | $\frac{100}{110}$ | $\frac{011}{110}$ | $\frac{010}{110}$ | $\frac{001}{110}$ | $\frac{000}{110}$ |
| Carry-free | ✓ | × | ✓ | × | ✓ | × | ✓ |

$$\begin{aligned} RHS &= X^{S_2(0)} Y^{S_2(6)} + X^{S_2(2)} Y^{S_2(4)} + X^{S_2(4)} Y^{S_2(2)} + X^{S_2(6)} Y^{S_2(0)} \\ &= Y^2 + XY + XY + X^2. \end{aligned}$$

Remark

Carry Free $\Leftrightarrow 1 + 1$ does not appear.

Example

Example

$n = 6 = (110)_2$. Therefore,

$$LHS = (X + Y)^{S_2(2)} = (X + Y)^2.$$

On the other hand,

| $k + (n - k)$ | 0 + 6 | 1 + 5 | 2 + 4 | 3 + 3 | 4 + 2 | 5 + 1 | 6 + 0 |
|---------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | 000 | 001 | 010 | 011 | 100 | 101 | 110 |
| Binary | $\frac{110}{110}$ | $\frac{101}{110}$ | $\frac{100}{110}$ | $\frac{011}{110}$ | $\frac{010}{110}$ | $\frac{001}{110}$ | $\frac{000}{110}$ |
| Carry-free | ✓ | × | ✓ | × | ✓ | × | ✓ |

$$\begin{aligned} RHS &= X^{S_2(0)} Y^{S_2(6)} + X^{S_2(2)} Y^{S_2(4)} + X^{S_2(4)} Y^{S_2(2)} + X^{S_2(6)} Y^{S_2(0)} \\ &= Y^2 + XY + XY + X^2. \end{aligned}$$

Remark

Carry Free $\Leftrightarrow 1 + 1$ does not appear.

Questions

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{I}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{I}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{I}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{I}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{I}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{I}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{I}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{i}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{i}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

Theorem

$$(X + Y)^{S_2(n)} = \sum_{S_2(k) + S_2(n-k) = S_2(n)} X^{S_2(k)} Y^{S_2(n-k)}.$$

carry-free $\Leftrightarrow \binom{S_2(n)}{i}$, i.e., distributing 1's in $(n)_2$.

- How to generalize this result to other bases rather than 2, for example 3?
- How to define

$$S_3(n) = ???$$

of 1's? # of 2's? # of non-zero digits?

- In binary, $0 + 1$, $0 + 0$ are OK and $1 + 1$ is not. \Leftrightarrow Carry Free.
In base 3,

$$1 + 1 = 2?$$

DEF.

$S_2(n) = \#$ of 1's in the binary expansion of $n =$ sum of all digits.

Thus,

$S_b(n) =$ sum of all digits of n in its expansion of base b .

This also implies $1 + 1 = 2$ is allowed.

Unfortunately $(X + Y)^{S_3(n)} \neq \sum_{\text{carry-free}} X^{S_3(k)} Y^{S_3(n-k)}$

$$S_3(2) = S_3((2)_3) = 2.$$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$LHS = (X + Y)^2 \neq X^2 + XY + Y^2 = RHS.$$

Solutions

DEF.

$S_2(n) = \#$ of 1's in the binary expansion of $n =$ sum of all digits.

Thus,

$S_b(n) =$ sum of all digits of n in its expansion of base b .

This also implies $1 + 1 = 2$ is allowed.

Unfortunately $(X + Y)^{S_3(n)} \neq \sum_{\text{carry-free}} X^{S_3(k)} Y^{S_3(n-k)}$

$$S_3(2) = S_3((2)_3) = 2.$$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$LHS = (X + Y)^2 \neq X^2 + XY + Y^2 = RHS.$$

Solutions

DEF.

$S_2(n) = \#$ of 1's in the binary expansion of $n =$ sum of all digits.

Thus,

$S_b(n) =$ sum of all digits of n in its expansion of base b .

This also implies $1 + 1 = 2$ is allowed.

Unfortunately $(X + Y)^{S_3(n)} \neq \sum_{\text{carry-free}} X^{S_3(k)} Y^{S_3(n-k)}$

$$S_3(2) = S_3((2)_3) = 2.$$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$LHS = (X + Y)^2 \neq X^2 + XY + Y^2 = RHS.$$

Solutions

DEF.

$S_2(n) = \#$ of 1's in the binary expansion of $n =$ sum of all digits.

Thus,

$S_b(n) =$ sum of all digits of n in its expansion of base b .

This also implies $1 + 1 = 2$ is allowed.

Unfortunately $(X + Y)^{S_3(n)} \neq \sum_{\text{carry-free}} X^{S_3(k)} Y^{S_3(n-k)}$

$$S_3(2) = S_3((2)_3) = 2.$$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$LHS = (X + Y)^2 \neq X^2 + XY + Y^2 = RHS.$$

$S_2(n) = \#$ of 1's in the binary expansion of $n =$ sum of all digits.

Thus,

$S_b(n) =$ sum of all digits of n in its expansion of base b .

This also implies $1 + 1 = 2$ is allowed.

Unfortunately $(X + Y)^{S_3(n)} \neq \sum_{\text{carry-free}} X^{S_3(k)} Y^{S_3(n-k)}$

$$S_3(2) = S_3((2)_3) = 2.$$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$LHS = (X + Y)^2 \neq X^2 + XY + Y^2 = RHS.$$

Result

$$(X + Y)^{S_b(n)} = \sum_{\square_1} \square_2 X^{S_b(k)} Y^{S_b(n-k)}$$

$$\begin{cases} \square_1 = \text{carry-free} & \mapsto ? \\ \square_2 = 1 & \mapsto ? \end{cases}$$

Modify \square_2 :

$$(X + Y)^{S_b(n)} = \sum_{\text{carry-free}} \left(\prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} \right) X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$S_b^{(j)}(m) = \# \text{ of } j\text{'s in } m.$$

Result

$$(X + Y)^{S_b(n)} = \sum_{\square_1} \square_2 X^{S_b(k)} Y^{S_b(n-k)}$$

$$\begin{cases} \square_1 = \text{carry-free} & \mapsto ? \\ \square_2 = 1 & \mapsto ? \end{cases}$$

Modify \square_2 :

$$(X + Y)^{S_b(n)} = \sum_{\text{carry-free}} \left(\prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} \right) X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$S_b^{(j)}(m) = \# \text{ of } j\text{'s in } m.$$

Result

$$(X + Y)^{S_b(n)} = \sum_{\square_1} \square_2 X^{S_b(k)} Y^{S_b(n-k)}$$

$$\begin{cases} \square_1 = \text{carry-free} & \mapsto ? \\ \square_2 = 1 & \mapsto ? \end{cases}$$

Modify \square_2 :

$$(X + Y)^{S_b(n)} = \sum_{\text{carry-free}} \left(\prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} \right) X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$S_b^{(j)}(m) = \# \text{ of } j\text{'s in } m.$$

Result

$$(X + Y)^{S_b(n)} = \sum_{\square_1} \square_2 X^{S_b(k)} Y^{S_b(n-k)}$$

$$\begin{cases} \square_1 = \text{carry-free} & \mapsto ? \\ \square_2 = 1 & \mapsto ? \end{cases}$$

Modify \square_2 :

$$(X + Y)^{S_b(n)} = \sum_{\text{carry-free}} \left(\prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} \right) X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$S_b^{(j)}(m) = \# \text{ of } j\text{'s in } m.$$

Result

$$n = 2$$

$$S_3(2) = 2.$$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$\begin{cases} LHS = (X + Y)^2 \\ RHS = X^2 + Y^2 + (2!)^{1-0-0} XY \end{cases}$$

Next

Define the b -ary binomial coefficients:

$$\binom{n}{k}_b := \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{if carry-free} \\ 0 & \text{otherwise} \end{cases}$$

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)}.$$

Result

$$n = 2$$

$$S_3(2) = 2.$$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$\begin{cases} LHS = (X + Y)^2 \\ RHS = X^2 + Y^2 + (2!)^{1-0-0} XY \end{cases}$$

Next

Define the b -ary binomial coefficients:

$$\binom{n}{k}_b := \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{if carry-free} \\ 0 & \text{otherwise} \end{cases}$$

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)}.$$

Result

$n = 2$

$S_3(2) = 2.$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$\begin{cases} LHS = (X + Y)^2 \\ RHS = X^2 + Y^2 + (2!)^{1-0-0} XY \end{cases}$$

Next

Define the b -ary binomial coefficients:

$$\binom{n}{k}_b := \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{if carry-free} \\ 0 & \text{otherwise} \end{cases}$$

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)}.$$

Result

$$n = 2$$

$$S_3(2) = 2.$$

| $k + (n - k)$ | $0 + 2$ | $1 + 1$ | $2 + 0$ |
|---------------|---------------|---------------|---------------|
| | 0 | 1 | 2 |
| Binary | $\frac{2}{2}$ | $\frac{1}{2}$ | $\frac{0}{2}$ |
| Carry-free | ✓ | ✓ | ✓ |

$$\begin{cases} LHS = (X + Y)^2 \\ RHS = X^2 + Y^2 + (2!)^{1-0-0} XY \end{cases}$$

Next

Define the b -ary binomial coefficients:

$$\binom{n}{k}_b := \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{if carry-free} \\ 0 & \text{otherwise} \end{cases}$$

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)}.$$

b -ary Binomial Coefficients

$$b = 4$$


b -ary Binomial Coefficients

Triangle

$$b = 4$$

b -ary Binomial Coefficients

Triangle

$$b = 4$$


b -ary Binomial Coefficients

Generating Function

Define

$$f(n, b, x) := \sum_{k=0}^n \binom{n}{k}_b x^k,$$

for $b = 4$,

| n | $f(n, 4, x)$ |
|-----|----------------------|
| 1 | $1 + x$ |
| 2 | $(1 + x)^2$ |
| 3 | $(1 + x)^3$ |
| 4 | $1 + x^4$ |
| 5 | $(1 + x)(1 + x^4)$ |
| 6 | $(1 + x)^2(1 + x^4)$ |
| 7 | $(1 + x)^3(1 + x^4)$ |
| 8 | $(1 + x^4)^2$ |

b -ary Binomial Coefficients

Generating Function

Define

$$f(n, b, x) := \sum_{k=0}^n \binom{n}{k}_b x^k,$$

for $b = 4$,

| n | $f(n, 4, x)$ |
|-----|----------------------|
| 1 | $1 + x$ |
| 2 | $(1 + x)^2$ |
| 3 | $(1 + x)^3$ |
| 4 | $1 + x^4$ |
| 5 | $(1 + x)(1 + x^4)$ |
| 6 | $(1 + x)^2(1 + x^4)$ |
| 7 | $(1 + x)^3(1 + x^4)$ |
| 8 | $(1 + x^4)^2$ |

b -ary Binomial Coefficients

Generating Function

Define

$$f(n, b, x) := \sum_{k=0}^n \binom{n}{k}_b x^k,$$

for $b = 4$,

| n | $f(n, 4, x)$ |
|-----|----------------------|
| 1 | $1 + x$ |
| 2 | $(1 + x)^2$ |
| 3 | $(1 + x)^3$ |
| 4 | $1 + x^4$ |
| 5 | $(1 + x)(1 + x^4)$ |
| 6 | $(1 + x)^2(1 + x^4)$ |
| 7 | $(1 + x)^3(1 + x^4)$ |
| 8 | $(1 + x^4)^2$ |

b -ary Binomial Coefficients

Generating Function

Define

$$f(n, b, x) := \sum_{k=0}^n \binom{n}{k}_b x^k,$$

for $b = 4$,

| n | $f(n, 4, x)$ |
|-----|----------------------|
| 1 | $1 + x$ |
| 2 | $(1 + x)^2$ |
| 3 | $(1 + x)^3$ |
| 4 | $1 + x^4$ |
| 5 | $(1 + x)(1 + x^4)$ |
| 6 | $(1 + x)^2(1 + x^4)$ |
| 7 | $(1 + x)^3(1 + x^4)$ |
| 8 | $(1 + x^4)^2$ |

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$\binom{n}{k}_b = \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{carry-free} \\ 0 & \text{otherwise} \end{cases} = \prod_{l=0}^{N-1} \binom{n_l}{k_l}.$$

In base b ,

$$n = n_{N-1}n_{N-2} \cdots n_0 \text{ and } k = k_{N-1}k_{N-2} \cdots k_0$$

i.e.,

$$n = n_{N-1}b^{N-1} + n_{N-2}b^{N-2} + \cdots + n_0b^0 \text{ and } k = \sum_{j=0}^{N-1} k_j b^j.$$

$$\text{carry-free} \Leftrightarrow k_i \leq n_i$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$\binom{n}{k}_b = \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{carry-free} \\ 0 & \text{otherwise} \end{cases} = \prod_{l=0}^{N-1} \binom{n_l}{k_l}.$$

In base b ,

$$n = n_{N-1}n_{N-2} \cdots n_0 \text{ and } k = k_{N-1}k_{N-2} \cdots k_0$$

i.e.,

$$n = n_{N-1}b^{N-1} + n_{N-2}b^{N-2} + \cdots + n_0b^0 \text{ and } k = \sum_{j=0}^{N-1} k_jb^j.$$

$$\text{carry-free} \Leftrightarrow k_i \leq n_i$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$\binom{n}{k}_b = \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{carry-free} \\ 0 & \text{otherwise} \end{cases} = \prod_{l=0}^{N-1} \binom{n_l}{k_l}.$$

In base b ,

$$n = n_{N-1}n_{N-2} \cdots n_0 \text{ and } k = k_{N-1}k_{N-2} \cdots k_0$$

i.e.,

$$n = n_{N-1}b^{N-1} + n_{N-2}b^{N-2} + \cdots + n_0b^0 \text{ and } k = \sum_{j=0}^{N-1} k_jb^j.$$

$$\text{carry-free} \Leftrightarrow k_i \leq n_i$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$\binom{n}{k}_b = \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{carry-free} \\ 0 & \text{otherwise} \end{cases} = \prod_{l=0}^{N-1} \binom{n_l}{k_l}.$$

In base b ,

$$n = n_{N-1}n_{N-2} \cdots n_0 \text{ and } k = k_{N-1}k_{N-2} \cdots k_0$$

i.e.,

$$n = n_{N-1}b^{N-1} + n_{N-2}b^{N-2} + \cdots + n_0b^0 \text{ and } k = \sum_{j=0}^{N-1} k_jb^j.$$

$$\text{carry-free} \Leftrightarrow k_i \leq n_i$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$\binom{n}{k}_b = \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{carry-free} \\ 0 & \text{otherwise} \end{cases} = \prod_{l=0}^{N-1} \binom{n_l}{k_l}.$$

In base b ,

$$n = n_{N-1}n_{N-2} \cdots n_0 \text{ and } k = k_{N-1}k_{N-2} \cdots k_0$$

i.e.,

$$n = n_{N-1}b^{N-1} + n_{N-2}b^{N-2} + \cdots + n_0b^0 \text{ and } k = \sum_{j=0}^{N-1} k_j b^j.$$

$$\text{carry-free} \Leftrightarrow k_i \leq n_i$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)},$$

where

$$\binom{n}{k}_b = \begin{cases} \prod_{j=0}^{b-1} (j!)^{S_b^{(j)}(n) - S_b^{(j)}(k) - S_b^{(j)}(n-k)} & \text{carry-free} \\ 0 & \text{otherwise} \end{cases} = \prod_{l=0}^{N-1} \binom{n_l}{k_l}.$$

In base b ,

$$n = n_{N-1}n_{N-2} \cdots n_0 \text{ and } k = k_{N-1}k_{N-2} \cdots k_0$$

i.e.,

$$n = n_{N-1}b^{N-1} + n_{N-2}b^{N-2} + \cdots + n_0b^0 \text{ and } k = \sum_{j=0}^{N-1} k_jb^j.$$

$$\text{carry-free} \Leftrightarrow k_i \leq n_i$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)}, \quad \binom{n}{k}_b = \prod_{l=0}^{N-1} \binom{n_l}{k_l}$$

Implies:

$$\sum_{k=0}^n \binom{n}{k}_b x^k = \prod_{l=0}^{N-1} (1 + x^{b^l})^{n_l}.$$

Remark[Lucas Theorem]

For a prime p ,

$$\binom{n}{k}_p \equiv \prod_{l=0}^{N-1} \binom{n_l}{k_l} \pmod{p}.$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)}, \quad \binom{n}{k}_b = \prod_{l=0}^{N-1} \binom{n_l}{k_l}$$

Implies:

$$\sum_{k=0}^n \binom{n}{k}_b x^k = \prod_{l=0}^{N-1} (1 + x^{b^l})^{n_l}.$$

Remark[Lucas Theorem]

For a prime p ,

$$\binom{n}{k}_p \equiv \prod_{l=0}^{N-1} \binom{n_l}{k_l} \pmod{p}.$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)}, \quad \binom{n}{k}_b = \prod_{l=0}^{N-1} \binom{n_l}{k_l}$$

Implies:

$$\sum_{k=0}^n \binom{n}{k}_b x^k = \prod_{l=0}^{N-1} (1 + x^{b^l})^{n_l}.$$

Remark[Lucas Theorem]

For a prime p ,

$$\binom{n}{k}_p \equiv \prod_{l=0}^{N-1} \binom{n_l}{k_l} \pmod{p}.$$

Theorem

$$(X + Y)^{S_b(n)} = \sum_{k=0}^n \binom{n}{k}_b X^{S_b(k)} Y^{S_b(n-k)}, \quad \binom{n}{k}_b = \prod_{l=0}^{N-1} \binom{n_l}{k_l}$$

Implies:

$$\sum_{k=0}^n \binom{n}{k}_b x^k = \prod_{l=0}^{N-1} (1 + x^{b^l})^{n_l}.$$

Remark[Lucas Theorem]

For a prime p ,

$$\binom{n}{k}_p \equiv \prod_{l=0}^{N-1} \binom{n_l}{k_l} \pmod{p}.$$

Thank You!