

The Method of Brackets (MoB) and Integrating by Differentiating (IbD) Method

Lin Jiu

Research Institute for Symbolic Computation
Johannes Kepler University
Dec. 9th 2016



RISC
Software GmbH



Acknowledgement

Joint Work with:



V. H. Moll



K. Kohl



I. Gonzalez

IMAGE
NOT
FOUND

C. Vignat

Outlines

1 The method of brackets (MoB)

- Rules
- Ramanujan's Master Theorem (RMT)
- Examples
- Recent result

2 Integration by Differentiating

- Formulas
- Recent proofs
- Connection

Rules

Idea

MoB evaluates the definite integral

$$\int_0^{\infty} f(x) dx$$

(most of the time) in terms of **SERIES**, with *ONLY SIX* rules:

Defintion [**Indicator**]

$$\phi_n := \frac{(-1)^n}{n!} = \frac{(-1)^n}{\Gamma(n+1)}$$

and

$$\phi_{1,\dots,r} := \phi_{n_1,\dots,n_r} = \phi_{n_1} \phi_{n_2} \cdots \phi_{n_r} = \prod_{i=1}^r \phi_{n_i}.$$

Rules

Idea

MoB evaluates the definite integral

$$\int_0^{\infty} f(x) dx$$

(most of the time) in terms of **SERIES**, with **ONLY SIX** rules:

Definition [Indicator]

$$\phi_n := \frac{(-1)^n}{n!} = \frac{(-1)^n}{\Gamma(n+1)}$$

and

$$\phi_{1,\dots,r} := \phi_{n_1,\dots,n_r} = \phi_{n_1} \phi_{n_2} \cdots \phi_{n_r} = \prod_{i=1}^r \phi_{n_i}.$$

Rules

Idea

MoB evaluates the definite integral

$$\int_0^{\infty} f(x) dx$$

(most of the time) in terms of **SERIES**, with **ONLY SIX** rules:

Definition [**Indicator**]

$$\phi_n := \frac{(-1)^n}{n!} = \frac{(-1)^n}{\Gamma(n+1)}$$

and

$$\phi_{1,\dots,r} := \phi_{n_1,\dots,n_r} = \phi_{n_1} \phi_{n_2} \cdots \phi_{n_r} = \prod_{i=1}^r \phi_{n_i}.$$

Rules

Idea

MoB evaluates the definite integral

$$\int_0^{\infty} f(x) dx$$

(most of the time) in terms of **SERIES**, with *ONLY SIX* rules:

Definition [**Indicator**]

$$\phi_n := \frac{(-1)^n}{n!} = \frac{(-1)^n}{\Gamma(n+1)}$$

and

$$\phi_{1,\dots,r} := \phi_{n_1,\dots,n_r} = \phi_{n_1} \phi_{n_2} \cdots \phi_{n_r} = \prod_{i=1}^r \phi_{n_i}.$$

Rules

Idea

MoB evaluates the definite integral

$$\int_0^{\infty} f(x) dx$$

(most of the time) in terms of **SERIES**, with *ONLY SIX* rules:

Definition [**Indicator**]

$$\phi_n := \frac{(-1)^n}{n!} = \frac{(-1)^n}{\Gamma(n+1)}$$

and

$$\phi_{1,\dots,r} := \phi_{n_1,\dots,n_r} = \phi_{n_1} \phi_{n_2} \cdots \phi_{n_r} = \prod_{i=1}^r \phi_{n_i}.$$

Rules (P-Production; E-Evaluation) $I = \int_0^\infty f(x) dx$

$$P_1: f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^\infty f(x) dx \mapsto \sum_n a_n \langle \alpha n + \beta \rangle \text{---Bracket Series;}$$

$$P_2: \text{For } \alpha < 0, (a_1 + \dots + a_r)^\alpha \mapsto \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \dots a_r^{n_r} \frac{\langle -\alpha + n_1 + \dots + n_r \rangle}{\Gamma(-\alpha)};$$

P_3 : For each bracket series, we assign index = # of sums - # of brackets;

$$E_1: \sum_n \phi_n f(n) \langle \alpha n + \beta \rangle = \frac{1}{|\alpha|} f(n^*) \Gamma(-n^*), \text{ where } n^* \text{ solves } \alpha n + \beta = 0;$$

$$E_2: \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} f(n_1, \dots, n_r) \prod_{i=1}^r \langle a_{i1} n_1 + \dots + a_{ir} n_r + c_i \rangle = \frac{f(n_1^*, \dots, n_r^*) \prod_{i=1}^r \Gamma(-n_i^*)}{|\det A|},$$

$$(n_1^*, \dots, n_r^*) \text{ solves } \begin{cases} a_{11} n_1 + \dots + a_{1r} n_r + c_1 = 0 \\ \dots & \dots \\ a_{r1} n_1 + \dots + a_{rr} n_r + c_r = 0 \end{cases}$$

E_3 : The value of a multi-dimensional bracket series of **POSITIVE** index is obtained by computing all the contributions of maximal rank by Rule E_2 . These contributions to the integral appear as series in the free parameters. Series converging in a common region are added and divergent series are discarded. Any series producing a non-real contribution is also discarded.

Ramanujan's Master Theorem[RMT]

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ a(0) - \frac{a(1)}{1!}x + \frac{a(2)}{2!}x^2 - \dots \right\} dx = a(-s) \Gamma(s)$$

(1)

$$\int_0^{\infty} x^{s-1} \left(\sum_{n=0}^{\infty} \phi_n a(n) x^n \right) dx = a(-s) \Gamma(s)$$

(2) [Hardy]

- $H(\delta) := \{s = \sigma + it : \sigma \geq -\delta, 0 < \delta < 1\}$;
- $\psi(x) \in C^\infty(H(\delta))$; $\exists C, P, A, A < \pi$ such that $|\psi(s)| \leq Ce^{P\delta + A|t|}$, $\forall s \in H(\delta)$;
- $0 < c < \delta$, $\Psi(x) := \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\pi}{\sin(\pi s)} \psi(-s) x^{-s} ds \stackrel{0 < x < e^{-P}}{=} \sum_{k=0}^{\infty} \psi(k) (-x)^k$;

$$\int_0^{\infty} \Psi(x) x^{s-1} dx = \frac{\pi}{\sin(\pi s)} \psi(-s).$$

Ramanujan's Master Theorem[RMT]

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ a(0) - \frac{a(1)}{1!}x + \frac{a(2)}{2!}x^2 - \dots \right\} dx = a(-s) \Gamma(s)$$

(1)

$$\int_0^{\infty} x^{s-1} \left(\sum_{n=0}^{\infty} \phi_n a(n) x^n \right) dx = a(-s) \Gamma(s)$$

(2) [Hardy]

- $H(\delta) := \{s = \sigma + it : \sigma \geq -\delta, 0 < \delta < 1\}$;
- $\psi(x) \in C^{\infty}(H(\delta))$; $\exists C, P, A, A < \pi$ such that $|\psi(s)| \leq Ce^{P\delta + A|t|}$, $\forall s \in H(\delta)$;
- $0 < c < \delta$, $\Psi(x) := \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\pi}{\sin(\pi s)} \psi(-s) x^{-s} ds \stackrel{0 \leq x < e^{-P}}{=} \sum_{k=0}^{\infty} \psi(k) (-x)^k$;

$$\int_0^{\infty} \Psi(x) x^{s-1} dx = \frac{\pi}{\sin(\pi s)} \psi(-s).$$

Ramanujan's Master Theorem[RMT]

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ a(0) - \frac{a(1)}{1!}x + \frac{a(2)}{2!}x^2 - \dots \right\} dx = a(-s) \Gamma(s)$$

(1)

$$\int_0^{\infty} x^{s-1} \left(\sum_{n=0}^{\infty} \phi_n a(n) x^n \right) dx = a(-s) \Gamma(s)$$

(2) [Hardy]

- $H(\delta) := \{s = \sigma + it : \sigma \geq -\delta, 0 < \delta < 1\}$;
- $\psi(x) \in C^{\infty}(H(\delta))$; $\exists C, P, A, A < \pi$ such that $|\psi(s)| \leq Ce^{P\delta + A|t|}$, $\forall s \in H(\delta)$;
- $0 < c < \delta$, $\Psi(x) := \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\pi}{\sin(\pi s)} \psi(-s) x^{-s} ds \stackrel{0 \leq x < e^{-P}}{=} \sum_{k=0}^{\infty} \psi(k) (-x)^k$;

$$\int_0^{\infty} \Psi(x) x^{s-1} dx = \frac{\pi}{\sin(\pi s)} \psi(-s).$$

Ramanujan's Master Theorem[RMT]

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ a(0) - \frac{a(1)}{1!}x + \frac{a(2)}{2!}x^2 - \dots \right\} dx = a(-s) \Gamma(s)$$

(1)

$$\int_0^{\infty} x^{s-1} \left(\sum_{n=0}^{\infty} \phi_n a(n) x^n \right) dx = a(-s) \Gamma(s)$$

(2) [Hardy]

- $H(\delta) := \{s = \sigma + it : \sigma \geq -\delta, 0 < \delta < 1\}$;
- $\psi(x) \in C^{\infty}(H(\delta))$; $\exists C, P, A, A < \pi$ such that $|\psi(s)| \leq Ce^{P\delta + A|t|}$, $\forall s \in H(\delta)$;
- $0 < c < \delta$, $\Psi(x) := \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\pi}{\sin(\pi s)} \psi(-s) x^{-s} ds \stackrel{0 \leq x < e^{-P}}{=} \sum_{k=0}^{\infty} \psi(k) (-x)^k$;

$$\int_0^{\infty} \Psi(x) x^{s-1} dx = \frac{\pi}{\sin(\pi s)} \psi(-s).$$

Ramanujan's Master Theorem[RMT]

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ a(0) - \frac{a(1)}{1!}x + \frac{a(2)}{2!}x^2 - \dots \right\} dx = a(-s) \Gamma(s)$$

(1)

$$\int_0^{\infty} x^{s-1} \left(\sum_{n=0}^{\infty} \phi_n a(n) x^n \right) dx = a(-s) \Gamma(s)$$

(2) [Hardy]

- $H(\delta) := \{s = \sigma + it : \sigma \geq -\delta, 0 < \delta < 1\}$;
- $\psi(x) \in C^\infty(H(\delta))$; $\exists C, P, A, A < \pi$ such that $|\psi(s)| \leq Ce^{P\delta + A|t|}$, $\forall s \in H(\delta)$;
- $0 < c < \delta$, $\Psi(x) := \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\pi}{\sin(\pi s)} \psi(-s) x^{-s} ds \stackrel{0 \leq x < e^{-P}}{=} \sum_{k=0}^{\infty} \psi(k) (-x)^k$;

$$\int_0^{\infty} \Psi(x) x^{s-1} dx = \frac{\pi}{\sin(\pi s)} \psi(-s).$$

Rules Again

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ \sum_{n=0}^{\infty} \phi_n a(n) x^n \right\} dx = a(-s) \Gamma(s)$$

(1) Integrand \mapsto Power Series;

(2) Keep Track of s ;

(3) Apply the Formula;

(4) Multiple Integrals;

$$\int_0^{\infty} \int_0^{\infty} \sum_{n,m} a(m,n) x^m y^n dx dy = ?$$

(5) More Sums than Integrals (brackets);

$$\int_0^{\infty} f_1(x) f_2(x) dx = \int_0^{\infty} \sum_{m,n} a(m,n) x^{m+n} dx = \sum_{m,n} a(m,n) (m+n+1) = ?$$

(6) Extra

Rules Again

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ \sum_{n=0}^{\infty} \phi_n a(n) x^n \right\} dx = a(-s) \Gamma(s)$$

(1) Integrand \mapsto Power Series;

(2) Keep Track of s ;

(3) Apply the Formula;

(4) Multiple Integrals;

$$\int_0^{\infty} \int_0^{\infty} \sum_{n,m} a(m,n) x^m y^n dx dy = ?$$

(5) More Sums than Integrals (brackets);

$$\int_0^{\infty} f_1(x) f_2(x) dx = \int_0^{\infty} \sum_{m,n} a(m,n) x^{m+n} dx = \sum_{m,n} a(m,n) (m+n+1) = ?$$

(6) Extra

Rules Again

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ \sum_{n=0}^{\infty} \phi_n a(n) x^n \right\} dx = a(-s) \Gamma(s)$$

- (1) Integrand \mapsto Power Series;
- (2) Keep Track of s ;
- (3) Apply the Formula;
- (4) Multiple Integrals;

$$\int_0^{\infty} \int_0^{\infty} \sum_{n,m} a(m,n) x^m y^n dx dy = ?$$

- (5) More Sums than Integrals (brackets);

$$\int_0^{\infty} f_1(x) f_2(x) dx = \int_0^{\infty} \sum_{m,n} a(m,n) x^{m+n} dx = \sum_{m,n} a(m,n) \langle m+n+1 \rangle = ?$$

- (6) Extra.

Rules Again

$$P_1: f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx \mapsto \sum_n a_n \langle \alpha n + \beta \rangle \boxed{s-1 \mapsto s}$$

$$P_2: \text{For } \alpha < 0, (a_1 + \dots + a_r)^{\alpha} \mapsto \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \dots a_r^{n_r} \frac{\langle -\alpha + n_1 + \dots + n_r \rangle}{\Gamma(-\alpha)};$$

$$P_3: \text{Index} = \# \text{ of sums} - \# \text{ of brackets}; \boxed{\text{Just a definition}}$$

$$E_1: \sum_n \phi_n f(n) \langle \alpha n + \beta \rangle = \frac{f(n^*) \Gamma(-n^*)}{|\alpha|}, \text{, } n^* \text{ solves } \alpha n + \beta = 0; \boxed{\text{RMT}}$$

$$E_2: \boxed{\text{Iteration of RMT}}$$

$$\sum_{n_1, \dots, n_r} \phi_{1, \dots, r} f(n_1, \dots, n_r) \prod_{i=1}^r \langle a_{i1} n_1 + \dots + a_{ir} n_r + c_i \rangle = \frac{f(n_1^*, \dots, n_r^*) \prod_{i=1}^r \Gamma(-n_i^*)}{|\det A|}$$

E_3 : The value of a multi-dimensional bracket series of **POSITIVE** index is obtained by computing all the contributions of maximal rank by Rule E_2 . These contributions to the integral appear as series in the free parameters. Series converging in a common region are added and divergent series are discarded. Any series producing a non-real contribution is also discarded.

Rules Again

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ \sum_{n=0}^{\infty} \phi_n a(n) x^n \right\} dx = a(-s) \Gamma(s)$$

- (1) Integrand \mapsto Power Series;
- (2) Keep Track of s ;
- (3) Apply the Formula;
- (4) Multiple Integrals;

$$\int_0^{\infty} \int_0^{\infty} \sum_{n,m} a(m,n) x^m y^n dx dy = ?$$

- (5) More Sums than Integrals (brackets);

$$\int_0^{\infty} f_1(x) f_2(x) dx = \int_0^{\infty} \sum_{m,n} a(m,n) x^{m+n} dx = \sum_{m,n} a(m,n) \langle m+n+1 \rangle = ?$$

- (6) Extra.

Rules Again

$$P_1: f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx \mapsto \sum_n a_n \langle \alpha n + \beta \rangle \boxed{s-1 \mapsto s}$$

$$P_2: \text{For } \alpha < 0, (a_1 + \dots + a_r)^{\alpha} \mapsto \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \dots a_r^{n_r} \frac{\langle -\alpha + n_1 + \dots + n_r \rangle}{\Gamma(-\alpha)};$$

$$P_3: \text{Index} = \# \text{ of sums} - \# \text{ of brackets}; \boxed{\text{Just a definition}}$$

$$E_1: \sum_n \phi_n f(n) \langle \alpha n + \beta \rangle = \frac{f(n^*) \Gamma(-n^*)}{|\alpha|}, n^* \text{ solves } \alpha n + \beta = 0; \boxed{\text{RMT}}$$

$$E_2: \boxed{\text{Iteration of RMT}}$$

$$\sum_{n_1, \dots, n_r} \phi_{1, \dots, r} f(n_1, \dots, n_r) \prod_{i=1}^r \langle a_{i1} n_1 + \dots + a_{ir} n_r + c_i \rangle = \frac{f(n_1^*, \dots, n_r^*) \prod_{i=1}^r \Gamma(-n_i^*)}{|\det A|}$$

E_3 : The value of a multi-dimensional bracket series of **POSITIVE** index is obtained by computing all the contributions of maximal rank by Rule E_2 . These contributions to the integral appear as series in the free parameters. Series converging in a common region are added and divergent series are discarded. Any series producing a non-real contribution is also discarded.

Rules Again

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ \sum_{n=0}^{\infty} \phi_n a(n) x^n \right\} dx = a(-s) \Gamma(s)$$

- (1) Integrand \mapsto Power Series;
- (2) Keep Track of s ;
- (3) Apply the Formula;
- (4) Multiple Integrals;

$$\int_0^{\infty} \int_0^{\infty} \sum_{n,m} a(m,n) x^m y^n dx dy = ?$$

- (5) More Sums than Integrals (brackets);

$$\int_0^{\infty} f_1(x) f_2(x) dx = \int_0^{\infty} \sum_{m,n} a(m,n) x^{m+n} dx = \sum_{m,n} a(m,n) \langle m+n+1 \rangle = ?$$

- (6) Extra.

Rules Again

$$P_1: f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx \mapsto \sum_n a_n \langle \alpha n + \beta \rangle \boxed{s-1 \mapsto s}$$

$$P_2: \text{For } \alpha < 0, (a_1 + \dots + a_r)^{\alpha} \mapsto \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \dots a_r^{n_r} \frac{\langle -\alpha + n_1 + \dots + n_r \rangle}{\Gamma(-\alpha)};$$

$$P_3: \text{Index} = \# \text{ of sums} - \# \text{ of brackets; } \boxed{\text{Just a definition}}$$

$$E_1: \sum_n \phi_n f(n) \langle \alpha n + \beta \rangle = \frac{f(n^*) \Gamma(-n^*)}{|\alpha|}, n^* \text{ solves } \alpha n + \beta = 0; \boxed{\text{RMT}}$$

$$E_2: \boxed{\text{Iteration of RMT}}$$

$$\sum_{n_1, \dots, n_r} \phi_{1, \dots, r} f(n_1, \dots, n_r) \prod_{i=1}^r \langle a_{i1} n_1 + \dots + a_{ir} n_r + c_i \rangle = \frac{f(n_1^*, \dots, n_r^*) \prod_{i=1}^r \Gamma(-n_i^*)}{|\det A|}$$

E_3 : The value of a multi-dimensional bracket series of **POSITIVE** index is obtained by computing all the contributions of maximal rank by Rule E_2 . These contributions to the integral appear as series in the free parameters. Series converging in a common region are added and divergent series are discarded. Any series producing a non-real contribution is also discarded.

Rules Again

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ \sum_{n=0}^{\infty} \phi_n a(n) x^n \right\} dx = a(-s) \Gamma(s)$$

- (1) Integrand \mapsto Power Series;
- (2) Keep Track of s ;
- (3) Apply the Formula;
- (4) Multiple Integrals;

$$\int_0^{\infty} \int_0^{\infty} \sum_{n,m} a(m,n) x^m y^n dx dy = ?$$

- (5) More Sums than Integrals (brackets);

$$\int_0^{\infty} f_1(x) f_2(x) dx = \int_0^{\infty} \sum_{m,n} a(m,n) x^{m+n} dx = \sum_{m,n} a(m,n) \langle m+n+1 \rangle = ?$$

- (6) Extra.

Rules Again

$$P_1: f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx \mapsto \sum_n a_n \langle \alpha n + \beta \rangle \boxed{s-1 \mapsto s}$$

$$P_2: \text{For } \alpha < 0, (a_1 + \dots + a_r)^{\alpha} \mapsto \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \dots a_r^{n_r} \frac{\langle -\alpha + n_1 + \dots + n_r \rangle}{\Gamma(-\alpha)};$$

$$P_3: \text{Index} = \# \text{ of sums} - \# \text{ of brackets; } \boxed{\text{Just a definition}}$$

$$E_1: \sum_n \phi_n f(n) \langle \alpha n + \beta \rangle = \frac{f(n^*) \Gamma(-n^*)}{|\alpha|}, n^* \text{ solves } \alpha n + \beta = 0; \boxed{\text{RMT}}$$

$$E_2: \boxed{\text{Iteration of RMT}}$$

$$\sum_{n_1, \dots, n_r} \phi_{1, \dots, r} f(n_1, \dots, n_r) \prod_{i=1}^r \langle a_{i1} n_1 + \dots + a_{ir} n_r + c_i \rangle = \frac{f(n_1^*, \dots, n_r^*) \prod_{i=1}^r \Gamma(-n_i^*)}{|\det A|}$$

E_3 : The value of a multi-dimensional bracket series of **POSITIVE** index is obtained by computing all the contributions of maximal rank by Rule E_2 . These contributions to the integral appear as series in the free parameters. Series converging in a common region are added and divergent series are discarded. Any series producing a non-real contribution is also discarded.

Rules Again

$$P_1: f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx \mapsto \sum_n a_n \langle \alpha n + \beta \rangle \boxed{s-1 \mapsto s}$$

$$P_2: \text{For } \alpha < 0, (a_1 + \dots + a_r)^{\alpha} \mapsto \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \dots a_r^{n_r} \frac{\langle -\alpha + n_1 + \dots + n_r \rangle}{\Gamma(-\alpha)};$$

$$P_3: \text{Index} = \# \text{ of sums} - \# \text{ of brackets; } \boxed{\text{Just a definition}}$$

$$E_1: \sum_n \phi_n f(n) \langle \alpha n + \beta \rangle = \frac{f(n^*) \Gamma(-n^*)}{|\alpha|}, n^* \text{ solves } \alpha n + \beta = 0; \boxed{\text{RMT}}$$

$$E_2: \boxed{\text{Iteration of RMT}}$$

$$\sum_{n_1, \dots, n_r} \phi_{1, \dots, r} f(n_1, \dots, n_r) \prod_{i=1}^r \langle a_{i1} n_1 + \dots + a_{ir} n_r + c_i \rangle = \frac{f(n_1^*, \dots, n_r^*) \prod_{i=1}^r \Gamma(-n_i^*)}{|\det A|}$$

E_3 : The value of a multi-dimensional bracket series of **POSITIVE** index is obtained by computing all the contributions of maximal rank by Rule E_2 . These contributions to the integral appear as series in the free parameters. Series converging in a common region are added and divergent series are discarded. Any series producing a non-real contribution is also discarded.

Rules Again

Theorem[RMT]

$$\int_0^{\infty} x^{s-1} \left\{ \sum_{n=0}^{\infty} \phi_n a(n) x^n \right\} dx = a(-s) \Gamma(s)$$

- (1) Integrand \mapsto Power Series;
- (2) Keep Track of s ;
- (3) Apply the Formula;
- (4) Multiple Integrals;

$$\int_0^{\infty} \int_0^{\infty} \sum_{n,m} a(m,n) x^m y^n dx dy = ?$$

- (5) More Sums than Integrals (brackets);

$$\int_0^{\infty} f_1(x) f_2(x) dx = \int_0^{\infty} \sum_{m,n} a(m,n) x^{m+n} dx = \sum_{m,n} a(m,n) \langle m+n+1 \rangle = ?$$

- (6) Extra.

Rules Again

$$P_1: f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx \mapsto \sum_n a_n \langle \alpha n + \beta \rangle \boxed{s-1 \mapsto s}$$

$$P_2: \text{For } \alpha < 0, (a_1 + \dots + a_r)^{\alpha} \mapsto \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \dots a_r^{n_r} \frac{\langle -\alpha + n_1 + \dots + n_r \rangle}{\Gamma(-\alpha)};$$

$$P_3: \text{Index} = \# \text{ of sums} - \# \text{ of brackets; } \boxed{\text{Just a definition}}$$

$$E_1: \sum_n \phi_n f(n) \langle \alpha n + \beta \rangle = \frac{f(n^*) \Gamma(-n^*)}{|\alpha|}, n^* \text{ solves } \alpha n + \beta = 0; \boxed{\text{RMT}}$$

$$E_2: \boxed{\text{Iteration of RMT}}$$

$$\sum_{n_1, \dots, n_r} \phi_{1, \dots, r} f(n_1, \dots, n_r) \prod_{i=1}^r \langle a_{i1} n_1 + \dots + a_{ir} n_r + c_i \rangle = \frac{f(n_1^*, \dots, n_r^*) \prod_{i=1}^r \Gamma(-n_i^*)}{|\det A|}$$

E_3 : The value of a multi-dimensional bracket series of **POSITIVE** index is obtained by computing all the contributions of maximal rank by Rule E_2 . These contributions to the integral appear as series in the free parameters. Series converging in a common region are added and divergent series are discarded. Any series producing a non-real contribution is also discarded.

Rule P_2

$$\begin{aligned}
 & \frac{\Gamma(-\alpha)}{(a_1 + \cdots + a_r)^{-\alpha}} \\
 = & \int_0^\infty x^{-\alpha-1} e^{-(a_1 + \cdots + a_r)x} dx \\
 = & \int_0^\infty x^{-\alpha-1} e^{-a_1 x} e^{-a_2 x} \cdots e^{-a_r x} dx \\
 = & \int_0^\infty x^{-\alpha-1} \prod_{i=1}^r \left(\sum_{n_i=0}^{\infty} \phi_{n_i} (ax)^{n_i} \right) dx \\
 = & \int_0^\infty \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \cdots a_r^{n_r} x^{n_1 + \cdots + n_r - \alpha - 1} dx \\
 = & \sum_{n_1, \dots, n_r} \phi_{1, \dots, r} a_1^{n_1} \cdots a_r^{n_r} \langle -\alpha + n_1 + \cdots + n_r \rangle
 \end{aligned}$$

Examples

$$I := \int_0^{\infty} x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay} \quad [y > 0 \text{ } Re(a) > 0]$$

Rule P_2 :

$$\frac{1}{\sqrt{a^2 + x^2}} = (a^2 + x^2)^{-\frac{1}{2}} = \sum_{n_1, n_2} \phi_{1,2} a^{2n_1} x^{2n_2} \frac{\langle \frac{1}{2} + n_1 + n_2 \rangle}{\Gamma(\frac{1}{2})}$$

$J_0(xy)$

$$J_0(xy) = \sum_{n_3} \phi_{n_3} \frac{y^{2n_3}}{\Gamma(n_3 + 1) 2^{2n_3}} x^{2n_3}$$

Rule P_1

$$\begin{aligned} I &= \int_0^{\infty} \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle x^{2n_2 + 2n_3 + 1} dx \\ &= \sum \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle \langle 2n_2 + 2n_3 + 2 \rangle \end{aligned}$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay} \quad [y > 0 \text{ } Re(a) > 0]$$

Rule P_2 :

$$\frac{1}{\sqrt{a^2 + x^2}} = (a^2 + x^2)^{-\frac{1}{2}} = \sum_{n_1, n_2} \phi_{1,2} a^{2n_1} x^{2n_2} \frac{\langle \frac{1}{2} + n_1 + n_2 \rangle}{\Gamma(\frac{1}{2})}$$

$J_0(xy)$

$$J_0(xy) = \sum_{n_3} \phi_{n_3} \frac{y^{2n_3}}{\Gamma(n_3 + 1) 2^{2n_3}} x^{2n_3}$$

Rule P_1

$$\begin{aligned} I &= \int_0^\infty \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle x^{2n_2 + 2n_3 + 1} dx \\ &= \sum \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle \langle 2n_2 + 2n_3 + 2 \rangle \end{aligned}$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay} \quad [y > 0 \text{ } \text{Re}(a) > 0]$$

Rule P_2 :

$$\frac{1}{\sqrt{a^2 + x^2}} = (a^2 + x^2)^{-\frac{1}{2}} = \sum_{n_1, n_2} \phi_{1,2} a^{2n_1} x^{2n_2} \frac{\langle \frac{1}{2} + n_1 + n_2 \rangle}{\Gamma(\frac{1}{2})}$$

$J_0(xy)$

$$J_0(xy) = \sum_{n_3} \phi_{n_3} \frac{y^{2n_3}}{\Gamma(n_3 + 1) 2^{2n_3}} x^{2n_3}$$

Rule P_1

$$\begin{aligned} I &= \int_0^\infty \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle x^{2n_2 + 2n_3 + 1} dx \\ &= \sum \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle \langle 2n_2 + 2n_3 + 2 \rangle \end{aligned}$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay} \quad [y > 0 \text{ } \text{Re}(a) > 0]$$

Rule P_2 :

$$\frac{1}{\sqrt{a^2 + x^2}} = (a^2 + x^2)^{-\frac{1}{2}} = \sum_{n_1, n_2} \phi_{1,2} a^{2n_1} x^{2n_2} \frac{\langle \frac{1}{2} + n_1 + n_2 \rangle}{\Gamma(\frac{1}{2})}$$

$J_0(xy)$

$$J_0(xy) = \sum_{n_3} \phi_{n_3} \frac{y^{2n_3}}{\Gamma(n_3 + 1) 2^{2n_3}} x^{2n_3}$$

Rule P_1

$$\begin{aligned} I &= \int_0^\infty \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle x^{2n_2 + 2n_3 + 1} dx \\ &= \sum \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle \langle 2n_2 + 2n_3 + 2 \rangle \end{aligned}$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay} \quad [y > 0 \text{ } Re(a) > 0]$$

Rule P_2 :

$$\frac{1}{\sqrt{a^2 + x^2}} = (a^2 + x^2)^{-\frac{1}{2}} = \sum_{n_1, n_2} \phi_{1,2} a^{2n_1} x^{2n_2} \frac{\langle \frac{1}{2} + n_1 + n_2 \rangle}{\Gamma(\frac{1}{2})}$$

$J_0(xy)$

$$J_0(xy) = \sum_{n_3} \phi_{n_3} \frac{y^{2n_3}}{\Gamma(n_3 + 1) 2^{2n_3}} x^{2n_3}$$

Rule P_1

$$\begin{aligned} I &= \int_0^\infty \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle x^{2n_2 + 2n_3 + 1} dx \\ &= \sum \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma(\frac{1}{2}) 2^{2n_3}} \langle n_1 + n_2 + \frac{1}{2} \rangle \langle 2n_2 + 2n_3 + 2 \rangle \end{aligned}$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

n_1 free: $n_2^* = -\frac{1}{2} - n_1$; $n_3^* = -\frac{1}{2} + n_1$; $\det = 2$:

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free} : I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free} : I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$$E_3 : \quad I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

$$n_1 \text{ free: } n_2^* = -\frac{1}{2} - n_1; n_3^* = -\frac{1}{2} + n_1; \det = 2:$$

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free : } I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free : } I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$$E_3 : \quad I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Examples

$$I := \int_0^{\infty} x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

$$n_1 \text{ free: } n_2^* = -\frac{1}{2} - n_1; n_3^* = -\frac{1}{2} + n_1; \det = 2;$$

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free : } I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free : } I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$$E_3 : \quad I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

n_1 free: $n_2^* = -\frac{1}{2} - n_1$; $n_3^* = -\frac{1}{2} + n_1$; $\det = 2$:

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free} : I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free} : I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$$E_3 : \quad I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

$$n_1 \text{ free: } n_2^* = -\frac{1}{2} - n_1; n_3^* = -\frac{1}{2} + n_1; \det = 2:$$

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free : } I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free : } I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$$E_3 : \quad I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

$$n_1 \text{ free: } n_2^* = -\frac{1}{2} - n_1; n_3^* = -\frac{1}{2} + n_1; \det = 2:$$

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free : } I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free : } I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$E_3 :$

$$I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

$$n_1 \text{ free: } n_2^* = -\frac{1}{2} - n_1; n_3^* = -\frac{1}{2} + n_1; \det = 2:$$

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free : } I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free : } I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$E_3 :$

$$I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

$$n_1 \text{ free: } n_2^* = -\frac{1}{2} - n_1; n_3^* = -\frac{1}{2} + n_1; \det = 2:$$

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free : } I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free : } I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$E_3 :$

$$I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Examples

$$I := \int_0^\infty x J_0(xy) \frac{dx}{\sqrt{a^2 + x^2}} = y^{-1} e^{-ay}$$

$$I = \sum_{n_1, n_2, n_3} \phi_{1,2,3} \frac{y^{2n_3} a^{2n_1}}{\Gamma(n_3 + 1) \Gamma\left(\frac{1}{2}\right) 2^{2n_3}} \left\langle n_1 + n_2 + \frac{1}{2} \right\rangle \langle 2n_2 + 2n_3 + 2 \rangle ;$$

$$n_1 \text{ free: } n_2^* = -\frac{1}{2} - n_1; n_3^* = -\frac{1}{2} + n_1; \det = 2:$$

$$\begin{aligned} I &= \frac{1}{2} \sum_{n_1} \phi_{n_1} \frac{y^{2n_1-1} a^{2n_1}}{\Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) 2^{2n_1-1}} \Gamma\left(n_1 + \frac{1}{2}\right) \Gamma\left(-n_1 + \frac{1}{2}\right) \\ &= \frac{1}{y} \sum_{n_1=0}^{\infty} \phi_{n_1} \left(\frac{ay}{2}\right)^{2n_1} \frac{\Gamma\left(\frac{1}{2} - n_1\right)}{\Gamma\left(\frac{1}{2}\right)} = \frac{1}{y} \cosh(ay); \end{aligned}$$

$$n_2 \text{ free : } I = \frac{1}{\sqrt{\pi y}} \sum_{n_2=0}^{\infty} \frac{\Gamma\left(n_2 + \frac{1}{2}\right)}{\Gamma(-n_2)} \left(\frac{2}{ay}\right)^{2n_2+1} = 0; \quad n_3 \text{ free : } I = \text{Series} = -\frac{\sinh(ay)}{y};$$

$$E_3 : \quad I = \frac{1}{y} \cosh(ay) - \frac{\sinh(ay)}{y} = y^{-1} e^{-ay}.$$

Multi-dim

$$I = \int_{\mathbb{R}^m} f(x_1^2 + \cdots + x_m^2) dx_1 \cdots dx_m$$

(I) Usual method: By the n -dim spherical coordinate that $r = x_1^2 + \cdots + x_m^2$ and

$$\begin{cases} x_1 = r \cos(\phi_1), & 0 \leq \phi_1 \leq \pi, \\ x_2 = r \sin(\phi_1) \cos(\phi_2), & 0 \leq \phi_2 \leq \pi, \\ \dots & \dots \\ x_{n-2} = r \sin(\phi_1) \cdots \sin(\phi_{m-3}) \cos(\phi_{m-2}), & 0 \leq \phi_{m-2} \leq \pi, \\ x_{n-1} = r \sin(\phi_1) \cdots \sin(\phi_{m-2}) \cos(\phi_{m-1}), & 0 \leq \phi_{m-1} \leq 2\pi, \\ x_n = r \sin(\phi_1) \cdots \sin(\phi_{m-2}) \sin(\phi_{m-1}), & 0 \leq r < \infty, \end{cases}$$

we have

$$dx_1 \cdots dx_m = r^{m-1} \sin^{m-2}(\phi_1) \cdots \sin(\phi_{m-2}) dr d\phi_1 \cdots d\phi_{m-1}.$$

Thus,

$$I = 2\pi^{\frac{m}{2}} \left[\int_0^\infty r^{m-1} f(r^2) dr \right] \frac{1}{\Gamma(\frac{m}{2})}.$$

Multi-dim

$$I = \int_{\mathbb{R}^m} f(x_1^2 + \cdots + x_m^2) dx_1 \cdots dx_m$$

(I) Usual method: By the n -dim spherical coordinate that $r = x_1^2 + \cdots + x_m^2$ and

$$\begin{cases} x_1 = r \cos(\phi_1), & 0 \leq \phi_1 \leq \pi, \\ x_2 = r \sin(\phi_1) \cos(\phi_2), & 0 \leq \phi_2 \leq \pi, \\ \dots & \dots \\ x_{n-2} = r \sin(\phi_1) \cdots \sin(\phi_{m-3}) \cos(\phi_{m-2}), & 0 \leq \phi_{m-2} \leq \pi, \\ x_{n-1} = r \sin(\phi_1) \cdots \sin(\phi_{m-2}) \cos(\phi_{m-1}), & 0 \leq \phi_{m-1} \leq 2\pi, \\ x_n = r \sin(\phi_1) \cdots \sin(\phi_{m-2}) \sin(\phi_{m-1}), & 0 \leq r < \infty, \end{cases}$$

we have

$$dx_1 \cdots dx_m = r^{m-1} \sin^{m-2}(\phi_1) \cdots \sin(\phi_{m-2}) dr d\phi_1 \cdots d\phi_{m-1}.$$

Thus,

$$I = 2\pi^{\frac{m}{2}} \left[\int_0^\infty r^{m-1} f(r^2) dr \right] \frac{1}{\Gamma\left(\frac{m}{2}\right)}.$$

Multi-dim

$$I = \int_{\mathbb{R}^m} f(x_1^2 + \cdots + x_m^2) dx_1 \cdots dx_m$$

(I) Usual method: By the n -dim spherical coordinate that $r = x_1^2 + \cdots + x_m^2$ and

$$\begin{cases} x_1 = r \cos(\phi_1), & 0 \leq \phi_1 \leq \pi, \\ x_2 = r \sin(\phi_1) \cos(\phi_2), & 0 \leq \phi_2 \leq \pi, \\ \dots & \dots \\ x_{n-2} = r \sin(\phi_1) \cdots \sin(\phi_{m-3}) \cos(\phi_{m-2}), & 0 \leq \phi_{m-2} \leq \pi, \\ x_{n-1} = r \sin(\phi_1) \cdots \sin(\phi_{m-2}) \cos(\phi_{m-1}), & 0 \leq \phi_{m-1} \leq 2\pi, \\ x_n = r \sin(\phi_1) \cdots \sin(\phi_{m-2}) \sin(\phi_{m-1}), & 0 \leq r < \infty, \end{cases}$$

we have

$$dx_1 \cdots dx_m = r^{m-1} \sin^{m-2}(\phi_1) \cdots \sin(\phi_{m-2}) dr d\phi_1 \cdots d\phi_{m-1}.$$

Thus,

$$I = 2\pi^{\frac{m}{2}} \left[\int_0^\infty r^{m-1} f(r^2) dr \right] \frac{1}{\Gamma\left(\frac{m}{2}\right)}.$$

Multi-dim

$$I = \int_{\mathbb{R}^m} f(x_1^2 + \cdots + x_m^2) dx_1 \cdots dx_m$$

(I) Usual method: By the n -dim spherical coordinate that $r = x_1^2 + \cdots + x_m^2$ and

$$\begin{cases} x_1 = r \cos(\phi_1), & 0 \leq \phi_1 \leq \pi, \\ x_2 = r \sin(\phi_1) \cos(\phi_2), & 0 \leq \phi_2 \leq \pi, \\ \dots & \dots \\ x_{n-2} = r \sin(\phi_1) \cdots \sin(\phi_{m-3}) \cos(\phi_{m-2}), & 0 \leq \phi_{m-2} \leq \pi, \\ x_{n-1} = r \sin(\phi_1) \cdots \sin(\phi_{m-2}) \cos(\phi_{m-1}), & 0 \leq \phi_{m-1} \leq 2\pi, \\ x_n = r \sin(\phi_1) \cdots \sin(\phi_{m-2}) \sin(\phi_{m-1}), & 0 \leq r < \infty, \end{cases}$$

we have

$$dx_1 \cdots dx_m = r^{m-1} \sin^{m-2}(\phi_1) \cdots \sin(\phi_{m-2}) dr d\phi_1 \cdots d\phi_{m-1}.$$

Thus,

$$I = 2\pi^{\frac{m}{2}} \left[\int_0^\infty r^{m-1} f(r^2) dr \right] \frac{1}{\Gamma\left(\frac{m}{2}\right)}.$$

Multi-dim

$$I = \int_{\mathbb{R}^m} f(x_1^2 + \cdots + x_m^2) dx_1 \cdots dx_m$$

(II) The method of brackets: Suppose

$$f(t) = \sum_{l=0}^{\infty} \phi_l a(l) t^l,$$

then,

$$\int_0^{\infty} r^{m-1} f(r^2) dr = \sum_l \phi_l a(l) \langle 2l + m \rangle = \sum_l \phi_l a(l) \langle 2l + m \rangle = \frac{1}{2} a\left(-\frac{m}{2}\right) \Gamma\left(\frac{m}{2}\right).$$

So it suffices to show that

$$I = 2\pi^{\frac{m}{2}} \left[\frac{1}{2} a\left(-\frac{m}{2}\right) \frac{\Gamma\left(-\frac{m}{2} + 1\right)}{(-1)^{-\frac{m}{2}}} \Gamma\left(\frac{m}{2}\right) \right] \frac{1}{\Gamma\left(\frac{m}{2}\right)} = \pi^{\frac{m}{2}} a\left(-\frac{m}{2}\right).$$

Multi-dim

$$I = \int_{\mathbb{R}^m} f(x_1^2 + \cdots + x_m^2) dx_1 \cdots dx_m$$

(II) The method of brackets: Suppose

$$f(t) = \sum_{l=0}^{\infty} \phi_l a(l) t^l,$$

then,

$$\int_0^{\infty} r^{m-1} f(r^2) dr = \sum_l \phi_l a(l) \langle 2l + m \rangle = \sum_l \phi_l a(l) \langle 2l + m \rangle = \frac{1}{2} a\left(-\frac{m}{2}\right) \Gamma\left(\frac{m}{2}\right).$$

So it suffices to show that

$$I = 2\pi^{\frac{m}{2}} \left[\frac{1}{2} a\left(-\frac{m}{2}\right) \frac{\Gamma\left(-\frac{m}{2} + 1\right)}{(-1)^{-\frac{m}{2}}} \Gamma\left(\frac{m}{2}\right) \right] \frac{1}{\Gamma\left(\frac{m}{2}\right)} = \pi^{\frac{m}{2}} a\left(-\frac{m}{2}\right).$$

Multi-dim

Direct computation shows:

$$\begin{aligned}
 I &= 2^m \int_{\mathbb{R}_+^m} \left[\sum_{l=0}^{\infty} \phi_l a(l) (x_1^2 + \cdots + x_m^2)^l \right] dx_1 \cdots dx_m \\
 &= 2^m \int_{\mathbb{R}_+^m} \sum_{l=0}^{\infty} \phi_l a(l) \sum_{\substack{n_1, \dots, n_m \\ n_1 + \cdots + n_m = l}} \binom{l}{n_1, \dots, n_m} x_1^{2n_1} \cdots x_m^{2n_m} dx_1 \cdots dx_m \\
 &= 2^m \sum_{l=n_1 + \cdots + n_m} \phi_l a(l) \sum_{n_1, \dots, n_m} \phi_{1, \dots, m} \binom{l}{n_1, \dots, n_m} \frac{1}{\phi_{1, \dots, m}} \prod_{j=1}^m \langle 2n_j + 1 \rangle \\
 &= AC \dots \\
 &= \pi^{\frac{m}{2}} a\left(-\frac{m}{2}\right),
 \end{aligned}$$

as desired.

Multi-dim

Direct computation shows:

$$\begin{aligned}
 I &= 2^m \int_{\mathbb{R}_+^m} \left[\sum_{l=0}^{\infty} \phi_l a(l) (x_1^2 + \cdots + x_m^2)^l \right] dx_1 \cdots dx_m \\
 &= 2^m \int_{\mathbb{R}_+^m} \sum_{l=0}^{\infty} \phi_l a(l) \sum_{\substack{n_1, \dots, n_m \\ n_1 + \cdots + n_m = l}} \binom{l}{n_1, \dots, n_m} x_1^{2n_1} \cdots x_m^{2n_m} dx_1 \cdots dx_m \\
 &= 2^m \sum_{l=n_1 + \cdots + n_m} \phi_l a(l) \sum_{n_1, \dots, n_m} \phi_{1, \dots, m} \binom{l}{n_1, \dots, n_m} \frac{1}{\phi_{1, \dots, m}} \prod_{j=1}^m \langle 2n_j + 1 \rangle \\
 &= AC \dots \\
 &= \pi^{\frac{m}{2}} a\left(-\frac{m}{2}\right),
 \end{aligned}$$

as desired.

Multi-dim

Direct computation shows:

$$\begin{aligned}
 I &= 2^m \int_{\mathbb{R}_+^m} \left[\sum_{l=0}^{\infty} \phi_l a(l) (x_1^2 + \cdots + x_m^2)^l \right] dx_1 \cdots dx_m \\
 &= 2^m \int_{\mathbb{R}_+^m} \sum_{l=0}^{\infty} \phi_l a(l) \sum_{\substack{n_1, \dots, n_m \\ n_1 + \cdots + n_m = l}} \binom{l}{n_1, \dots, n_m} x_1^{2n_1} \cdots x_m^{2n_m} dx_1 \cdots dx_m \\
 &= 2^m \sum_{l=n_1 + \cdots + n_m} \phi_l a(l) \sum_{n_1, \dots, n_m} \phi_{1, \dots, m} \binom{l}{n_1, \dots, n_m} \frac{1}{\phi_{1, \dots, m}} \prod_{j=1}^m \langle 2n_j + 1 \rangle \\
 &= AC \dots \\
 &= \pi^{\frac{m}{2}} a\left(-\frac{m}{2}\right),
 \end{aligned}$$

as desired.

Multi-dim

$$I = \int_{\mathbb{R}_+^m} \frac{x_1^{p_1-1} \cdots x_m^{p_m-1} dx_1 \cdots dx_m}{(r_0 + r_1 x_1 + \cdots + r_m x_m)^s} = \frac{\Gamma(p_1) \cdots \Gamma(p_m) \Gamma(s - p_1 - p_2 - \cdots - p_m)}{r_1^{p_1} \cdots r_m^{p_m} r_0^{s-p_1-\cdots-p_m} \Gamma(s)}$$

$$(r_0 + r_1 x_1 + \cdots + r_m x_m)^{-s} = \sum_{n_0, n_1, \dots, n_m} \phi_{0,1,\dots,m} r_0^{n_0} r_1^{n_1} x_1^{n_1} \cdots r_m^{n_m} x_m^{n_m} \frac{\langle s + n_0 + \cdots + n_m \rangle}{\Gamma(s)}$$

$$I = \frac{1}{\Gamma(s)} \sum_{n_0, n_1, \dots, n_m} \phi_{0,1,\dots,m} r_0^{n_0} \cdots r_m^{n_m} \langle s + n_0 + \cdots + n_m \rangle \prod_{j=1}^m \langle n_m + p_m \rangle.$$

$$\begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 0 & 1 & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} n_0 \\ n_1 \\ \cdots \\ n_m \end{bmatrix} + \begin{bmatrix} s \\ p_1 \\ \cdots \\ p_m \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \cdots \\ 0 \end{bmatrix},$$

$$\det A = 1, \quad n_j^* = -p_j, \forall j = 1, \dots, m \quad \text{and} \quad n_0^* = p_1 + \cdots + p_m - s.$$

Multi-dim

$$I = \int_{\mathbb{R}_+^m} \frac{x_1^{p_1-1} \cdots x_m^{p_m-1} dx_1 \cdots dx_m}{(r_0 + r_1 x_1 + \cdots + r_m x_m)^s} = \frac{\Gamma(p_1) \cdots \Gamma(p_m) \Gamma(s - p_1 - p_2 - \cdots - p_m)}{r_1^{p_1} \cdots r_m^{p_m} r_0^{s-p_1-\cdots-p_m} \Gamma(s)}$$

$$(r_0 + r_1 x_1 + \cdots + r_m x_m)^{-s} = \sum_{n_0, n_1, \dots, n_m} \phi_{0,1,\dots,m} r_0^{n_0} r_1^{n_1} x_1^{n_1} \cdots r_m^{n_m} x_m^{n_m} \frac{\langle s + n_0 + \cdots + n_m \rangle}{\Gamma(s)}$$

$$I = \frac{1}{\Gamma(s)} \sum_{n_0, n_1, \dots, n_m} \phi_{0,1,\dots,m} r_0^{n_0} \cdots r_m^{n_m} \langle s + n_0 + \cdots + n_m \rangle \prod_{j=1}^m \langle n_m + p_m \rangle.$$

$$\begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 0 & 1 & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} n_0 \\ n_1 \\ \cdots \\ n_m \end{bmatrix} + \begin{bmatrix} s \\ p_1 \\ \cdots \\ p_m \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \cdots \\ 0 \end{bmatrix},$$

$$\det A = 1, \quad n_j^* = -p_j, \forall j = 1, \dots, m \quad \text{and} \quad n_0^* = p_1 + \cdots + p_m - s.$$

Null/Divergent Series

$$K_0(x) = \int_0^\infty \frac{\cos(tx) dt}{\sqrt{1+t^2}}.$$

$$K_0(x) = \frac{1}{2} \sum_n \phi_n \Gamma(-n) \frac{x^{2n}}{4^n} \text{ and } K_0(x) = \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2}{\Gamma(-n)} \cdot \frac{4^n}{x^{2n+1}}$$

$$\mathcal{M}(K_0)(s) = \int_0^\infty x^{s-1} K_0(x) dx = 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2$$

$$\begin{aligned} & \int_0^\infty x^{s-1} K_0(x) dx \\ &= \int_0^\infty x^{s-1} \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2}{\Gamma(-n)} \cdot \frac{4^n}{x^{2n+1}} dx \\ &= \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2 4^n}{\Gamma(-n)} \langle s - 2n - 1 \rangle \\ &= 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2, \end{aligned}$$

$$\begin{aligned} & \int_0^\infty x^{s-1} K_0(x) dx \\ &= \int_0^\infty \frac{x^{s-1}}{2} \sum_n \phi_n \Gamma(-n) \frac{x^{2n}}{4^n} dx \\ &= \frac{1}{2} \sum_n \phi_n \frac{\Gamma(-n)}{4^n} \langle 2n + s \rangle \\ &= 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2. \end{aligned}$$

Null/Divergent Series

$$K_0(x) = \int_0^\infty \frac{\cos(tx) dt}{\sqrt{1+t^2}}.$$

$$K_0(x) = \frac{1}{2} \sum_n \phi_n \Gamma(-n) \frac{x^{2n}}{4^n} \text{ and } K_0(x) = \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2}{\Gamma(-n)} \cdot \frac{4^n}{x^{2n+1}}$$

$$\mathcal{M}(K_0)(s) = \int_0^\infty x^{s-1} K_0(x) dx = 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2$$

$$\begin{aligned} & \int_0^\infty x^{s-1} K_0(x) dx \\ &= \int_0^\infty x^{s-1} \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2}{\Gamma(-n)} \cdot \frac{4^n}{x^{2n+1}} dx \\ &= \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2 4^n}{\Gamma(-n)} \langle s - 2n - 1 \rangle \\ &= 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2, \end{aligned}$$

$$\begin{aligned} & \int_0^\infty x^{s-1} K_0(x) dx \\ &= \int_0^\infty \frac{x^{s-1}}{2} \sum_n \phi_n \Gamma(-n) \frac{x^{2n}}{4^n} dx \\ &= \frac{1}{2} \sum_n \phi_n \frac{\Gamma(-n)}{4^n} \langle 2n + s \rangle \\ &= 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2. \end{aligned}$$

Null/Divergent Series

$$K_0(x) = \int_0^\infty \frac{\cos(tx) dt}{\sqrt{1+t^2}}.$$

$$K_0(x) = \frac{1}{2} \sum_n \phi_n \Gamma(-n) \frac{x^{2n}}{4^n} \text{ and } K_0(x) = \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2}{\Gamma(-n)} \cdot \frac{4^n}{x^{2n+1}}$$

$$\mathcal{M}(K_0)(s) = \int_0^\infty x^{s-1} K_0(x) dx = 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2$$

$$\begin{aligned} & \int_0^\infty x^{s-1} K_0(x) dx \\ &= \int_0^\infty x^{s-1} \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2}{\Gamma(-n)} \cdot \frac{4^n}{x^{2n+1}} dx \\ &= \sum_n \phi_n \frac{\Gamma\left(n + \frac{1}{2}\right)^2 4^n}{\Gamma(-n)} \langle s - 2n - 1 \rangle \\ &= 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2, \end{aligned}$$

$$\begin{aligned} & \int_0^\infty x^{s-1} K_0(x) dx \\ &= \int_0^\infty \frac{x^{s-1}}{2} \sum_n \phi_n \Gamma(-n) \frac{x^{2n}}{4^n} dx \\ &= \frac{1}{2} \sum_n \phi_n \frac{\Gamma(-n)}{4^n} \langle 2n + s \rangle \\ &= 2^{s-2} \Gamma\left(\frac{s}{2}\right)^2. \end{aligned}$$

DEF

A. Kempf et al. study Dirac-delta function to obtain the following formulas:

$$\begin{aligned}\int_a^b f(x) dx &= \lim_{\varepsilon \rightarrow 0} f(\partial_\varepsilon) \frac{e^{\varepsilon b} - e^{\varepsilon a}}{\varepsilon}, \\ \int_{-\infty}^{\infty} f(x) dx &= \lim_{\varepsilon \rightarrow 0} 2\pi f(-\iota \partial_\varepsilon) \delta(\varepsilon) = 2\pi \delta(\iota \partial_\varepsilon) f(\varepsilon), \\ \int_0^{\infty} f(x) dx &= \lim_{\varepsilon \rightarrow 0} f(-\partial_\varepsilon) \frac{1}{\varepsilon}, \\ \int_{-\infty}^0 f(x) dx &= \lim_{\varepsilon \rightarrow 0} f(\partial_\varepsilon) \frac{1}{\varepsilon}, \\ \int_{-\infty}^{\infty} f(x) dx &= \lim_{\varepsilon \rightarrow 0} [f(-\partial_\varepsilon) + f(\partial_\varepsilon)] \frac{1}{\varepsilon},\end{aligned}$$

where ∂_ε denotes the derivative with respect to ε .

DEF

A. Kempf et al. study Dirac-delta function to obtain the following formulas:

$$\begin{aligned}\int_a^b f(x) dx &= \lim_{\varepsilon \rightarrow 0} f(\partial_\varepsilon) \frac{e^{\varepsilon b} - e^{\varepsilon a}}{\varepsilon}, \\ \int_{-\infty}^{\infty} f(x) dx &= \lim_{\varepsilon \rightarrow 0} 2\pi f(-\iota \partial_\varepsilon) \delta(\varepsilon) = 2\pi \delta(\iota \partial_\varepsilon) f(\varepsilon), \\ \int_0^{\infty} f(x) dx &= \lim_{\varepsilon \rightarrow 0} f(-\partial_\varepsilon) \frac{1}{\varepsilon}, \\ \int_{-\infty}^0 f(x) dx &= \lim_{\varepsilon \rightarrow 0} f(\partial_\varepsilon) \frac{1}{\varepsilon}, \\ \int_{-\infty}^{\infty} f(x) dx &= \lim_{\varepsilon \rightarrow 0} [f(-\partial_\varepsilon) + f(\partial_\varepsilon)] \frac{1}{\varepsilon},\end{aligned}$$

where ∂_ε denotes the derivative with respect to ε .

Example

$$I = \int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

$$f(x) = \frac{1}{x} \cdot \frac{1}{2l} (e^{lx} - e^{-lx})$$

$$I = \lim_{\varepsilon \rightarrow 0} f(-\partial_\varepsilon) \frac{1}{\varepsilon} = \frac{1}{2l} \lim_{\varepsilon \rightarrow 0} (e^{-l\partial_\varepsilon} - e^{l\partial_\varepsilon}) \frac{1}{\partial_\varepsilon} \circ \frac{1}{\varepsilon}.$$

Note that $1/\partial_\varepsilon$ is the inverse operation of derivative, i.e., integration.

$$I = \frac{1}{2l} \lim_{\varepsilon \rightarrow 0} (e^{-l\partial_\varepsilon} - e^{l\partial_\varepsilon}) \circ (\ln \varepsilon + c)$$

Recall that for the derivative operator ∂_ε , so that

$$e^{a\partial_\varepsilon} \circ f(\varepsilon) = f(\varepsilon + a).$$

$$\begin{aligned} I &= \frac{1}{2l} \lim_{\varepsilon \rightarrow 0} [(\ln(\varepsilon - l) + c) - (\ln(\varepsilon + l) + c)] \\ &= \frac{1}{2l} \lim_{\varepsilon \rightarrow 0} [\ln(\varepsilon - l) - \ln(\varepsilon + l)] = \frac{1}{2l} \left(\frac{-l\pi}{2} - \frac{l\pi}{2} \right) = \frac{\pi}{2}. \end{aligned}$$

Example

$$I = \int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

$$f(x) = \frac{1}{x} \cdot \frac{1}{2\iota} (e^{\iota x} - e^{-\iota x})$$

$$I = \lim_{\varepsilon \rightarrow 0} f(-\partial_\varepsilon) \frac{1}{\varepsilon} = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_\varepsilon} - e^{\iota \partial_\varepsilon}) \frac{1}{\partial_\varepsilon} \circ \frac{1}{\varepsilon}.$$

Note that $1/\partial_\varepsilon$ is the inverse operation of derivative, i.e., integration.

$$I = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_\varepsilon} - e^{\iota \partial_\varepsilon}) \circ (\ln \varepsilon + c)$$

Recall that for the derivative operator ∂_ε , so that

$$e^{a\partial_\varepsilon} \circ f(\varepsilon) = f(\varepsilon + a).$$

$$\begin{aligned} I &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [(\ln(\varepsilon - \iota) + c) - (\ln(\varepsilon + \iota) + c)] \\ &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [\ln(\varepsilon - \iota) - \ln(\varepsilon + \iota)] = \frac{1}{2\iota} \left(\frac{-\iota\pi}{2} - \frac{\iota\pi}{2} \right) = \frac{\pi}{2}. \end{aligned}$$

Example

$$I = \int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

$$f(x) = \frac{1}{x} \cdot \frac{1}{2\iota} (e^{\iota x} - e^{-\iota x})$$

$$I = \lim_{\varepsilon \rightarrow 0} f(-\partial_{\varepsilon}) \frac{1}{\varepsilon} = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_{\varepsilon}} - e^{\iota \partial_{\varepsilon}}) \frac{1}{\partial_{\varepsilon}} \circ \frac{1}{\varepsilon}.$$

Note that $1/\partial_{\varepsilon}$ is the inverse operation of derivative, i.e., integration.

$$I = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_{\varepsilon}} - e^{\iota \partial_{\varepsilon}}) \circ (\ln \varepsilon + c)$$

Recall that for the derivative operator ∂_{ε} , so that

$$e^{a\partial_{\varepsilon}} \circ f(\varepsilon) = f(\varepsilon + a).$$

$$\begin{aligned} I &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [(\ln(\varepsilon - \iota) + c) - (\ln(\varepsilon + \iota) + c)] \\ &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [\ln(\varepsilon - \iota) - \ln(\varepsilon + \iota)] = \frac{1}{2\iota} \left(\frac{-\iota\pi}{2} - \frac{\iota\pi}{2} \right) = \frac{\pi}{2}. \end{aligned}$$

Example

$$I = \int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

$$f(x) = \frac{1}{x} \cdot \frac{1}{2l} (e^{lx} - e^{-lx})$$

$$I = \lim_{\varepsilon \rightarrow 0} f(-\partial_\varepsilon) \frac{1}{\varepsilon} = \frac{1}{2l} \lim_{\varepsilon \rightarrow 0} (e^{-l\partial_\varepsilon} - e^{l\partial_\varepsilon}) \frac{1}{\partial_\varepsilon} \circ \frac{1}{\varepsilon}.$$

Note that $1/\partial_\varepsilon$ is the inverse operation of derivative, i.e., integration.

$$I = \frac{1}{2l} \lim_{\varepsilon \rightarrow 0} (e^{-l\partial_\varepsilon} - e^{l\partial_\varepsilon}) \circ (\ln \varepsilon + c)$$

Recall that for the derivative operator ∂_ε , so that

$$e^{a\partial_\varepsilon} \circ f(\varepsilon) = f(\varepsilon + a).$$

$$\begin{aligned} I &= \frac{1}{2l} \lim_{\varepsilon \rightarrow 0} [(\ln(\varepsilon - l) + c) - (\ln(\varepsilon + l) + c)] \\ &= \frac{1}{2l} \lim_{\varepsilon \rightarrow 0} [\ln(\varepsilon - l) - \ln(\varepsilon + l)] = \frac{1}{2l} \left(\frac{-l\pi}{2} - \frac{l\pi}{2} \right) = \frac{\pi}{2}. \end{aligned}$$

Example

$$I = \int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

$$f(x) = \frac{1}{x} \cdot \frac{1}{2\iota} (e^{\iota x} - e^{-\iota x})$$

$$I = \lim_{\varepsilon \rightarrow 0} f(-\partial_\varepsilon) \frac{1}{\varepsilon} = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_\varepsilon} - e^{\iota \partial_\varepsilon}) \frac{1}{\partial_\varepsilon} \circ \frac{1}{\varepsilon}.$$

Note that $1/\partial_\varepsilon$ is the inverse operation of derivative, i.e., integration.

$$I = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_\varepsilon} - e^{\iota \partial_\varepsilon}) \circ (\ln \varepsilon + c)$$

Recall that for the derivative operator ∂_ε , so that

$$e^{a\partial_\varepsilon} \circ f(\varepsilon) = f(\varepsilon + a).$$

$$\begin{aligned} I &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [(\ln(\varepsilon - \iota) + c) - (\ln(\varepsilon + \iota) + c)] \\ &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [\ln(\varepsilon - \iota) - \ln(\varepsilon + \iota)] = \frac{1}{2\iota} \left(\frac{-\iota\pi}{2} - \frac{\iota\pi}{2} \right) = \frac{\pi}{2}. \end{aligned}$$

Example

$$I = \int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

$$f(x) = \frac{1}{x} \cdot \frac{1}{2\iota} (e^{\iota x} - e^{-\iota x})$$

$$I = \lim_{\varepsilon \rightarrow 0} f(-\partial_\varepsilon) \frac{1}{\varepsilon} = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_\varepsilon} - e^{\iota \partial_\varepsilon}) \frac{1}{\partial_\varepsilon} \circ \frac{1}{\varepsilon}.$$

Note that $1/\partial_\varepsilon$ is the inverse operation of derivative, i.e., integration.

$$I = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_\varepsilon} - e^{\iota \partial_\varepsilon}) \circ (\ln \varepsilon + c)$$

Recall that for the derivative operator ∂_ε , so that

$$e^{a\partial_\varepsilon} \circ f(\varepsilon) = f(\varepsilon + a).$$

$$\begin{aligned} I &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [(\ln(\varepsilon - \iota) + c) - (\ln(\varepsilon + \iota) + c)] \\ &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [\ln(\varepsilon - \iota) - \ln(\varepsilon + \iota)] = \frac{1}{2\iota} \left(\frac{-\iota\pi}{2} - \frac{\iota\pi}{2} \right) = \frac{\pi}{2}. \end{aligned}$$

Example

$$I = \int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

$$f(x) = \frac{1}{x} \cdot \frac{1}{2\iota} (e^{\iota x} - e^{-\iota x})$$

$$I = \lim_{\varepsilon \rightarrow 0} f(-\partial_\varepsilon) \frac{1}{\varepsilon} = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_\varepsilon} - e^{\iota \partial_\varepsilon}) \frac{1}{\partial_\varepsilon} \circ \frac{1}{\varepsilon}.$$

Note that $1/\partial_\varepsilon$ is the inverse operation of derivative, i.e., integration.

$$I = \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} (e^{-\iota \partial_\varepsilon} - e^{\iota \partial_\varepsilon}) \circ (\ln \varepsilon + c)$$

Recall that for the derivative operator ∂_ε , so that

$$e^{a\partial_\varepsilon} \circ f(\varepsilon) = f(\varepsilon + a).$$

$$\begin{aligned} I &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [(\ln(\varepsilon - \iota) + c) - (\ln(\varepsilon + \iota) + c)] \\ &= \frac{1}{2\iota} \lim_{\varepsilon \rightarrow 0} [\ln(\varepsilon - \iota) - \ln(\varepsilon + \iota)] = \frac{1}{2\iota} \left(\frac{-\iota\pi}{2} - \frac{\iota\pi}{2} \right) = \frac{\pi}{2}. \end{aligned}$$

Remark

$$\begin{aligned}
 I &= \int_0^\infty \frac{1}{x} \sum_n \phi_n \frac{\Gamma(n+1)}{\Gamma(2n+2)} x^{2n+1} dx \\
 &= \sum_n \phi_n \frac{\Gamma(n+1)}{\Gamma(2n+2)} \langle 2n+1 \rangle \\
 &= \frac{1}{2} \cdot \frac{\Gamma\left(-\frac{1}{2}+1\right)}{\Gamma\left(2\left(-\frac{1}{2}\right)+2\right)} \Gamma\left(\frac{1}{2}\right) \\
 &= \frac{\pi}{2}.
 \end{aligned}$$

Remark

$$\begin{aligned} I &= \int_0^\infty \frac{1}{x} \sum_n \phi_n \frac{\Gamma(n+1)}{\Gamma(2n+2)} x^{2n+1} dx \\ &= \sum_n \phi_n \frac{\Gamma(n+1)}{\Gamma(2n+2)} \langle 2n+1 \rangle \\ &= \frac{1}{2} \cdot \frac{\Gamma\left(-\frac{1}{2}+1\right)}{\Gamma\left(2\left(-\frac{1}{2}\right)+2\right)} \Gamma\left(\frac{1}{2}\right) \\ &= \frac{\pi}{2}. \end{aligned}$$

Proofs

D. Jia , E. Tang, A. Kempf, “present a list of propositions that put the above integration by differentiation methods on a rigorous footing.”

$$\int_0^{\infty} f(x) e^{-xy} dx = \lim_{a \rightarrow \infty} f(-\partial_y) \frac{1 - e^{-ay}}{y},$$

provided that $f : \mathbb{R} \rightarrow \mathbb{R}$ is entire and Laplace transformable on \mathbb{R}_+ . Formal/Key idea:

$$\begin{aligned} \int_0^{\infty} f(x) e^{-xy} dx &= \int_0^{\infty} \sum_{n=0}^{\infty} c_n x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a (-\partial_y)^n e^{-xy} dx. \end{aligned}$$

Proofs

D. Jia , E. Tang, A. Kempf, “present a list of propositions that put the above integration by differentiation methods on a rigorous footing.”

$$\int_0^{\infty} f(x) e^{-xy} dx = \lim_{a \rightarrow \infty} f(-\partial_y) \frac{1 - e^{-ay}}{y},$$

provided that $f : \mathbb{R} \rightarrow \mathbb{R}$ is entire and Laplace transformable on \mathbb{R}_+ . Formal/Key idea:

$$\begin{aligned} \int_0^{\infty} f(x) e^{-xy} dx &= \int_0^{\infty} \sum_{n=0}^{\infty} c_n x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a (-\partial_y)^n e^{-xy} dx. \end{aligned}$$

Proofs

D. Jia , E. Tang, A. Kempf, “present a list of propositions that put the above integration by differentiation methods on a rigorous footing.”

$$\int_0^{\infty} f(x) e^{-xy} dx = \lim_{a \rightarrow \infty} f(-\partial_y) \frac{1 - e^{-ay}}{y},$$

provided that $f : \mathbb{R} \rightarrow \mathbb{R}$ is entire and Laplace transformable on \mathbb{R}_+ . Formal/Key idea:

$$\begin{aligned} \int_0^{\infty} f(x) e^{-xy} dx &= \int_0^{\infty} \sum_{n=0}^{\infty} c_n x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a (-\partial_y)^n e^{-xy} dx. \end{aligned}$$

Proofs

D. Jia , E. Tang, A. Kempf, “present a list of propositions that put the above integration by differentiation methods on a rigorous footing.”

$$\int_0^{\infty} f(x) e^{-xy} dx = \lim_{a \rightarrow \infty} f(-\partial_y) \frac{1 - e^{-ay}}{y},$$

provided that $f : \mathbb{R} \rightarrow \mathbb{R}$ is entire and Laplace transformable on \mathbb{R}_+ . Formal/Key idea:

$$\begin{aligned} \int_0^{\infty} f(x) e^{-xy} dx &= \int_0^{\infty} \sum_{n=0}^{\infty} c_n x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a (-\partial_y)^n e^{-xy} dx. \end{aligned}$$

Proofs

D. Jia , E. Tang, A. Kempf, “present a list of propositions that put the above integration by differentiation methods on a rigorous footing.”

$$\int_0^{\infty} f(x) e^{-xy} dx = \lim_{a \rightarrow \infty} f(-\partial_y) \frac{1 - e^{-ay}}{y},$$

provided that $f : \mathbb{R} \rightarrow \mathbb{R}$ is entire and Laplace transformable on \mathbb{R}_+ . Formal/Key idea:

$$\begin{aligned} \int_0^{\infty} f(x) e^{-xy} dx &= \int_0^{\infty} \sum_{n=0}^{\infty} c_n x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a x^n e^{-xy} dx \\ &= \sum_{n=0}^{\infty} c_n \lim_{a \rightarrow \infty} \int_0^a (-\partial_y)^n e^{-xy} dx. \end{aligned}$$

Formal Connection

$$\begin{aligned}
 f(x) &= \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx = \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx. \\
 \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx &= \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} \left(\sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \right) dx \\
 &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n \int_0^{\infty} e^{-\varepsilon x} x^{\alpha n + \beta - 1} dx \\
 &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n \int_0^{\infty} \left((-\partial_{\varepsilon})^{\alpha n + \beta - 1} \circ e^{-\varepsilon x} \right) dx \\
 &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n (-\partial_{\varepsilon})^{\alpha n + \beta - 1} \circ \int_0^{\infty} e^{-\varepsilon x} dx \\
 &= \lim_{\varepsilon \rightarrow 0} f(-\partial_{\varepsilon}) \circ \frac{1}{\varepsilon}.
 \end{aligned}$$

Formal Connection

$$f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx = \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx.$$

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx &= \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} \left(\sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \right) dx \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n \int_0^{\infty} e^{-\varepsilon x} x^{\alpha n + \beta - 1} dx \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n \int_0^{\infty} \left((-\partial_{\varepsilon})^{\alpha n + \beta - 1} \circ e^{-\varepsilon x} \right) dx \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n (-\partial_{\varepsilon})^{\alpha n + \beta - 1} \circ \int_0^{\infty} e^{-\varepsilon x} dx \\ &= \lim_{\varepsilon \rightarrow 0} f(-\partial_{\varepsilon}) \circ \frac{1}{\varepsilon}. \end{aligned}$$

Formal Connection

$$f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx = \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx.$$

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx &= \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} \left(\sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \right) dx \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n \int_0^{\infty} e^{-\varepsilon x} x^{\alpha n + \beta - 1} dx \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n \int_0^{\infty} \left((-\partial_{\varepsilon})^{\alpha n + \beta - 1} \circ e^{-\varepsilon x} \right) dx \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{n=0}^{\infty} a_n (-\partial_{\varepsilon})^{\alpha n + \beta - 1} \circ \int_0^{\infty} e^{-\varepsilon x} dx \\ &= \lim_{\varepsilon \rightarrow 0} f(-\partial_{\varepsilon}) \circ \frac{1}{\varepsilon}. \end{aligned}$$

Formal Connection

Recall P_1 :

$$f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx = \sum_n a_n \langle \alpha n + \beta \rangle.$$

Formally,

$$\langle a \rangle := \int_0^{\infty} x^{a-1} dx.$$

and

$$\langle a \rangle_{\varepsilon} := \int_0^{\infty} x^{a-1} e^{-\varepsilon x} dx \Rightarrow \lim_{\varepsilon \rightarrow 0} \langle a \rangle_{\varepsilon} = \langle a \rangle.$$

Therefore

$$\begin{aligned} \sum_n a_n \langle \alpha n + \beta \rangle &= \lim_{\varepsilon \rightarrow 0} \sum_n a_n \langle \alpha n + \beta \rangle_{\varepsilon} \\ &= \lim_{\varepsilon \rightarrow 0} \sum_n a_n \int_0^{\infty} x^{\alpha n + \beta - 1} e^{-\varepsilon x} dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx. \end{aligned}$$

Formal Connection

Recall P_1 :

$$f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx = \sum_n a_n \langle \alpha n + \beta \rangle.$$

Formally,

$$\langle a \rangle := \int_0^{\infty} x^{a-1} dx.$$

and

$$\langle a \rangle_{\varepsilon} := \int_0^{\infty} x^{a-1} e^{-\varepsilon x} dx \Rightarrow \lim_{\varepsilon \rightarrow 0} \langle a \rangle_{\varepsilon} = \langle a \rangle.$$

Therefore

$$\begin{aligned} \sum_n a_n \langle \alpha n + \beta \rangle &= \lim_{\varepsilon \rightarrow 0} \sum_n a_n \langle \alpha n + \beta \rangle_{\varepsilon} \\ &= \lim_{\varepsilon \rightarrow 0} \sum_n a_n \int_0^{\infty} x^{\alpha n + \beta - 1} e^{-\varepsilon x} dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx. \end{aligned}$$

Formal Connection

Recall P_1 :

$$f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx = \sum_n a_n \langle \alpha n + \beta \rangle.$$

Formally,

$$\langle a \rangle := \int_0^{\infty} x^{a-1} dx.$$

and

$$\langle a \rangle_{\varepsilon} := \int_0^{\infty} x^{a-1} e^{-\varepsilon x} dx \Rightarrow \lim_{\varepsilon \rightarrow 0} \langle a \rangle_{\varepsilon} = \langle a \rangle.$$

Therefore

$$\begin{aligned} \sum_n a_n \langle \alpha n + \beta \rangle &= \lim_{\varepsilon \rightarrow 0} \sum_n a_n \langle \alpha n + \beta \rangle_{\varepsilon} \\ &= \lim_{\varepsilon \rightarrow 0} \sum_n a_n \int_0^{\infty} x^{\alpha n + \beta - 1} e^{-\varepsilon x} dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx. \end{aligned}$$

Formal Connection

Recall P_1 :

$$f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^{\infty} f(x) dx = \sum_n a_n \langle \alpha n + \beta \rangle.$$

Formally,

$$\langle a \rangle := \int_0^{\infty} x^{a-1} dx.$$

and

$$\langle a \rangle_{\varepsilon} := \int_0^{\infty} x^{a-1} e^{-\varepsilon x} dx \Rightarrow \lim_{\varepsilon \rightarrow 0} \langle a \rangle_{\varepsilon} = \langle a \rangle.$$

Therefore

$$\begin{aligned} \sum_n a_n \langle \alpha n + \beta \rangle &= \lim_{\varepsilon \rightarrow 0} \sum_n a_n \langle \alpha n + \beta \rangle_{\varepsilon} \\ &= \lim_{\varepsilon \rightarrow 0} \sum_n a_n \int_0^{\infty} x^{\alpha n + \beta - 1} e^{-\varepsilon x} dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} e^{-\varepsilon x} f(x) dx. \end{aligned}$$

Possible Future Work

- Connect MoB to IbD, and provide rigorous proofs of the former.
- Ramanujan's Master Theorem.
- Extension to other intervals.
- etc.

Possible Future Work

- Connect MoB to IbD, and provide rigorous proofs of the former.
- Ramanujan's Master Theorem.
- Extension to other intervals.
- etc.

Possible Future Work

- Connect MoB to IbD, and provide rigorous proofs of the former.
- Ramanujan's Master Theorem.
- Extension to other intervals.
- etc.

Possible Future Work

- Connect MoB to IbD, and provide rigorous proofs of the former.
- Ramanujan's Master Theorem.
- Extension to other intervals.
- etc.

Possible Future Work

- Connect MoB to IbD, and provide rigorous proofs of the former.
- Ramanujan's Master Theorem.
- Extension to other intervals.
- etc.

References



B. C. Berndt, *Ramanujan's Notebooks Part I*, Springer-Verlag, 1991.



K. Boyadzhiev, V. H. Moll. The integrals in Gradshteyn and Ryzhik. Part 21: hyperbolic functions. *Scientia*, 22 (2013), 109–127, 2013.



I. Gonzalez, K. Kohl, and V. H. Moll. Evaluation of entries in Gradshteyn and Ryzhik employing the method of brackets. *Scientia*, 25 (2014), 65–84.



I. Gonzalez and V. H. Moll. Definite integrals by the method of brackets. Part 1. *Adv. Appl. Math.*, 45 (2010), 50–73.



I. Gonzalez and I. Schmidt. Optimized negative dimensional integration method (NDIM) and multiloop Feynman diagram calculation. *Nuclear Phys. B*, 769 (2017), 124–173.



I. Gonzalez and I. Schmidt. Modular application of an integration by fractional expansion (IBFE) method to multiloop Feynman diagrams. *Phys. Rev. D*, 78 (2008), 086003.



I. G. Halliday and R. M. Ricotta. Negative dimensional integrals. I. Feynman graphs. *Phys. Lett. B*, 193 (1987), 241–246.



I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products*, Academic Press, 2015.



G. H. Hardy. *Ramanujan. Twelve Lectures on Subjects Suggested by His Life and Work*. Chelsea Publishing Company, 1987.



K. Kohl. *Algorithmic Methods for Definite Integration*. PhD thesis, Tulane University, 2011.

References



B. C. Berndt, *Ramanujan's Notebooks Part I*, Springer-Verlag, 1991.



K. Boyadzhiev, V. H. Moll. The integrals in Gradshteyn and Ryzhik. Part 21: hyperbolic functions. *Scientia*. **22** (2013), 109–127, 2013.



I. Gonzalez, K. Kohl, and V. H. Moll. Evaluation of entries in Gradshteyn and Ryzhik employing the method of brackets. *Scientia*, **25** (2014), 65–84.



I. Gonzalez and V. H. Moll. Definite integrals by the method of brackets. Part 1. *Adv. Appl. Math.*, **45** (2010), 50–73.



I. Gonzalez and I. Schmidt. Optimized negative dimensional integration method (NDIM) and multiloop Feynman diagram calculation. *Nuclear Phys. B*, **769** (2017), 124–173.



I. Gonzalez and I. Schmidt. Modular application of an integration by fractional expansion (IBFE) method to multiloop Feynman diagrams. *Phys. Rev. D*, **78** (2008), 086003.



I. G. Halliday and R. M. Ricotta. Negative dimensional integrals. I. Feynman graphs. *Phys. Lett. B*, **193** (1987), 241–246.



I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products*, Academic Press, 2015.



G. H. Hardy. *Ramanujan. Twelve Lectures on Subjects Suggested by His Life and Work*. Chelsea Publishing Company, 1987.



K. Kohl. *Algorithmic Methods for Definite Integration*. PhD thesis, Tulane University, 2011.

References



A. Kempf, D. M. Jackson, and A. H. Morales, *New Dirac Delta Function Based Methods with Applications to Perturbative Expansions in Quantum Field Theory*, Journal of Physics A: Mathematical and Theoretical 47 (41): 415-204, 2014



A. Kempf, D. M. Jackson, and A. H. Morales, *How to (Path-) Integrate by Differentiating*, preprint, arXiv:math/1507.04348, 2015.



D. Jia, E. Tang and A. Kempf, *Integration by differentiation: new proofs, methods and examples*, preprint, <https://arxiv.org/abs/1610.09702>, 2016.

References



A. Kempf, D. M. Jackson, and A. H. Morales, *New Dirac Delta Function Based Methods with Applications to Perturbative Expansions in Quantum Field Theory*, Journal of Physics A: Mathematical and Theoretical 47 (41): 415-204, 2014



A. Kempf, D. M. Jackson, and A. H. Morales, *How to (Path-) Integrate by Differentiating*, preprint, arXiv:math/1507.04348, 2015.



D. Jia, E. Tang and A. Kempf, *Integration by differentiation: new proofs, methods and examples*, preprint, <https://arxiv.org/abs/1610.09702>, 2016.

End

Thank you!