

The Probabilistic and Combinatorial Interpretations of the Bernoulli Symbol \mathcal{B}

Lin JIU

CMS Winter Conference 2017 @ University of Waterloo
Department of Mathematics and Statistics, Dalhousie University
Joint work with Diane Yahui Shi, Tianjin University

December 10th, 2017

Outlines

Bernoulli Symbol and Umbral Calculus

Probabilistic Aspect

Combinatorial Interpretation

Future Work

Definition

The Bernoulli numbers $(B_n)_{n=0}^{\infty}$ and Bernoulli polynomials $(B_n(x))_{n=0}^{\infty}$ can be defined by

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \text{ and } \frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$

Definition

The Bernoulli numbers $(B_n)_{n=0}^{\infty}$ and Bernoulli polynomials $(B_n(x))_{n=0}^{\infty}$ can be defined by

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \text{ and } \frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$

with the relation

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k.$$

Definition

The Bernoulli numbers $(B_n)_{n=0}^{\infty}$ and Bernoulli polynomials $(B_n(x))_{n=0}^{\infty}$ can be defined by

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \text{ and } \frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$

with the relation

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k.$$

The Bernoulli symbol \mathcal{B} satisfies the evaluation rule that

$$\mathcal{B}^n = B_n.$$

Definition

The Bernoulli numbers $(B_n)_{n=0}^{\infty}$ and Bernoulli polynomials $(B_n(x))_{n=0}^{\infty}$ can be defined by

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \text{ and } \frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$

with the relation

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k.$$

The Bernoulli symbol \mathcal{B} satisfies the evaluation rule that

$$\mathcal{B}^n = B_n.$$

Treat $t = \partial_x$, and

$$\frac{t}{e^t - 1} \bullet x^n = B_n(x)$$

Definition

The Bernoulli numbers $(B_n)_{n=0}^{\infty}$ and Bernoulli polynomials $(B_n(x))_{n=0}^{\infty}$ can be defined by

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \text{ and } \frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$

with the relation

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k.$$

The Bernoulli symbol \mathcal{B} satisfies the evaluation rule that

$$\mathcal{B}^n = B_n.$$

Treat $t = \partial_x$, and

$$\frac{t}{e^t - 1} \bullet x^n = B_n(x)$$

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} \mathcal{B}^{n-k} x^k = (\mathcal{B} + x)^n.$$

Examples



$$B'_n(x) = nB_{n-1}(x)$$

Examples



$$B'_n(x) = nB_{n-1}(x) \Leftrightarrow \frac{d}{dx}(\mathcal{B} + x)^n = n(\mathcal{B} + x)^{n-1}.$$

Examples



$$B'_n(x) = nB_{n-1}(x) \Leftrightarrow \frac{d}{dx}(\mathcal{B} + x)^n = n(\mathcal{B} + x)^{n-1}.$$



$$e^{\mathcal{B}t} = \frac{t}{e^t - 1}$$

Examples



$$B'_n(x) = nB_{n-1}(x) \Leftrightarrow \frac{d}{dx} (\mathcal{B} + x)^n = n(\mathcal{B} + x)^{n-1}.$$



$$\begin{aligned} e^{\mathcal{B}t} = \frac{t}{e^t - 1} &\Rightarrow e^{-\mathcal{B}t} = \frac{-t}{e^{-t} - 1} = \frac{te^t}{e^t - 1} = e^{(\mathcal{B}+1)t} \\ &\Rightarrow -\mathcal{B} = \mathcal{B} + 1 \end{aligned}$$

Examples



$$B'_n(x) = nB_{n-1}(x) \Leftrightarrow \frac{d}{dx}(\mathcal{B} + x)^n = n(\mathcal{B} + x)^{n-1}.$$



$$\begin{aligned} e^{\mathcal{B}t} = \frac{t}{e^t - 1} &\Rightarrow e^{-\mathcal{B}t} = \frac{-t}{e^{-t} - 1} = \frac{te^t}{e^t - 1} = e^{(\mathcal{B}+1)t} \\ &\Rightarrow -\mathcal{B} = \mathcal{B} + 1 \\ &\Rightarrow (-1)^n B_n(-x) = (-1)^n (\mathcal{B} - x)^n = B_n(x + 1) \end{aligned}$$

Examples



$$B'_n(x) = nB_{n-1}(x) \Leftrightarrow \frac{d}{dx}(\mathcal{B} + x)^n = n(\mathcal{B} + x)^{n-1}.$$



$$\begin{aligned} e^{\mathcal{B}t} = \frac{t}{e^t - 1} &\Rightarrow e^{-\mathcal{B}t} = \frac{-t}{e^{-t} - 1} = \frac{te^t}{e^t - 1} = e^{(\mathcal{B}+1)t} \\ &\Rightarrow -\mathcal{B} = \mathcal{B} + 1 \\ &\Rightarrow (-1)^n B_n(-x) = (-1)^n (\mathcal{B} - x)^n = B_n(x + 1) \end{aligned}$$

Bernoulli (Random) Symbol

A. Dixit et al. show that let $L \sim \frac{\pi}{2} \operatorname{sech}^2(\pi t)$ and $\mathcal{B} \sim \imath L - \frac{1}{2}$, then

Bernoulli (Random) Symbol

A. Dixit et al. show that let $L \sim \frac{\pi}{2} \operatorname{sech}^2(\pi t)$ and $\mathcal{B} \sim \imath L - \frac{1}{2}$, then

$$B_n = \mathbb{E}[\mathcal{B}^n] = \frac{\pi}{2} \int_{\mathbb{R}} \left(\imath t - \frac{1}{2} \right)^n \operatorname{sech}^2(\pi t) \, dt$$

Bernoulli (Random) Symbol

A. Dixit et al. show that let $L \sim \frac{\pi}{2} \operatorname{sech}^2(\pi t)$ and $\mathcal{B} \sim \imath L - \frac{1}{2}$, then

$$B_n = \mathbb{E}[\mathcal{B}^n] = \frac{\pi}{2} \int_{\mathbb{R}} \left(\imath t - \frac{1}{2} \right)^n \operatorname{sech}^2(\pi t) dt$$

and

$$B_n(x) = \frac{\pi}{2} \int_{\mathbb{R}} \left(x + \imath t - \frac{1}{2} \right)^n \operatorname{sech}^2(\pi t) dt = \mathbb{E}[(\mathcal{B} + x)^n].$$

Bernoulli (Random) Symbol

A. Dixit et al. show that let $L \sim \frac{\pi}{2} \operatorname{sech}^2(\pi t)$ and $\mathcal{B} \sim \imath L - \frac{1}{2}$, then

$$B_n = \mathbb{E}[\mathcal{B}^n] = \frac{\pi}{2} \int_{\mathbb{R}} \left(\imath t - \frac{1}{2} \right)^n \operatorname{sech}^2(\pi t) dt$$

and

$$B_n(x) = \frac{\pi}{2} \int_{\mathbb{R}} \left(x + \imath t - \frac{1}{2} \right)^n \operatorname{sech}^2(\pi t) dt = \mathbb{E}[(\mathcal{B} + x)^n].$$

By omitting expectation operator \mathbb{E} , we have

$$B_n = \mathcal{B}^n \text{ and } B_n(x) = (\mathcal{B} + x)^n.$$

Bernoulli (Random) Symbol

A. Dixit et al. show that let $L \sim \frac{\pi}{2} \operatorname{sech}^2(\pi t)$ and $\mathcal{B} \sim \imath L - \frac{1}{2}$, then

$$B_n = \mathbb{E}[\mathcal{B}^n] = \frac{\pi}{2} \int_{\mathbb{R}} \left(\imath t - \frac{1}{2} \right)^n \operatorname{sech}^2(\pi t) dt$$

and

$$B_n(x) = \frac{\pi}{2} \int_{\mathbb{R}} \left(x + \imath t - \frac{1}{2} \right)^n \operatorname{sech}^2(\pi t) dt = \mathbb{E}[(\mathcal{B} + x)^n].$$

By omitting expectation operator \mathbb{E} , we have

$$B_n = \mathcal{B}^n \text{ and } B_n(x) = (\mathcal{B} + x)^n.$$

Namely

$$\frac{t}{e^t - 1} \bullet = \mathbb{E}[\bullet]$$

Probabilistic Interpretation

For independent random variables X and Y , if $\mathbb{E} [e^{tX}] = F(x)$ and $\mathbb{E} [e^{tY}] = G(x)$, then

$$\mathbb{E} [e^{t(X+Y)}] = F(x) G(x).$$

Probabilistic Interpretation

For independent random variables X and Y , if $\mathbb{E} [e^{tX}] = F(x)$ and $\mathbb{E} [e^{tY}] = G(x)$, then

$$\mathbb{E} [e^{t(X+Y)}] = F(x) G(x).$$

Choose $X = x$ and $Y = \mathcal{B}$, then

$$\mathbb{E} [e^{tX}] = e^{tx} \text{ and } \mathbb{E} [e^{t\mathcal{B}}] = \frac{t}{e^t - 1}$$

Probabilistic Interpretation

For independent random variables X and Y , if $\mathbb{E}[e^{tX}] = F(x)$ and $\mathbb{E}[e^{tY}] = G(x)$, then

$$\mathbb{E}[e^{t(X+Y)}] = F(x) G(x).$$

Choose $X = x$ and $Y = \mathcal{B}$, then

$$\mathbb{E}[e^{tX}] = e^{tx} \text{ and } \mathbb{E}[e^{t\mathcal{B}}] = \frac{t}{e^t - 1}$$

$$\mathbb{E}[e^{t(x+\mathcal{B})}] = \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} \frac{\mathbb{E}[(x+\mathcal{B})^n]}{n!} t^n.$$

Probabilistic Interpretation

For independent random variables X and Y , if $\mathbb{E}[e^{tX}] = F(x)$ and $\mathbb{E}[e^{tY}] = G(x)$, then

$$\mathbb{E}[e^{t(X+Y)}] = F(x) G(x).$$

Choose $X = x$ and $Y = \mathcal{B}$, then

$$\mathbb{E}[e^{tX}] = e^{tx} \text{ and } \mathbb{E}[e^{t\mathcal{B}}] = \frac{t}{e^t - 1}$$

$$\mathbb{E}[e^{t(x+\mathcal{B})}] = \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} \frac{\mathbb{E}[(x+\mathcal{B})^n]}{n!} t^n.$$

$$B_n(x) = \mathbb{E}[(\mathcal{B} + x)^n] = \frac{[t^n] e^{\mathcal{B}t} e^{xt}}{n!} = \frac{[t^n] \frac{te^{xt}}{e^t - 1}}{n!}.$$

Generalization

- Bernoulli:

$$\frac{t}{e^t - 1} e^{tx} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} \Leftrightarrow B_n(x) = (x + \mathcal{B})^n$$

- Nörlund:

$$\left(\frac{t}{e^t - 1} \right)^p e^{tx} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} \Leftrightarrow B_n^{(p)}(x) = (x + \mathcal{B}_1 + \dots + \mathcal{B}_p)^n$$

- Bernoulli-Barnes: for $\mathbf{a} = (a_1, \dots, a_k)$.

$$e^{tx} \prod_{i=1}^k \frac{t}{e^{a_i t} - 1} = \sum_{n=0}^{\infty} B_n(\mathbf{a}; x) \frac{t^n}{n!}$$

$$\Leftrightarrow B_n(\mathbf{a}; x) = \frac{1}{|\mathbf{a}|} \left(x + \mathbf{a} \cdot \vec{\mathcal{B}} \right)^n, \text{ where } \begin{cases} \mathbf{a} = (a_1, \dots, a_k) \\ \vec{\mathcal{B}} = (\mathcal{B}_1, \dots, \mathcal{B}_k) \\ \mathbf{a} \cdot \vec{\mathcal{B}} = \sum_{l=1}^k a_l \mathcal{B}_l \\ |\mathbf{a}| = \prod_{l=1}^k a_l \end{cases}$$

Several Results

Bernoulli-Barnes

$$e^{tx} \prod_{i=1}^k \frac{t}{e^{a_i t} - 1} = \sum_{n=0}^{\infty} B_n(\mathbf{a}; x) \frac{t^n}{n!} \Leftrightarrow B_n(\mathbf{a}; x) = \frac{1}{|\mathbf{a}|} \left(x + \mathbf{a} \cdot \vec{B} \right)^n$$

Theorem[L. Jiu, V. Moll and C. Vignat]

$$f\left(x - \mathbf{a} \cdot \vec{B}\right) = \sum_{\ell=0}^n \sum_{|L|=\ell} |\mathbf{a}|_{L*} f^{(n-\ell)}\left(x + \left(\mathbf{a} \cdot \vec{B}\right)_L\right).$$

The multiple zeta function

$$\zeta_r(n_1, \dots, n_r) = \sum_{0 < k_1 < \dots < k_r} \frac{1}{k_1^{n_1} \dots k_r^{n_r}} = \sum_{k_1, \dots, k_r=1}^{\infty} \frac{1}{k_1^{n_1} (k_1 + k_2)^{n_2} \dots (k_1 + \dots + k_r)^{n_r}}$$

Theorem[L. Jiu, V. Moll and C. Vignat]

$$\zeta_r(-n_1, \dots, -n_r) = \prod_{k=1}^r (-1)^{n_k} c_{1, \dots, k}^{n_k+1},$$

where

$$c_1^n = \frac{\mathcal{B}_1^n}{n}, c_{1,2}^n = \frac{(\mathcal{C}_1 + \mathcal{B}_2)^n}{n}, \dots, c_{1, \dots, k+1}^n = \frac{(\mathcal{C}_{1, \dots, k} + \mathcal{B}_{k+1})^n}{n}$$

Uniqueness of Hyperbolic Secant Square

Uniqueness of Hyperbolic Secant Square

$$\mathcal{B} \sim \imath L - \frac{1}{2}.$$

Uniqueness of Hyperbolic Secant Square

$\mathcal{B} \sim \imath L - \frac{1}{2}$. Define $\bar{B}_n := |B_n(\frac{1}{2})|$, then

$$\bar{B}_n = \frac{\pi}{2} \int_{\mathbb{R}} t^n \operatorname{sech}^2(\pi t) dt.$$

Uniqueness of Hyperbolic Secant Square

$\mathcal{B} \sim \imath L - \frac{1}{2}$. Define $\bar{B}_n := |B_n(\frac{1}{2})|$, then

$$\bar{B}_n = \frac{\pi}{2} \int_{\mathbb{R}} t^n \operatorname{sech}^2(\pi t) dt. \left(\frac{x/2}{\sin(x/2)} = \sum_{n=0}^{\infty} \bar{B}_n \frac{x^n}{n!} \right)$$

Uniqueness of Hyperbolic Secant Square

$\mathcal{B} \sim \imath L - \frac{1}{2}$. Define $\bar{B}_n := |B_n(\frac{1}{2})|$, then

$$\bar{B}_n = \frac{\pi}{2} \int_{\mathbb{R}} t^n \operatorname{sech}^2(\pi t) dt. \left(\frac{x/2}{\sin(x/2)} = \sum_{n=0}^{\infty} \bar{B}_n \frac{x^n}{n!} \right)$$

Theorem

$\frac{\pi}{2} \operatorname{sech}^2(\pi t) dt$ is the *UNIQUE* density on \mathbb{R} for $(\bar{B}_n)_{n=0}^{\infty}$.

Uniqueness of Hyperbolic Secant Square

$\mathcal{B} \sim \imath L - \frac{1}{2}$. Define $\bar{B}_n := |B_n(\frac{1}{2})|$, then

$$\bar{B}_n = \frac{\pi}{2} \int_{\mathbb{R}} t^n \operatorname{sech}^2(\pi t) dt. \left(\frac{x/2}{\sin(x/2)} = \sum_{n=0}^{\infty} \bar{B}_n \frac{x^n}{n!} \right)$$

Theorem

$\frac{\pi}{2} \operatorname{sech}^2(\pi t) dt$ is the *UNIQUE* density on \mathbb{R} for $(\bar{B}_n)_{n=0}^{\infty}$.

Proof.

$$(-1)^{n+1} B_{2n} \sim \frac{2(2n)!}{(2\pi)^{2n}}.$$



Lemma

Uniqueness is equivalent to existence of constants C and D , such that

$$|\bar{B}_n| \leq CD^n n!.$$

Cumulants

$$K(t) := \sum_{n=1}^{\infty} \kappa_n \frac{t^n}{n!} = \log \mathbb{E} \left[e^{tX} \right] = \log \left(\sum_{n=0}^{\infty} \frac{\mathbb{E}[X^n]}{n!} t^n \right).$$

Cumulants

$$K(t) := \sum_{n=1}^{\infty} \kappa_n \frac{t^n}{n!} = \log \mathbb{E} \left[e^{tX} \right] = \log \left(\sum_{n=0}^{\infty} \frac{\mathbb{E}[X^n]}{n!} t^n \right).$$

Theorem

[Faà di Bruno's formula] For moments $(m_n)_{n=0}^{\infty}$ and cumulants $(\kappa_n)_{n=1}^{\infty}$ it holds that

$$m_n = Y_n(\kappa_1, \dots, \kappa_n) \text{ and } \kappa_n = \sum_{k=1}^n (-1)^{k-1} (k-1)! Y_{n,k}(m_1, \dots, m_{n-k+1}),$$

where, the partial or incomplete exponential Bell polynomial is given by

$$Y_{n,k}(x_1, \dots, x_{n-k+1}) := \sum_{\substack{j_1 + \dots + j_{n-k+1} = k \\ j_1 + 2j_2 + \dots + (n-k+1)j_{n-k+1} = n}} \frac{n!}{j_1! \dots j_{n-k+1}!} \left(\frac{x_1}{1!} \right)^{j_1} \dots \left(\frac{x_{n-k+1}}{(n-k+1)!} \right)^{j_{n-k+1}},$$

and the n^{th} complete exponential Bell polynomial is given by the sum

$$Y_n(x_1, \dots, x_n) := \sum_{k=1}^n Y_{n,k}(x_1, \dots, x_{n-k+1}) = \sum_{k = \left(\underbrace{1, \dots, 1}_{k_1}, \dots, \underbrace{n, \dots, n}_{k_n} \right) \vdash n} \frac{n!}{k_1! \dots k_n!} \left(\frac{x_1}{1!} \right)^{k_1} \dots \left(\frac{x_n}{n!} \right)^{k_n}.$$

Cumulants

Theorem

$$B_n \left(\frac{1}{2} \right) = Y_n \left(0, -\frac{B_2}{2}, -\frac{B_3}{3}, \dots, -\frac{B_n}{n} \right),$$

and

$$B_n = -n \sum_{k=1}^n (-1)^{k-1} (k-1)! Y_{n,k} \left(B_0 \left(\frac{1}{2} \right), \dots, B_{n-k+1} \left(\frac{1}{2} \right) \right).$$

Cumulants

Theorem

$$B_n \left(\frac{1}{2} \right) = Y_n \left(0, -\frac{B_2}{2}, -\frac{B_3}{3}, \dots, -\frac{B_n}{n} \right),$$

and

$$B_n = -n \sum_{k=1}^n (-1)^{k-1} (k-1)! Y_{n,k} \left(B_0 \left(\frac{1}{2} \right), \dots, B_{n-k+1} \left(\frac{1}{2} \right) \right).$$

$$Y_k \left(-\frac{B_2 \cdot 1!}{2 \cdot 2!}, \dots, -\frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k! B_{2k} \left(\frac{1}{2} \right)}{(2k)!} = \frac{k!}{(2k)!} \cdot (2^{2k-1} - 1) B_{2k}.$$

Cumulants

Theorem

$$B_n \left(\frac{1}{2} \right) = Y_n \left(0, -\frac{B_2}{2}, -\frac{B_3}{3}, \dots, -\frac{B_n}{n} \right),$$

and

$$B_n = -n \sum_{k=1}^n (-1)^{k-1} (k-1)! Y_{n,k} \left(B_0 \left(\frac{1}{2} \right), \dots, B_{n-k+1} \left(\frac{1}{2} \right) \right).$$

$$Y_k \left(-\frac{B_2 \cdot 1!}{2 \cdot 2!}, \dots, -\frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k! B_{2k} \left(\frac{1}{2} \right)}{(2k)!} = \frac{k!}{(2k)!} \cdot \left(2^{2k-1} - 1 \right) B_{2k}.$$

[M. Hoffman]

$$Y_k \left(\frac{B_2 \cdot 1!}{2 \cdot 2!}, \frac{B_4 \cdot 2!}{4 \cdot 4!}, \dots, \frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k!}{2^{2k} (2k+1)!}.$$

Cumulants

Theorem

$$B_n \left(\frac{1}{2} \right) = Y_n \left(0, -\frac{B_2}{2}, -\frac{B_3}{3}, \dots, -\frac{B_n}{n} \right),$$

and

$$B_n = -n \sum_{k=1}^n (-1)^{k-1} (k-1)! Y_{n,k} \left(B_0 \left(\frac{1}{2} \right), \dots, B_{n-k+1} \left(\frac{1}{2} \right) \right).$$

$$Y_k \left(-\frac{B_2 \cdot 1!}{2 \cdot 2!}, \dots, -\frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k! B_{2k} \left(\frac{1}{2} \right)}{(2k)!} = \frac{k!}{(2k)!} \cdot \left(2^{2k-1} - 1 \right) B_{2k}.$$

[M. Hoffman]

$$Y_k \left(\frac{B_2 \cdot 1!}{2 \cdot 2!}, \frac{B_4 \cdot 2!}{4 \cdot 4!}, \dots, \frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k!}{2^{2k} (2k+1)!}.$$

Remark

The first one is a special result of B. Robinstein:

<https://arxiv.org/abs/0911.3069>.

Cumulants

Consider different moment generating function

$$M_Y(t) = \mathbb{E}[e^{tY}] = \frac{\sinh \frac{t}{2}}{\frac{t}{2}}$$

Theorem

It also holds that

$$B_n = n \sum_{k=1}^n (-1)^{k-1} (k-1)! Y_{n,k} \left(0, \frac{1}{4 \cdot 3}, 0, \dots, \frac{1 + (-1)^{n-k+2}}{2^{n-k+2} (n-k+2)} \right).$$

Cumulants

Consider different moment generating function

$$M_Y(t) = \mathbb{E}[e^{tY}] = \frac{\sinh \frac{t}{2}}{\frac{t}{2}}$$

Theorem

It also holds that

$$B_n = n \sum_{k=1}^n (-1)^{k-1} (k-1)! Y_{n,k} \left(0, \frac{1}{4 \cdot 3}, 0, \dots, \frac{1 + (-1)^{n-k+2}}{2^{n-k+2} (n-k+2)} \right).$$

$$Y_k \left(-\frac{B_2 \cdot 1!}{2 \cdot 2!}, \dots, -\frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k!}{(2k)!} \cdot (2^{2k-1} - 1) B_{2k}$$

$$Y_k \left(\frac{B_2 \cdot 1!}{2 \cdot 2!}, \frac{B_4 \cdot 2!}{4 \cdot 4!}, \dots, \frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k!}{2^{2k} (2k+1)!}$$

Cumulants

$$Y_k \left(-\frac{B_2 \cdot 1!}{2 \cdot 2!}, \dots, -\frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k!}{(2k)!} \cdot (2^{2k-1} - 1) B_{2k}$$
$$Y_k \left(\frac{B_2 \cdot 1!}{2 \cdot 2!}, \frac{B_4 \cdot 2!}{4 \cdot 4!}, \dots, \frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k!}{2^{2k} (2k+1)!}$$

Cumulants

$$Y_k \left(-\frac{B_2 \cdot 1!}{2 \cdot 2!}, \dots, -\frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k!}{(2k)!} \cdot (2^{2k-1} - 1) B_{2k}$$
$$Y_k \left(\frac{B_2 \cdot 1!}{2 \cdot 2!}, \frac{B_4 \cdot 2!}{4 \cdot 4!}, \dots, \frac{B_{2k} \cdot k!}{(2k) \cdot (2k)!} \right) = \frac{k!}{2^{2k} (2k+1)!}$$

$$f(x) := \sum_{k=0}^{\infty} \frac{B_{2k} k!}{(2k) (2k)!} = \log \left(\frac{e^x - 1}{x} \right) - \frac{x}{2}$$

and

$$\sum_{k=0}^{\infty} \frac{x^{2k}}{2^{2k} (2k+1)!} = \frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} = e^{f(x)}.$$

Continued Fractions & Orthogonal Polynomials

$$(m_n)_{n=0}^{\infty} \sim m_n = \int_{\mathbb{R}} x^n d\mu(x)$$

Continued Fractions & Orthogonal Polynomials

$$(m_n)_{n=0}^{\infty} \sim m_n = \int_{\mathbb{R}} x^n d\mu(x) \quad \stackrel{?}{\Rightarrow} \quad (P_n(x))_{n=1}^{\infty} \sim \int_{\mathbb{R}} P_n(x) P_m(x) d\mu(x) = C_n \delta_{m,n}$$

Continued Fractions & Orthogonal Polynomials

$$\begin{aligned}
 (m_n)_{n=0}^\infty \sim m_n = \int_{\mathbb{R}} x^n d\mu(x) &\stackrel{?}{\Rightarrow} (P_n(x))_{n=1}^\infty \sim \int_{\mathbb{R}} P_n(x) P_m(x) d\mu(x) = C_n \delta_{m,n} \\
 &\Rightarrow P_{n+1}(x) = (x + s_n) P_n(x) - t_n P_{n-1}(x) \\
 &\Rightarrow \sum_{n=0}^{\infty} m_n x^n = \frac{m_0}{1 - s_0 x - \frac{t_1 x^2}{1 - s_1 x - \frac{t_2 x^2}{1 - \dots}}}
 \end{aligned}$$

Continued Fractions & Orthogonal Polynomials

$$\begin{aligned}
 (m_n)_{n=0}^\infty \sim m_n = \int_{\mathbb{R}} x^n d\mu(x) &\stackrel{?}{\Rightarrow} (P_n(x))_{n=1}^\infty \sim \int_{\mathbb{R}} P_n(x) P_m(x) d\mu(x) = C_n \delta_{m,n} \\
 &\Rightarrow P_{n+1}(x) = (x + s_n) P_n(x) - t_n P_{n-1}(x) \\
 &\Rightarrow \sum_{n=0}^{\infty} m_n x^n = \frac{m_0}{1 - s_0 x - \frac{t_1 x^2}{1 - s_1 x - \frac{t_2 x^2}{1 - \dots}}}
 \end{aligned}$$

Theorem [J. Touchard]

The polynomial sequence (ϕ_n) , define by

$$\phi_{n+1}(z) = \left(z + \frac{1}{2}\right) \phi_n(z) + \omega_n \phi_{n-1}(z)$$

satisfies for any $0 \leq r < n$, $\mathcal{B}^r \phi_n(\mathcal{B}) = 0$, where

$$\omega_n = \frac{n^4}{4(2n+1)(2n-1)}.$$

How

How

$$\psi_1(z) := \psi'(z) := (\log(\Gamma(z)))''$$

How

$$\psi_1(z) := \psi'(z) := (\log(\Gamma(z)))''$$

$$\sum_{n=0}^{\infty} \frac{B_n}{z^{n+1}} \sim \psi_1(z+1) = \frac{1}{z + \frac{1}{2} + \frac{\omega_1}{z + \frac{1}{2} + \frac{\omega_2}{z + \frac{1}{2} + \dots}}}$$

How

$$\psi_1(z) := \psi'(z) := (\log(\Gamma(z)))''$$

$$\sum_{n=0}^{\infty} \frac{B_n}{z^{n+1}} \sim \psi_1(z+1) = \frac{1}{z + \frac{1}{2} + \frac{\frac{\omega_1}{z + \frac{1}{2}} + \frac{\omega_2}{z + \frac{1}{2} + \dots}}{}}$$

$$\psi(z+x) \sim \log(z) - \sum_{n=1}^{\infty} \frac{(-1)^n B_n(x)}{nz^n}$$

How

$$\psi_1(z) := \psi'(z) := (\log(\Gamma(z)))''$$

$$\sum_{n=0}^{\infty} \frac{B_n}{z^{n+1}} \sim \psi_1(z+1) = \frac{1}{z + \frac{1}{2} + \frac{\frac{\omega_1}{z + \frac{1}{2}} + \frac{\omega_2}{z + \frac{1}{2} + \dots}}{}}$$

$$\psi(z+x) \sim \log(z) - \sum_{n=1}^{\infty} \frac{(-1)^n B_n(x)}{nz^n}$$

A. Dixit et al. showed

$$\log(\mathcal{B} + z) = \psi\left(\left|z - \frac{1}{2}\right| + \frac{1}{2}\right).$$

How

$$\psi_1(z) := \psi'(z) := (\log(\Gamma(z)))''$$

$$\sum_{n=0}^{\infty} \frac{B_n}{z^{n+1}} \sim \psi_1(z+1) = \frac{1}{z + \frac{1}{2} + \frac{\omega_1}{z + \frac{1}{2} + \frac{\omega_2}{z + \frac{1}{2} + \dots}}}$$

$$\psi(z+x) \sim \log(z) - \sum_{n=1}^{\infty} \frac{(-1)^n B_n(x)}{nz^n}$$

A. Dixit et al. showed

$$\log(B+z) = \psi\left(\left|z - \frac{1}{2}\right| + \frac{1}{2}\right).$$

Theorem

$$\varphi_{n+1}(z, x) := \left(z + \frac{1}{2} - x\right) \varphi_n(z, x) + \omega_n \varphi_{n-1}(z, x)$$

How

$$\psi_1(z) := \psi'(z) := (\log(\Gamma(z)))''$$

$$\sum_{n=0}^{\infty} \frac{B_n}{z^{n+1}} \sim \psi_1(z+1) = \frac{1}{z + \frac{1}{2} + \frac{\omega_1}{z + \frac{1}{2} + \frac{\omega_2}{z + \frac{1}{2} + \dots}}}$$

$$\psi(z+x) \sim \log(z) - \sum_{n=1}^{\infty} \frac{(-1)^n B_n(x)}{nz^n}$$

A. Dixit et al. showed

$$\log(\mathcal{B} + z) = \psi\left(\left|z - \frac{1}{2}\right| + \frac{1}{2}\right).$$

Theorem

$$\varphi_{n+1}(z, x) := \left(z + \frac{1}{2} - x\right) \varphi_n(z, x) + \omega_n \varphi_{n-1}(z, x)$$


$$z^r \varphi_n(z, x) \Big|_{z=\mathcal{B}+r} = (\mathcal{B} + x)^r \varphi_n(\mathcal{B} + x, x) = 0, \quad \forall 0 \leq r < n.$$

(Generalized) Motzkin Numbers

$$M_{n+1,k} = M_{n,k-1} + s_k M_{n,k} + t_{k+1} M_{n,k+1}$$


(Generalized) Motzkin Numbers

$$M_{n+1,k} = M_{n,k-1} + s_k M_{n,k} + t_{k+1} M_{n,k+1}$$




(Generalized) Motzkin Numbers

$$M_{n+1,k} = M_{n,k-1} + s_k M_{n,k} + t_{k+1} M_{n,k+1}$$



(Generalized) Motzkin Numbers

$$M_{n+1,k} = M_{n,k-1} + s_k M_{n,k} + t_{k+1} M_{n,k+1}$$



$$\sum_{n=0}^{\infty} M_{n,0} z^n = \frac{1}{1 - s_0 z - \frac{t_1 z^2}{1 - s_1 z - \frac{t_2 z^2}{\dots}}}$$

Combinatorial Interpretation

Theorem

Define $\left(M_{n,k}^{\mathbf{x},\omega}\right)_{n,k=0}^{\infty}$, by $M_{0,0}^{\mathbf{x},\omega} = 1$, $M_{n,k}^{\mathbf{x},\omega} = 0$ if $k > n$, and the recurrence

$$M_{n+1,k}^{\mathbf{x},\omega} = M_{n,k-1}^{\mathbf{x},\omega} + x_k M_{n,k}^{\mathbf{x},\omega} - \omega_{k+1} M_{n,k+1}^{\mathbf{x},\omega},$$

where $\mathbf{x} = (x_n)_{n=0}^{\infty}$ is given by $x_n = x - \frac{1}{2}$, and $\omega = (\omega_n)_{n=1}^{\infty}$ by $\omega_n = \frac{n^4}{4(2n+1)(2n-1)}$. Then, $M_{n,0}^{\mathbf{x},\omega} = B_n(x)$.

Combinatorial Interpretation

Theorem

Define $\left(M_{n,k}^{\mathbf{x},\omega}\right)_{n,k=0}^{\infty}$, by $M_{0,0}^{\mathbf{x},\omega} = 1$, $M_{n,k}^{\mathbf{x},\omega} = 0$ if $k > n$, and the recurrence

$$M_{n+1,k}^{\mathbf{x},\omega} = M_{n,k-1}^{\mathbf{x},\omega} + x_k M_{n,k}^{\mathbf{x},\omega} - \omega_{k+1} M_{n,k+1}^{\mathbf{x},\omega},$$

where $\mathbf{x} = (x_n)_{n=0}^{\infty}$ is given by $x_n = x - \frac{1}{2}$, and $\omega = (\omega_n)_{n=1}^{\infty}$ by $\omega_n = \frac{n^4}{4(2n+1)(2n-1)}$. Then, $M_{n,0}^{\mathbf{x},\omega} = B_n(x)$. In addition, the lattice path interpretation allows us to define the infinite-dimensional matrix

$$R_{\mathbf{x},\omega} := \begin{pmatrix} x - \frac{1}{2} & -\omega_1 & 0 & 0 & \cdots & 0 & \cdots \\ 1 & x - \frac{1}{2} & -\omega_2 & 0 & \cdots & 0 & \cdots \\ 0 & 1 & x - \frac{1}{2} & \ddots & \ddots & \vdots & \cdots \\ 0 & 0 & 1 & \ddots & -\omega_n & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots & x - \frac{1}{2} & -\omega_{n+1} & \cdots \\ 0 & 0 & 0 & \ddots & 1 & \ddots & \ddots \\ \vdots & \vdots & \vdots & \cdots & \vdots & \ddots & \ddots \end{pmatrix}.$$

Matrix Computation

Direct computations shows

$$R_{\mathbf{x},\omega,4} = \begin{pmatrix} x - 1/2 & -\frac{1}{12} & 0 & 0 \\ 1 & x - 1/2 & -\frac{4}{15} & 0 \\ 0 & 1 & x - 1/2 & -\frac{81}{140} \\ 0 & 0 & 1 & x - 1/2 \end{pmatrix}$$

Matrix Computation

Direct computations shows

$$R_{\mathbf{x},\omega,4} = \begin{pmatrix} x - 1/2 & -\frac{1}{12} & 0 & 0 \\ 1 & x - 1/2 & -\frac{4}{15} & 0 \\ 0 & 1 & x - 1/2 & -\frac{81}{140} \\ 0 & 0 & 1 & x - 1/2 \end{pmatrix}$$

and

$$(R_{\mathbf{x},\omega,4}^4) = \begin{pmatrix} x^4 - 2x^3 + x^2 - \frac{1}{30} & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{pmatrix},$$

Matrix Computation

Direct computations shows

$$R_{\mathbf{x},\omega,4} = \begin{pmatrix} x - 1/2 & -\frac{1}{12} & 0 & 0 \\ 1 & x - 1/2 & -\frac{4}{15} & 0 \\ 0 & 1 & x - 1/2 & -\frac{81}{140} \\ 0 & 0 & 1 & x - 1/2 \end{pmatrix}$$

and

$$(R_{\mathbf{x},\omega,4}^4) = \begin{pmatrix} x^4 - 2x^3 + x^2 - \frac{1}{30} & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{pmatrix},$$

where noting

$$B_4(x) = x^4 - 2x^3 + x^2 - \frac{1}{30}.$$

Euler Analogue

Definition

Euler numbers $(E_n)_{n=0}^{\infty}$ and Euler polynomials $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

Euler Analogue

Definition

Euler numbers $(E_n)_{n=0}^{\infty}$ and Euler polynomials $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

In addition,

$$E_n(x) = \int_{\mathbb{R}} \left(x - \frac{1}{2} + it \right)^n \operatorname{sech}(\pi t) dt.$$

Euler Analogue

Definition

Euler numbers $(E_n)_{n=0}^{\infty}$ and Euler polynomials $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

In addition,

$$E_n(x) = \int_{\mathbb{R}} \left(x - \frac{1}{2} + it \right)^n \operatorname{sech}(\pi t) dt.$$

$$E_n = 2^n E_n\left(\frac{1}{2}\right)$$

Euler Analogue

Definition

Euler numbers $(E_n)_{n=0}^{\infty}$ and Euler polynomials $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

In addition,

$$E_n(x) = \int_{\mathbb{R}} \left(x - \frac{1}{2} + it \right)^n \operatorname{sech}(\pi t) dt.$$

$E_n = 2^n E_n\left(\frac{1}{2}\right) \Rightarrow \mathcal{E} \sim 2\iota L_E$, where L_E has its density function $\operatorname{sech}(\pi t)$.

Euler Analogue

Definition

Euler numbers $(E_n)_{n=0}^{\infty}$ and Euler polynomials $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

In addition,

$$E_n(x) = \int_{\mathbb{R}} \left(x - \frac{1}{2} + it \right)^n \operatorname{sech}(\pi t) dt.$$

$E_n = 2^n E_n\left(\frac{1}{2}\right) \Rightarrow \mathcal{E} \sim 2\iota L_E$, where L_E has its density function $\operatorname{sech}(\pi t)$.

$$\mathcal{E}^n := \mathbb{E}[\mathcal{E}^n] = E_n$$

Conversely, it holds that $\mathbb{E}[L_E^n] = \left(\frac{\iota}{2}\right)^n E_n$ and $\mathbb{E}[e^{tL_E}] = \sec\left(\frac{t}{2}\right)$.

Euler Analogue

- Uniqueness of $\operatorname{sech}(\pi t)$ for L_E

Euler Analogue

- Uniqueness of $\operatorname{sech}(\pi t)$ for L_E ✓

Euler Analogue

- Uniqueness of $\operatorname{sech}(\pi t)$ for $L_E \checkmark (-1)^n E_{2n} \sim 8\sqrt{n/\pi} (4n/\pi/e)^{2n}$

Euler Analogue

- ▶ Uniqueness of $\operatorname{sech}(\pi t)$ for $L_E \checkmark (-1)^n E_{2n} \sim 8\sqrt{n/\pi} (4n/\pi/e)^{2n}$
- ▶ Faà di Bruno's formula:

Euler Analogue

- ▶ Uniqueness of $\operatorname{sech}(\pi t)$ for $L_E \checkmark (-1)^n E_{2n} \sim 8\sqrt{n/\pi} (4n/\pi/e)^{2n}$
- ▶ Faà di Bruno's formula:

$$\begin{cases} E_{2n} = 1 - \sum_{k=1}^n \binom{2n}{2k-1} \frac{2^{2k}(2^{2k}-1)B_{2k}}{2k} \\ B_{2n} = \frac{2n}{2^{2n}(2^{2n}-1)} \sum_{k=0}^{n-1} \binom{2n-1}{2k} E_{2k} \end{cases}$$

Euler Analogue

- ▶ Uniqueness of $\operatorname{sech}(\pi t)$ for $L_E \checkmark (-1)^n E_{2n} \sim 8\sqrt{n/\pi} (4n/\pi/e)^{2n}$
- ▶ Faà di Bruno's formula:

$$\begin{cases} E_{2n} = 1 - \sum_{k=1}^n \binom{2n}{2k-1} \frac{2^{2k}(2^{2k}-1)B_{2k}}{2k} \\ B_{2n} = \frac{2n}{2^{2n}(2^{2n}-1)} \sum_{k=0}^{n-1} \binom{2n-1}{2k} E_{2k} \end{cases}$$

- ▶ Orthogonal polynomials, Motzkin number, continued fractions

Euler Analogue

- ▶ Uniqueness of $\operatorname{sech}(\pi t)$ for $L_E \checkmark (-1)^n E_{2n} \sim 8\sqrt{n/\pi} (4n/\pi/e)^{2n}$
- ▶ Faà di Bruno's formula:

$$\begin{cases} E_{2n} = 1 - \sum_{k=1}^n \binom{2n}{2k-1} \frac{2^{2k}(2^{2k}-1)B_{2k}}{2k} \\ B_{2n} = \frac{2n}{2^{2n}(2^{2n}-1)} \sum_{k=0}^{n-1} \binom{2n-1}{2k} E_{2k} \end{cases}$$

- ▶ Orthogonal polynomials, Motzkin number, continued fractions

$$2\beta\left(\frac{s+1}{2}\right) \sim \sum_{j=1}^{\infty} \frac{E_j}{s^{j+1}}$$

Possible Extension to Nörlund Polynomials

$$\left(\frac{t}{e^t - 1}\right)^p e^{xt} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} \Leftrightarrow B_n^{(p)}(x) = (\mathcal{B}_1 + \cdots + \mathcal{B}_p + x)^n.$$

Possible Extension to Nörlund Polynomials

$$\left(\frac{t}{e^t - 1}\right)^p e^{xt} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} \Leftrightarrow B_n^{(p)}(x) = (\mathcal{B}_1 + \cdots + \mathcal{B}_p + x)^n.$$

$$\frac{\Gamma(z+x)}{\Gamma(z+x+1-p)z^p} \sim \sum_{n=0}^{\infty} \frac{(p-n)_n}{n!} B_n^{(p)}(x) \frac{1}{z^{n+1}}$$

where $(a)_n = a(a+1) \cdots (a+n-1)$.

Possible Extension to Nörlund Polynomials

$$\left(\frac{t}{e^t - 1}\right)^p e^{xt} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} \Leftrightarrow B_n^{(p)}(x) = (\mathcal{B}_1 + \cdots + \mathcal{B}_p + x)^n.$$

$$\frac{\Gamma(z+x)}{\Gamma(z+x+1-p)z^p} \sim \sum_{n=0}^{\infty} \frac{(p-n)_n}{n!} B_n^{(p)}(x) \frac{1}{z^{n+1}}$$

where $(a)_n = a(a+1)\cdots(a+n-1)$.

$$\log(\mathcal{B}_1 + \cdots + \mathcal{B}_p + z) = -H_{p-1} + \frac{d^{p-1}}{dz^{p-1}} \left[\binom{z-1}{p-1} \psi\left(z - \lfloor \frac{p}{2} \rfloor\right) \right]$$

where $H_n := 1 + 1/2 + \cdots + 1/n$, is the n -th harmonic number and $\lfloor \cdot \rfloor$ is the floor function.

End

Thank you