

I am an experimental mathematician working mainly in **number theory**, also intersecting with **combinatorics**, **probability**, and **differential geometry**. Computer software, programming languages, especially certain algorithms play an essential role in my research, by providing calculations, simulations, visualizations, and etc., for empirical evidence of conjectures. Primarily applying probabilistic and symbolic methods, I focus on certain *numbers, sequences, polynomials, special functions, combinatorial identities, geometric structures, algorithms*, and their properties that computer algorithms can (in-)directly guide or help to prove.

In the next sections, I will briefly describe current and future projects.

1 Probabilistic Approach

Large portion of my research topics, methods, and background involve probability: either by probabilistic approach, or some direct results in probability.

1.1 The Bernoulli and Euler Symbol

For all the special functions and polynomials that are either important, or have deep connections with other fields of mathematics, **Bernoulli and Euler polynomials** (e.g., [10, Chpt. 24]) are among my favorite ones. Some early results are summarized in the second part of my Ph. D thesis [Jiu16]. The main tools here are the *Bernoulli* and *Euler symbols*, denoted by \mathcal{B} and \mathcal{E} , respectively. Each of the two symbols satisfies a simple evaluation rule: $\mathcal{B}^n = B_n$ and $\mathcal{E}^n = E_n/2$, where B_n and E_n are *Bernoulli* and *Euler numbers*. Originally, both \mathcal{B} and \mathcal{E} arise from the traditional umbral calculus (see, e.g., [12]). Meanwhile, the probabilistic interpretations (See, e.g., [4, Thm. 2.3] and [JMV14, Prop. 2.1]) view both symbols as random variables:

$$\mathcal{B} = iL_B - 1/2 \quad \text{and} \quad \mathcal{E} = iL_E - 1/2, \quad (1)$$

where $i^2 = -1$, and two random variables L_B and L_E have density functions p_B and p_E , respectively, as follows.

$$p_B(t) := \pi \operatorname{sech}^2(\pi t)/2 \quad \text{and} \quad p_E(t) := \operatorname{sech}(\pi t). \quad (2)$$

Note that, (1) implies the evaluation rules are exactly calculating the expectation. This probabilistic setup not only provides a rigorous background, also largely extends the application of symbolic computations.

1.1.1 Extensions

As random variables, we can consider the sum of independent and identically distributed (i.i.d.) sequence

$$\mathcal{B}^{(p)} := \mathcal{B}_1 + \cdots + \mathcal{B}_p \quad \text{and} \quad \mathcal{E}^{(p)} := \mathcal{E}_1 + \cdots + \mathcal{E}_p,$$

where $\mathcal{B}_i \sim \mathcal{B}$ and similarly, $\mathcal{E}_i \sim \mathcal{E}$. Then, extensions on higher-order polynomials and Bernoulli-Barnes polynomials also admit symbolic expressions, which do not appear in traditional umbral calculus.

- Bernoulli and Euler polynomials of order p , $B_n^{(p)}(x)$ and $E_n^{(p)}(x)$, are given by

$$B_n^{(p)}(x) = \mathbb{E} \left[\left(x + \mathcal{B}^{(p)} \right)^n \right] \quad \text{and} \quad E_n^{(p)}(x) = \mathbb{E} \left[\left(x + \mathcal{E}^{(p)} \right)^n \right]; \quad (3)$$

- and Bernoulli-Barnes polynomials $B_n(\mathbf{a}; x)$, with $\mathbf{a} = (a_1, \dots, a_k)$ and $a_l \neq 0$ can be expressed as

$$B_n(\mathbf{a}; x) = \mathbb{E} \left[\frac{(x + a_1 \mathcal{B}_1 + \cdots + a_k \mathcal{B}_k)^n}{a_1 a_2 \cdots a_k} \right], \quad (4)$$

where $\mathcal{B} = (\mathcal{B}_1, \dots, \mathcal{B}_k)$ and $\mathbf{a} \cdot \mathcal{B} = \sum_{l=1}^k a_l \mathcal{B}_l$.

Results based on symbolic expressions extends traditional ones. For instance, In [JMV16, Thm. 2.2], we extended a difference formula by Bayad and Beck [2, Thm. 5.1], symbolically, that for any polynomial P ,

$$P(x - \mathbf{a} \cdot \mathcal{B}) = \sum_{j=0}^n \sum_{|J|=j} |a|_{J^*} P^{(n-j)}(x + (\mathbf{a} \cdot \mathcal{B})_J),$$

where $J \subset [n] := \{1, \dots, n\}$ and $J^* = [n] \setminus J$.

While in [JMV14], not only did we apply the random variable interpretation to establish and prove identities, an expected observation (see [JMV14, Note 4.8]) triggers more probabilistic models involving objects in combinatorics and number theory, especially for finding and proving identities involving Bernoulli and Euler polynomials with their higher-order extensions. This will be stated in Subsection 1.2.1.

1.1.2 Orthogonal polynomials, Hankel determinants, and continued fractions

Admittedly, one does not need the probabilistic background, e.g., random variable, probability measure, moments, etc., to study orthogonal polynomials, Hankel determinants, continued fractions, weighted lattice path, generalized Motzkin number, and other related topics. Comprehensive introduction on basics and facts can be found, e.g., in [5, 7]. Personally, it is the random variable interpretation of $E_n^{(p)}(x)$, Euler polynomials of higher-order defined in (3), leading me to study the corresponding orthogonal polynomials, denoted by $\Omega_n^{(p)}(y)$. In [JS19b], we recognize that $\Omega_n^{(p)}(y) = i^n n! P_n^{(p/2)}(-i(y - x + \frac{p}{2}); \frac{\pi}{2}) / 2^n$, as the Meixner-Pollaczek polynomial $P_n^{(\lambda)}(y; \phi)$ (; see also [8, eq. 9.7.1]). Similarly, let $\varrho_n(y)$ be the monic orthogonal polynomials with respect to the Bernoulli polynomial $B_n(x)$, then we identify that $\varrho_n(y) = n! p_n(y; \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}) / (n+1)_n$, where $p_n(y; a, b, c, d)$ is the continuous Hahn polynomial [8, pp. 200–202]. This work opens the door for me to continue on Hankel determinants, especially of sequences related to Bernoulli and Euler polynomials, due to the following two reasons.

1. The classical formula for calculating the monic orthogonal polynomials of a given sequence contains the Hankel determinants (, see, e.g., [7, Eq. 2.1.6]);
2. Lemma 1 in [JS19b] actually provides an alternative proof, in the aspect of random variables and orthogonal polynomials, that Hankel determinants are invariant under binomial transforms [9, Item 445]. More precisely, the sequence c_n and its binomial transformed polynomial $c_n(x) := \sum_{k=0}^n \binom{n}{k} c_k x^{n-k}$ share the same Hankel determinants.

Therefore, in recent years, I accomplished a series studies on Hankel determinants [DJ21, DJ22, DJ23, JL, CJS, CJLW], involving mainly Bernoulli and Euler polynomials, with their extensions, in, e.g., Dirichlet characters and q -analogues. Besides major computations of new and important Hankel determinants, other highlights include the following.

- An incomplete table of Hankel determinant identities for numerous sequences containing Bernoulli and Euler numbers and polynomials was collected at the end of [DJ22].
- New general formulas of Hankel determinants of right-shifted sequence [DJ23, Lem. 2.5] and derivatives of Apell sequence [DJ22, Thm. 5.1] are proven.
- The Hankel determinants we studied in [JL] originally comes from statistically estimating the variance in non-parametric regression. Also the symbolic computation, or more precisely WZ method, was used and applied in the proofs.
- In the latest work [CJLW], we study the q -binomial transform sequence and its Hankel determinants, with partial result on the degree of polynomials and the leading coefficients. And some proofs make fully use of the Bernoulli and Euler symbols in (1).

1.2 Probabilistic Models

In [JMV14], based on the probabilistic interpretation, we obtained, for any positive integer N ,

$$E_n(x) = \frac{1}{N^n} \sum_{l=N}^{\infty} p_l^{(N)} E_n^{(l)} \left(\frac{l-N}{2} + Nx \right),$$

where the coefficients $p_l^{(N)}$ appear in the series expansion of the reciprocal of the N th Chebychev polynomial of the first kind T_N

$$\frac{1}{T_N(1/z)} = \sum_{\ell=0}^{\infty} p_{\ell}^{(N)} z^{\ell}.$$

Moreover, $p_{\ell}^{(N)}$ can also be viewed as transition probabilities in the context of a random walk over a finite number of sites [JMV14, Note 4.8], based on which, we continued to explore more of such probabilistic models.

1.2.1 Random walk, Brownian motion, and Bessel process

We consider both 1-dimensional reflected Brownian motion and 3-dimensional Bessel process. More specifically, we decompose the hitting times for consecutive level sites. For instance, considering in the figure below: starting from site a_0 , the hitting time to the site a_{m+1} can be decomposed, combinatorially, by the hitting times among sites a_0, a_1, \dots, a_m in between, where the back and forth walks between neighboring sites form loops, denoted by L_1, L_2, \dots, L_m .




Figure 1: Loop Decomposition

With the combinatorial loop and hitting time decomposition, we can further derive non-trivial identities, involving $B_n^{(p)}(x)$ and $E_n^{(p)}(x)$. In the early work [JV19], results on only 1 or 2 loops, i.e. $m = 1$ or 2 in Figure 1, were obtained. Later in [JSY22], we successfully extended to the general n loops, by both induction and a combinatorial interpretation. Further identities are derived from equally distributed level sites. For instance, the 3-loop cases in 3-dim Bessel process implies

$$B_{n+1}\left(\frac{x+2}{5}\right) - B_{n+1}\left(\frac{x}{5}\right) = \frac{n+1}{5^n} \sum_{k=0}^{\infty} \frac{3^k}{2^{2k+3}} \sum_{\ell=0}^k \binom{k}{\ell} (-1)^\ell \frac{1}{12^\ell} E_n^{(2k+2\ell+3)}(k+\ell+x).$$

1.2.2 Shuffling model

The very recent work [CJS] studies the uniform shuffling on n cards, i.e., each result has probability $1/n!$. Then, merge the maximal consecutive integer subsequences and reduce the number of cards into the current remaining one. The model was introduced by Rao et al. [11] to model the number of times that catalysts must be added to n molecules to bond into a single lump. The molecules have a given hierarchical order which led to the above mathematical formulation of the process. They studied the number of permutations needed for the process to end, X_n ; and obtained the asymptotic behavior of the mean as

$$n \leq \mathbb{E}[X_n] \leq n + \sqrt{n} \Rightarrow \mathbb{E}[X_n] \sim n.$$

Our work [CJS] not only improved it by

$$\mathbb{E}[X_n] = n + \left(1 + \frac{1}{2} + \cdots + \frac{1}{n-1}\right) + \varepsilon_n, \quad \text{with } 0 \leq \varepsilon_n - \varepsilon_{n+1} \leq \frac{1}{n^2}, \quad (5)$$

which also indicates that ε_n has a limit. More importantly, the simulation in [11], for X_2, \dots, X_{100} hinted that the asymptotic distribution of X_n would tend to be normal. While, we completely solve it in the affirmative, i.e., we proved that

$$\frac{X_n - n}{\sqrt{n}} \xrightarrow{w} \mathcal{N}(0, 1),$$

by estimating all the central moments of X_n . As a by product, certain limits related to the sequence involve the Bell numbers.

1.3 Differential and Information Geometry

I studied differential and information geometry for my master's degree, which can be seen from some of my early work, e.g., [WJ06, JS07, PSJ07, ZSJP13, ZSJP14, LZJS16]. Switching to studying experimental mathematics in my Ph. D program does not prevent me from continuing my research in information geometry; in fact, I am gradually combining some work together.

- In both [TJKZ20] and [JK20], the Wishart distribution plays the essential rule, in defining the zonal polynomials and hence hypergeometric functions with matrix arguments. Alternatively, an equivalent definition of the zonal polynomials is the eigenfunction of the Laplace–Beltrami operator on the space of symmetric, positive definite matrices.
- I contributed to an awarded joint grant “Wuhan University—Duke Kunshan University—University of Minnesota, Twin Cities Joint Research Platform”, together with my colleague Dr. Dongmian Zou.
- In the recent work [JP], we use the information geometry tools to study the exponential generalized beta of the second kind (EGB2) distribution or the beta-logistic distribution, with density function

$$f(x; \beta_1, \beta_2) = \frac{1}{B(\beta_1, \beta_2)} \cdot \frac{\exp(\beta_1 x)}{(1 + \exp(x))^{\beta_1 + \beta_2}},$$

where $\beta_1, \beta_2 > 0$ and $B(x, y)$ is the beta function. With some simplification, it is not hard to recognize that this family include both densities for Bernoulli and Euler symbols (2). Besides basic geometric structures, we also uncovers that the beta-logistic distribution admits an α -parallel prior for any real number α , that has the potential for application in geometric statistical inference.

2 Symbolic Approach

In general, computer or machine proofs, as the major aim in symbolic computation, always contribute to mathematical proofs. For example, the WZ-method is applied in the proofs of [DJMV15, JK20, JL]. Meanwhile, Bernoulli and Euler symbols, mentioned above, can also be considered as generalized symbolic computation method.

2.1 Multiple zeta value at non-positive integers

The multiple zeta function

$$\zeta_r(n_1, \dots, n_r) = \sum_{k_1, \dots, k_r > 0} \frac{1}{k_1^{n_1} (k_1 + k_2)^{n_2} \dots (k_1 + \dots + k_r)^{n_r}}$$

has more than one analytic continuations at non-positive integers. For instance, Sadaoui [13, Thm. 1] used the Raabe's identity while Akiyama and Yanigawa [1, p. 350] considered the Euler-Maclaurin summation formula. Since both results involve Bernoulli number, applying Bernoulli symbol reveals, to our surprise, that both analytic continuations coincide. More precisely, for non-negative integers n_1, \dots, n_k , we have, symbolically,

$$\zeta_r(-n_1, \dots, -n_r) = \prod_{k=1}^r (-1)^{n_k} \mathcal{C}_{1, \dots, k}^{n_k+1} \quad (6)$$

where $\mathcal{C}_1^n = \mathcal{B}_1^n/n$ and recursively, $\mathcal{C}_{1, \dots, k+1} = (\mathcal{C}_{1, \dots, k} + \mathcal{B}_{k+1})^n/n$. This shows the two approaches, by Raabe's identity and Euler-Maclaurin summation formula that lead to analytic continuations of MZVs, coincide on non-positive integer values. Other results such as recurrence [JMV18, Thm. 3.1], contiguity identities [JMV18, Thm. 4.1], and generating functions [JMV18, Thm. 5.1] follow naturally from (6).

Also in [JWV20], we further symbolically expressed the r -fold harmonic sums at negative indices, which is similar to the multiple zeta value expression. For example [JWV20, Thm. 3.1], by defining the \mathcal{H} symbol as

$$(\mathcal{H}(N))^n = H_{-n}(N) = 1^n + 2^n + \dots + (N-1)^n,$$

the r -fold multiple power sums can be expressed as

$$H_{-n_1, \dots, -n_r}(N) := \sum_{N > i_1 > \dots > i_r > 0} i_1^{n_1} \dots i_r^{n_r} = \prod_{k=1}^r \mathcal{H}_{1, \dots, k}^{n_k},$$

where $\mathcal{H}_1 = \mathcal{H}(N)$ and recursively $\mathcal{H}_{1, \dots, k} = \mathcal{H}(\mathcal{H}_{1, \dots, k-1})$ for $k = 2, 3, \dots, r$. This is compatible with (6) by letting $N \rightarrow \infty$.

2.2 The method of brackets

The method of bracket is an efficient method for the evaluation of a large class of definite integrals on the half-line, i.e.,

$$\int_0^\infty f(x) dx,$$

with only 6 simple rules, with applications to evaluate certain Feynman integrals, which arise from Feynman diagrams. In theoretical physics, a Feynman diagram is a pictorial representation of the mathematical expressions describing the behavior and interaction of subatomic particles. It is also related to Dr. Moll, my Ph. D supervisor's long term project in proving all entries of [6]. Some other work of this project include [ADG⁺16a]

Some early work and summary can be found in the first half of my Ph. D thesis [Jiu16]. Further generalization, novel evaluation techniques, and related discussion can be found, e.g., in [GJM16, GKJM17, GKJM18, GKJM20, BGJ⁺23]. For instance, in [GJM16], we first observed several examples, all missing a common factor of 2. This is

due to the analytic continuation of the Pochhammer symbol $(a)_k = \Gamma(a+k)/\Gamma(a)$, when both a and k are negative integers. In order to get the correct answer, the evaluation

$$(-km)_{-m} = \frac{k}{k+1} \frac{(-1)^m (km)!}{((k+1)m)!},$$

should be applied.

3 Miscellaneous and final remark

Please find the publication list below, including some other work that does not fit into my major research projects. Among them, some are also suitable for undergraduate students, e.g., [BJMV14, JV16, JS22, JS].

List of Publications

- [ADG⁺14] Tewodros Amdeberhan, Atul Dixit, Xiao Guan, Lin Jiu, and Victor H. Moll, *The unimodality of a polynomial coming from a rational integral. back to the original proof*, Journal of Mathematical Analysis and Applications **420** (2014), no. 5, 1154–1166.
- [ADG⁺16a] Tewodros Amdeberhan, Atul Dixit, Xiao Guan, Lin Jiu, Alexey Kuznetsov, Victor H. Moll, and Christophe Vignat, *The integrals in gradshiteyn and ryzhik. part 30: trigonometric functions*, Scientia Series A: Mathematical Sciences **27** (2016), 47–74.
- [ADG⁺16b] Tewodros Amdeberhan, Atul Dixit, Xiao Guan, Lin Jiu, Victor H. Moll, and Christophe Vignat, *A series involving catalan numbers. proofs and demonstrations*, Elemente der Mathematik **71** (2016), no. 3, 109–121.
- [BGJ⁺23] Zachary Bradshaw, Ivan Gonzalez, Lin Jiu, Victor H. Moll, and Christophe Vignat, *Compatibility of the method of brackets with classical integration rules*, Open Mathematics **21** (2023), no. 1, Article 20220581.
- [BJMV14] Alyssa Byrnes, Lin Jiu, Victor H. Moll, and Christophe Vignat, *Recursion rules for the hypergeometric zeta functions*, International Journal of Number Theory **10** (2014), no. 7, 1761–1782.
- [CJ24] Shane Chern and Lin Jiu, *Hankel determinants and jacobi continued fractions for q-euler number*, Comptes Rendus Mathématique **362** (2024), 203–216.
- [CJLW] Shane Chern, Lin Jiu, Shuhan Li, and Liuquan Wang, *Leading coefficient in the hankel determinants related to binomial and q-binomial transforms*, under submission.
- [CJS] Shane Chern, Lin Jiu, and Italo Simonelli, *A central limit theorem for a card shuffling problem*, under submission.
- [DJ21] Karl Dicher and Lin Jiu, *Orthogonal polynomials and hankel determinants for certain bernoulli and euler polynomials*, Journal of Mathematical Analysis and Applications **497** (2021), no. 1, Article 124855.
- [DJ22] ———, *Hankel determinants of sequences related to bernoulli and euler polynomials*, International Journal of Number Theory **18** (2022), no. 2, 331–359.
- [DJ23] ———, *Hankel determinants of shifted sequences of bernoulli and euler numbers*, Contributions to Discrete Mathematics **18** (2023), no. 2, 146–175.
- [DJMV15] Atul Dixit, Lin Jiu, Victor H. Moll, and Christophe Vignat, *The finite fourier transform of classical polynomials*, Journal of the Australian Mathematical Society **98** (2015), no. 2, 145–160.
- [GJM16] Ivan Gonzalez, Lin Jiu, and Victor H. Moll, *Pochhammer symbol with negative indices. a new rule for the method of brackets*, Open Mathematics **14** (2016), no. 1, 681–686.
- [GKJM17] Ivan Gonzalez, Karen Kohl, Lin Jiu, and Victor H. Moll, *An extension of the method of brackets. part 1*, Open Mathematics **15** (2017), no. 1, 1181–1211.
- [GKJM18] ———, *The method of brackets in experimental mathematics*, ch. 16, pp. 307–318, World Scientific Publishers, 2018.
- [GKJM20] ———, *An extension of the method of brackets. part 2*, Open Mathematics **18** (2020), no. 1, 983–955.
- [Jiu16] Lin Jiu, *The method of brackets and the bernoulli symbol*, Ph.D. thesis, New Orleans, LA, USA, 2016.
- [Jiu17] ———, *Integral representations of equally positive integer-indexed harmonic sums at infinity*, Research in Number Theory **3** (2017), Article 10.
- [JK20] Lin Jiu and Christoph Koutschan, *Calculation and properties of zonal polynomials*, Mathematics in Computer Science **14** (2020), 623–640.
- [JL] Lin Jiu and Ye Li, *Hankel determinants of certain sequences of bernoulli polynomials: A direct proof of an inverse matrix entry from statistics*, Contributions to Discrete Mathematics, to appear.

- [JMV14] Lin Jiu, Victor H. Moll, and Christophe Vignat, *Identities for generalized euler polynomials*, Integral Transforms and Special Functions **25** (2014), no. 10, 777–789.
- [JMV16] ———, *A symbolic approach to some identities for bernoulli-barnes polynomials*, International Journal of Number Theory **12** (2016), no. 3, 649–662.
- [JMV18] ———, *A symbolic approach to multiple zeta values at the negative integers*, Journal of Symbolic Computation **84** (2018), 1–13.
- [JP] Lin Jiu and Linyu Peng, *Information geometry and α -parallel prior of the beta-logistic distribution*, under submission.
- [JS] Lin Jiu and Diane Yahui Shi, *On b-ary binomial coefficients with negative entries*, under submission.
- [JS07] Lin Jiu and Huafei Sun, *On minimal homothetical hypersurfaces*, Colloquium Mathematicum **109** (2007), no. 2, 239–249.
- [JS19a] Lin Jiu and Diane Yahui Shi, *Matrix representation for multiplicative nested sums*, Colloquium Mathematicum **158** (2019), no. 2, 183–194.
- [JS19b] ———, *Orthogonal polynomials and connection to generalized motzkin numbers for higher-order euler polynomials*, Journal of Number Theory **199** (2019), 389–402.
- [JS22] ———, *Moments and cumulants on identities for bernoulli and euler numbers*, Mathematical Reports **24** (2022), no. 4, 643–650.
- [JSY22] Lin Jiu, Italo Simonelli, and Heng Yue, *Loop decompositions of random walks and nontrivial identities of bernoulli and euler polynomials*, INTEGERS **22** (2022), Article 91.
- [JV16] Lin Jiu and Christophe Vignat, *On binomial identities in arbitrary bases*, Journal of Integer Sequences **19** (2016), no. 5, Article 16.5.5.
- [JV19] ———, *Connection coefficients for higher-order bernoulli and euler polynomials: a random walk approach*, Fibonacci Quarterly **57** (2019), no. 5, 84–95.
- [JVV20] Lin Jiu, Christophe Vignat, and Tanay Wakhare, *Analytic continuation for multiple zeta values using symbolic representations*, International Journal of Number Theory **16** (2020), no. 3, 579–602.
- [LZJS16] Chunhui Li, Erchuan Zhang, Lin Jiu, and Huafei Sun, *Optimal control on special euclidean group via natural gradient descent algorithm*, Science China Information Sciences **59** (2016), no. 11, Article 112203.
- [PSJ07] Linyu Peng, Huafei Sun, and Lin Jiu, *The geometric structure of the pareto distribution*, Boletín de la Asociación Matemática Venezolana **14** (2007), no. 1, 5–13.
- [TJKZ20] Nobuki Takayama, Lin Jiu, Satoshi Kuriki, and Yi Zhang, *Computation of the expected euler characteristic for the largest eigenvalue of a real non-central wishart matrix*, Journal of Multivariate Analysis **179** (2020), Article 104642.
- [WJ06] Xiaojie Wang and Lin Jiu, *Characterizing hypersurfaces of generalized rotation through its normal lines*, Journal of Ningde Normal University (Natural Science) **02** (2006), 117–119.
- [ZSJP13] Fengyun Zhang, Huafei Sun, Lin Jiu, and Linyu Peng, *The arc length variational formula on the exponential manifold*, Mathematica Slovaca **63** (2013), no. 5, 1101–1112.
- [ZSJP14] Zhengning Zhang, Huafei Sun, Lin Jiu, and Linyu Peng, *A natural gradient algorithm for stochastic distribution systems*, Entropy **18** (2014), no. 8, 4338–4352.

References

- [1] S. Akiyama and Y. Tanigawa, Multiple zeta values at non-positive integers, *Ramanujan J.*, **5** (2001), 327–351.
- [2] A. Bayad and M. Beck, Relations for Bernoulli-Barnes numbers and Barnes zeta functions, *Int. J. Number Theory*, **10** (2014), 1321–1335.
- [3] T. Budd, Winding of simple walks on the square lattice, *J. Combin. Theory Ser. A*, 172(2020) Article 105191.
- [4] A. Dixit, V. H. Moll, and C. Vignat, The Zagier modification of Bernoulli numbers and a polynomial extension. Part I, *Ramanujan J.*, **33** (2014), 379–422.
- [5] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge Univ. Press, 2010.
- [6] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Academic Press, 2015.
- [7] M. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, Cambridge Univ. Press, 2005.
- [8] R. Koekoek, P. A. Lesky, and R. F. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their q -Analogues*, Springer, 2010.
- [9] T. Muir and W. H. Metzler, *A Treatise on the Theory of Determinants*, Dover Publications, 1960.
- [10] F. W. J. Oliver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, (eds.), *NIST Handbook of Mathematical Functions*, Cambridge Univ. Press, 2010.

- [11] M. Rao, H. Zhang, C. Huang, and F.-C. Cheng, A discrete probability problem in card shuffling, *Comm. Statist. Theory Methods*, **45** (2016), , 612–620.
- [12] S. Roman, *The Umbral Calculus*, Academic Press Inc., 1984.
- [13] B. Sadaoui, Multiple zeta values at the non-positive integers, *C. R. Acad. Sci. Paris, Ser. 1* **12** (2014), 977–984.