

# Matrix Representations for Bernoulli and Euler Polynomials

Lin JIU (Joint work with Diane Yahui Shi)

Department of Mathematics and Statistics, Dalhousie University  
2018 CMS Summer Meeting @ University of New Brunswick

June 3rd, 2018

# Objects

# Objects

Bernoulli polynomials:

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}.$$

# Objects

Bernoulli polynomials:

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}.$$

Euler polynomials:

$$\frac{2e^{zx}}{e^z + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{z^n}{n!}.$$

# Objects

Bernoulli polynomials:

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}.$$

Euler polynomials:

$$\frac{2e^{zx}}{e^z + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{z^n}{n!}.$$

Bernoulli numbers:  $B_n = B_n(0)$

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \qquad B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k.$$

# Objects

Bernoulli polynomials:

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}.$$

Euler polynomials:

$$\frac{2e^{zx}}{e^z + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{z^n}{n!}.$$

Bernoulli numbers:  $B_n = B_n(0)$

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \qquad B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k.$$

$n$	0	1	2	3	4
$B_n$	1	$-\frac{1}{2}$	$\frac{1}{6}$	0	$-\frac{1}{30}$
$B_n(x)$	1	$x - \frac{1}{2}$	$x^2 - x + \frac{1}{6}$	$x^3 - \frac{3}{2}x^2 + \frac{1}{2}x$	$x^4 - 2x^3 + x^2 - \frac{1}{30}$

# Objects

Bernoulli polynomials:

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}.$$

Euler polynomials:

$$\frac{2e^{zx}}{e^z + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{z^n}{n!}.$$

Bernoulli numbers:  $B_n = B_n(0)$

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k.$$

$n$	0	1	2	3	4
$B_n$	1	$-\frac{1}{2}$	$\frac{1}{6}$	0	$-\frac{1}{30}$
$B_n(x)$	1	$x - \frac{1}{2}$	$x^2 - x + \frac{1}{6}$	$x^3 - \frac{3}{2}x^2 + \frac{1}{2}x$	$x^4 - 2x^3 + x^2 - \frac{1}{30}$

$$B_n^+ := \begin{cases} B_n, & \text{if } n \neq 1; \\ 1/2, & \text{if } n = 1. \end{cases}$$

# Objects

Bernoulli polynomials:

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}.$$

Euler polynomials:

$$\frac{2e^{zx}}{e^z + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{z^n}{n!}.$$

Bernoulli numbers:  $B_n = B_n(0)$

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k.$$

$n$	0	1	2	3	4
$B_n$	1	$-\frac{1}{2}$	$\frac{1}{6}$	0	$-\frac{1}{30}$
$B_n(x)$	1	$x - \frac{1}{2}$	$x^2 - x + \frac{1}{6}$	$x^3 - \frac{3}{2}x^2 + \frac{1}{2}x$	$x^4 - 2x^3 + x^2 - \frac{1}{30}$

$$B_n^+ := \begin{cases} B_n, & \text{if } n \neq 1; \\ 1/2, & \text{if } n = 1. \end{cases} \quad B_n^+ = B_n(1).$$



# Matrix & Determinants

# Matrix & Determinants

$$B_n^+ = \frac{\det(a_{i,j})_{n \times n}}{n!}, \quad a_{i,j} := \begin{cases} 0, & \text{if } j > i + 1; \\ \binom{i+1}{j-1}, & \text{otherwise.} \end{cases}$$

# Matrix & Determinants

$$B_n^+ = \frac{\det(a_{i,j})_{n \times n}}{n!}, \quad a_{i,j} := \begin{cases} 0, & \text{if } j > i + 1; \\ \binom{i+1}{j-1}, & \text{otherwise.} \end{cases}$$

$$B_3 = B_3^+ = \frac{\det \begin{pmatrix} 1 & 2 & 0 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{pmatrix}}{3!} = 0.$$

# Matrix & Determinants

$$B_n^+ = \frac{\det(a_{i,j})_{n \times n}}{n!}, \quad a_{i,j} := \begin{cases} 0, & \text{if } j > i + 1; \\ \binom{i+1}{j-1}, & \text{otherwise.} \end{cases}$$

$$B_3 = B_3^+ = \frac{\det \begin{pmatrix} 1 & 2 & 0 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{pmatrix}}{3!} = 0.$$

$$B_n = (-1)^n n! \det \begin{pmatrix} \frac{1}{2!} & 1 & 0 & 0 & \cdots & 0 \\ \frac{1}{3!} & \frac{1}{2!} & 1 & 0 & \cdots & 0 \\ \frac{1}{4!} & \frac{1}{3!} & \frac{1}{2!} & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{1}{n!} & \frac{1}{(n-1)!} & \frac{1}{(n-2)!} & \frac{1}{(n-3)!} & \cdots & 1 \\ \frac{1}{(n+1)!} & \frac{1}{n!} & \frac{1}{(n-1)!} & \frac{1}{(n-2)!} & \cdots & \frac{1}{2!} \end{pmatrix}$$

# Matrix Representation

# Matrix Representation

$$R := \begin{pmatrix} x - \frac{1}{2} & \omega_1 & 0 & 0 & \cdots & \cdots \\ 1 & x - \frac{1}{2} & \omega_2 & 0 & \cdots & \cdots \\ 0 & 1 & x - \frac{1}{2} & \ddots & \ddots & \cdots \\ 0 & 0 & 1 & \ddots & \omega_n & \cdots \\ 0 & 0 & 0 & \ddots & x - \frac{1}{2} & \cdots \\ 0 & 0 & 0 & \ddots & 1 & \ddots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \text{ where } \omega_n = -\frac{n^4}{4(2n+1)(2n-1)}.$$

# Matrix Representation

$$R := \begin{pmatrix} x - \frac{1}{2} & \omega_1 & 0 & 0 & \cdots & \cdots \\ 1 & x - \frac{1}{2} & \omega_2 & 0 & \cdots & \cdots \\ 0 & 1 & x - \frac{1}{2} & \ddots & \ddots & \cdots \\ 0 & 0 & 1 & \ddots & \omega_n & \cdots \\ 0 & 0 & 0 & \ddots & x - \frac{1}{2} & \cdots \\ 0 & 0 & 0 & \ddots & 1 & \ddots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \text{ where } \omega_n = -\frac{n^4}{4(2n+1)(2n-1)}.$$

For example,

$$R_4 := \begin{pmatrix} x - 1/2 & -1/12 & 0 & 0 \\ 1 & x - 1/2 & -4/15 & 0 \\ 0 & 1 & x - 1/2 & -81/140 \\ 0 & 0 & 1 & x - 1/2 \end{pmatrix}$$

# Matrix Representation

$$R := \begin{pmatrix} x - \frac{1}{2} & \omega_1 & 0 & 0 & \cdots & \cdots \\ 1 & x - \frac{1}{2} & \omega_2 & 0 & \cdots & \cdots \\ 0 & 1 & x - \frac{1}{2} & \ddots & \ddots & \cdots \\ 0 & 0 & 1 & \ddots & \omega_n & \cdots \\ 0 & 0 & 0 & \ddots & x - \frac{1}{2} & \cdots \\ 0 & 0 & 0 & \ddots & 1 & \ddots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \text{ where } \omega_n = -\frac{n^4}{4(2n+1)(2n-1)}.$$

For example,

$$R_4 := \begin{pmatrix} x - 1/2 & -1/12 & 0 & 0 \\ 1 & x - 1/2 & -4/15 & 0 \\ 0 & 1 & x - 1/2 & -81/140 \\ 0 & 0 & 1 & x - 1/2 \end{pmatrix}$$

$$(R_4)^4 = \begin{pmatrix} x^4 - 2x^3 + x^2 - \frac{1}{30} & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{pmatrix}$$



# Matrix Representation

$$R := \begin{pmatrix} x - \frac{1}{2} & \omega_1 & 0 & 0 & \cdots & \cdots \\ 1 & x - \frac{1}{2} & \omega_2 & 0 & \cdots & \cdots \\ 0 & 1 & x - \frac{1}{2} & \ddots & \ddots & \cdots \\ 0 & 0 & 1 & \ddots & \omega_n & \cdots \\ 0 & 0 & 0 & \ddots & x - \frac{1}{2} & \cdots \\ 0 & 0 & 0 & \ddots & 1 & \ddots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \text{ where } \omega_n = -\frac{n^4}{4(2n+1)(2n-1)}.$$

For example,

$$R_4 := \begin{pmatrix} x - 1/2 & -1/12 & 0 & 0 \\ 1 & x - 1/2 & -4/15 & 0 \\ 0 & 1 & x - 1/2 & -81/140 \\ 0 & 0 & 1 & x - 1/2 \end{pmatrix}$$

$$(R_4)^4 = \begin{pmatrix} x^4 - 2x^3 + x^2 - \frac{1}{30} & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{pmatrix}$$

If letting  $\omega_n = -n^2/4$ , we could similarly obtain  $E_n(x)$ .

# Main Result

## Theorem [L. J and D. Shi]

Define  $M_{n,k}$  by  $M_{0,0} = 1$  and for  $n > 0$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1},$$

where  $\omega_k = -\frac{k^4}{4(2k+1)(2k-1)}$  and  $M_{n,k} = 0$  if  $k > n$ . Then,  $M_{n,0} = B_n(x)$ .

# Main Result

## Theorem [L. J and D. Shi]

Define  $M_{n,k}$  by  $M_{0,0} = 1$  and for  $n > 0$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1},$$

where  $\omega_k = -\frac{k^4}{4(2k+1)(2k-1)}$  and  $M_{n,k} = 0$  if  $k > n$ . Then,  $M_{n,0} = B_n(x)$ .

$$R := \begin{pmatrix} x - \frac{1}{2} & \omega_1 & 0 & 0 & \cdots & \cdots \\ 1 & x - \frac{1}{2} & \omega_2 & 0 & \cdots & \cdots \\ 0 & 1 & x - \frac{1}{2} & \ddots & \ddots & \cdots \\ 0 & 0 & 1 & \ddots & \omega_n & \cdots \\ 0 & 0 & 0 & \ddots & x - \frac{1}{2} & \cdots \\ 0 & 0 & 0 & \ddots & 1 & \ddots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ ,

# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ , i.e.,

$$m_n = \mathbb{E}[X^n] = \int_{\mathbb{R}} t^n p(t) dt.$$

# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ , i.e.,

$$m_n = \mathbb{E}[X^n] = \int_{\mathbb{R}} t^n p(t) dt.$$

Let  $P_n(y)$  be the monic orthogonal polynomials with respect to  $X$  (or w. r. t.  $m_n$ ),

# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ , i.e.,

$$m_n = \mathbb{E}[X^n] = \int_{\mathbb{R}} t^n p(t) dt.$$

Let  $P_n(y)$  be the monic orthogonal polynomials with respect to  $X$  (or w. r. t.  $m_n$ ), i.e.,  $\deg P_n = n$ ,

# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ , i.e.,

$$m_n = \mathbb{E}[X^n] = \int_{\mathbb{R}} t^n p(t) dt.$$

Let  $P_n(y)$  be the monic orthogonal polynomials with respect to  $X$  (or w. r. t.  $m_n$ ), i.e.,  $\deg P_n = n$ ,  $\text{LC}[P_n] = 1$ ,



# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ , i.e.,

$$m_n = \mathbb{E}[X^n] = \int_{\mathbb{R}} t^n p(t) dt.$$

Let  $P_n(y)$  be the monic orthogonal polynomials with respect to  $X$  (or w. r. t.  $m_n$ ), i.e.,  $\deg P_n = n$ ,  $\text{LC}[P_n] = 1$ , and

$$\int_{\mathbb{R}} P_m(t) P_n(t) p(t) dt = c_n \delta_{m,n} = \begin{cases} c_n, & \text{if } m = n; \\ 0, & \text{otherwise.} \end{cases}$$

# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ , i.e.,

$$m_n = \mathbb{E}[X^n] = \int_{\mathbb{R}} t^n p(t) dt.$$

Let  $P_n(y)$  be the monic orthogonal polynomials with respect to  $X$  (or w. r. t.  $m_n$ ), i.e.,  $\deg P_n = n$ ,  $\text{LC}[P_n] = 1$ , and

$$\int_{\mathbb{R}} P_m(t) P_n(t) p(t) dt = c_n \delta_{m,n} = \begin{cases} c_n, & \text{if } m = n; \\ 0, & \text{otherwise.} \end{cases}$$

$$c_n \delta_{m,n} = P_m(t) P_n(t) \Big|_{t^k = m_k}.$$

# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ , i.e.,

$$m_n = \mathbb{E}[X^n] = \int_{\mathbb{R}} t^n p(t) dt.$$

Let  $P_n(y)$  be the monic orthogonal polynomials with respect to  $X$  (or w. r. t.  $m_n$ ), i.e.,  $\deg P_n = n$ ,  $\text{LC}[P_n] = 1$ , and

$$\int_{\mathbb{R}} P_m(t) P_n(t) p(t) dt = c_n \delta_{m,n} = \begin{cases} c_n, & \text{if } m = n; \\ 0, & \text{otherwise.} \end{cases}$$

$$c_n \delta_{m,n} = P_m(t) P_n(t) \Big|_{t^k = m_k}.$$

$P_n$  satisfies a three-term recurrence: for some sequences  $(s_n)_{n \geq 0}$  and  $(t_n)_{n \geq 1}$ ,

$$P_{n+1}(y) = (y - s_n) P_n(y) - t_n P_{n-1}(y).$$

# Random Variable

Let  $X$  be a random variable with density function  $p(t)$  on  $\mathbb{R}$  and with moments  $m_n$ , i.e.,

$$m_n = \mathbb{E}[X^n] = \int_{\mathbb{R}} t^n p(t) dt.$$

Let  $P_n(y)$  be the monic orthogonal polynomials with respect to  $X$  (or w. r. t.  $m_n$ ), i.e.,  $\deg P_n = n$ ,  $\text{LC}[P_n] = 1$ , and

$$\int_{\mathbb{R}} P_m(t) P_n(t) p(t) dt = c_n \delta_{m,n} = \begin{cases} c_n, & \text{if } m = n; \\ 0, & \text{otherwise.} \end{cases}$$

$$c_n \delta_{m,n} = P_m(t) P_n(t) \Big|_{t^k = m_k}.$$

$P_n$  satisfies a three-term recurrence: for some sequences  $(s_n)_{n \geq 0}$  and  $(t_n)_{n \geq 1}$ ,

$$P_{n+1}(y) = (y - s_n) P_n(y) - t_n P_{n-1}(y).$$

## Theorem


$$\sum_{n=0}^{\infty} m_n x^n = \frac{m_0}{1 - s_0 x - \frac{t_1 x^2}{1 - s_1 x - \frac{t_2 x^2}{\ddots}}}.$$

# Generalized Motzkin Numbers

$$M_{n+1,k} = M_{n,k-1} + \sigma_k M_{n,k} + \tau_{k+1} M_{n,k+1}$$


# Generalized Motzkin Numbers

$$M_{n+1,k} = M_{n,k-1} + \sigma_k M_{n,k} + \tau_{k+1} M_{n,k+1}$$




# Generalized Motzkin Numbers

$$M_{n+1,k} = M_{n,k-1} + \sigma_k M_{n,k} + \tau_{k+1} M_{n,k+1}$$



# Generalized Motzkin Numbers

$$M_{n+1,k} = M_{n,k-1} + \sigma_k M_{n,k} + \tau_{k+1} M_{n,k+1}$$



$$\sum_{n=0}^{\infty} M_{n,0} z^n = \frac{1}{1 - s_0 z - \frac{t_1 z^2}{1 - s_1 z - \frac{t_2 z^2}{\dots}}}$$



# General Identification

## Theorem

For random variable  $X$  with moments  $m_n$  and monic orthogonal polynomials  $P_n$ , satisfying recurrence

$$P_{n+1}(y) = (y - s_n)P_n(y) - t_n P_{n-1}(y),$$

we define generalized Motzkin numbers  $M_{n,k}$  by letting  $(\sigma_k, \tau_k) = (s_k, t_k)$ . If further assuming  $m_0 = 1$ , we have

$$M_{n,0} = m_n = \mathbb{E}[X^n].$$

# General Identification

## Theorem

For random variable  $X$  with moments  $m_n$  and monic orthogonal polynomials  $P_n$ , satisfying recurrence

$$P_{n+1}(y) = (y - s_n)P_n(y) - t_n P_{n-1}(y),$$

we define generalized Motzkin numbers  $M_{n,k}$  by letting  $(\sigma_k, \tau_k) = (s_k, t_k)$ . If further assuming  $m_0 = 1$ , we have

$$M_{n,0} = m_n = \mathbb{E}[X^n].$$

$$M_{n+1,k} = M_{n,k-1} + s_k M_{n,k} + t_{k+1} M_{n,k+1}$$

# General Identification

## Theorem

For random variable  $X$  with moments  $m_n$  and monic orthogonal polynomials  $P_n$ , satisfying recurrence

$$P_{n+1}(y) = (y - s_n)P_n(y) - t_n P_{n-1}(y),$$

we define generalized Motzkin numbers  $M_{n,k}$  by letting  $(\sigma_k, \tau_k) = (s_k, t_k)$ . If further assuming  $m_0 = 1$ , we have

$$M_{n,0} = m_n = \mathbb{E}[X^n].$$

$$M_{n+1,k} = M_{n,k-1} + s_k M_{n,k} + t_{k+1} M_{n,k+1}$$

$$M_{n+1,0} = s_0 M_{n,0} + t_1 M_{n,1}.$$

# Probabilistic Interpretation of $B_n(x)$

## Probabilistic Interpretation of $B_n(x)$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1} \Rightarrow M_{n,0} = B_n(x).$$

# Probabilistic Interpretation of $B_n(x)$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1} \Rightarrow M_{n,0} = B_n(x).$$

[Question1] Are  $B_n(x)$  moments of some random variable?

# Probabilistic Interpretation of $B_n(x)$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1} \Rightarrow M_{n,0} = B_n(x).$$

[Question1] Are  $B_n(x)$  moments of some random variable?

[Answer] Let  $L_B \sim \pi \operatorname{sech}^2(\pi t)/2$ , then

$$B_n(x) = \frac{\pi}{2} \int_{\mathbb{R}} \left(x + it - \frac{1}{2}\right)^n \operatorname{sech}^2(\pi t) dt = \mathbb{E} \left[ \left(iL_B + x - \frac{1}{2}\right)^n \right].$$

# Probabilistic Interpretation of $B_n(x)$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1} \Rightarrow M_{n,0} = B_n(x).$$

[Question1] Are  $B_n(x)$  moments of some random variable?

[Answer] Let  $L_B \sim \pi \operatorname{sech}^2(\pi t)/2$ , then

$$B_n(x) = \frac{\pi}{2} \int_{\mathbb{R}} \left(x + it - \frac{1}{2}\right)^n \operatorname{sech}^2(\pi t) dt = \mathbb{E} \left[ \left(iL_B + x - \frac{1}{2}\right)^n \right].$$

[Question2] What are the monic orthogonal polynomials w. r. t.  $B_n(x)$ ?



# Probabilistic Interpretation of $B_n(x)$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1} \Rightarrow M_{n,0} = B_n(x).$$

[Question1] Are  $B_n(x)$  moments of some random variable?

[Answer] Let  $L_B \sim \pi \operatorname{sech}^2(\pi t)/2$ , then

$$B_n(x) = \frac{\pi}{2} \int_{\mathbb{R}} \left(x + it - \frac{1}{2}\right)^n \operatorname{sech}^2(\pi t) dt = \mathbb{E} \left[ \left(iL_B + x - \frac{1}{2}\right)^n \right].$$

[Question2] What are the monic orthogonal polynomials w. r. t.  $B_n(x)$ ?

[Answer] Well...

# Probabilistic Interpretation of $B_n(x)$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1} \Rightarrow M_{n,0} = B_n(x).$$

[Question1] Are  $B_n(x)$  moments of some random variable?

[Answer] Let  $L_B \sim \pi \operatorname{sech}^2(\pi t)/2$ , then

$$B_n(x) = \frac{\pi}{2} \int_{\mathbb{R}} \left(x + it - \frac{1}{2}\right)^n \operatorname{sech}^2(\pi t) dt = \mathbb{E} \left[ \left(iL_B + x - \frac{1}{2}\right)^n \right].$$

[Question2] What are the monic orthogonal polynomials w. r. t.  $B_n(x)$ ?

[Answer] Well...

- ▶ 1st observation:  $B_n = \mathbb{E} \left[ \left(iL_B - \frac{1}{2}\right)^n \right]$

# Probabilistic Interpretation of $B_n(x)$

$$M_{n+1,k} = M_{n,k-1} + \left(x - \frac{1}{2}\right) M_{n,k} + \omega_{k+1} M_{n,k+1} \Rightarrow M_{n,0} = B_n(x).$$

[Question1] Are  $B_n(x)$  moments of some random variable?

[Answer] Let  $L_B \sim \pi \operatorname{sech}^2(\pi t)/2$ , then

$$B_n(x) = \frac{\pi}{2} \int_{\mathbb{R}} \left(x + it - \frac{1}{2}\right)^n \operatorname{sech}^2(\pi t) dt = \mathbb{E} \left[ \left(iL_B + x - \frac{1}{2}\right)^n \right].$$

[Question2] What are the monic orthogonal polynomials w. r. t.  $B_n(x)$ ?

[Answer] Well...

- ▶ 1st observation:  $B_n = \mathbb{E} \left[ \left(iL_B - \frac{1}{2}\right)^n \right] = B_n(0).$

# One More Setup

## Theorem (J. Touchard)

Let  $\phi_n$  be the monic orthogonal polynomials w. r. t.  $B_n$ . Then, they satisfy

$$\phi_{n+1}(y) = \left(y + \frac{1}{2}\right) \phi_n(y) - \omega_n \phi_{n-1}(y)$$

# One More Setup

## Theorem (J. Touchard)

Let  $\phi_n$  be the monic orthogonal polynomials w. r. t.  $B_n$ . Then, they satisfy

$$\phi_{n+1}(y) = \left(y + \frac{1}{2}\right) \phi_n(y) - \omega_n \phi_{n-1}(y) \quad \left[(s_n, t_n) = \left(\frac{1}{2}, \omega_n\right)\right].$$

# One More Setup

## Theorem (J. Touchard)

Let  $\phi_n$  be the monic orthogonal polynomials w. r. t.  $B_n$ . Then, they satisfy

$$\phi_{n+1}(y) = \left(y + \frac{1}{2}\right) \phi_n(y) - \omega_n \phi_{n-1}(y) \quad \left[(s_n, t_n) = \left(\frac{1}{2}, \omega_n\right)\right].$$

## Lemma[L. J and D. Shi]

	Moments	Monic Orthogonal Polynomials
$X$	$m_n$	$P_{n+1}(y) = (y - s_n)P_n(y) - t_n P_{n-1}(y)$
$X + c$	$\sum_{k=0}^n \binom{n}{k} m_k c^{n-k}$	$Q_{n+1}(y) = (y - s_n - c)Q_n(y) - t_n Q_{n-1}(y)$
$CX$	$C^n m_n$	$R_{n+1}(y) = (y - Cs_n)R_n(y) - C^2 t_n R_{n-1}(y)$

# One More Setup

## Theorem (J. Touchard)

Let  $\phi_n$  be the monic orthogonal polynomials w. r. t.  $B_n$ . Then, they satisfy

$$\phi_{n+1}(y) = \left(y + \frac{1}{2}\right) \phi_n(y) - \omega_n \phi_{n-1}(y) \quad \left[(s_n, t_n) = \left(\frac{1}{2}, \omega_n\right)\right].$$

## Lemma[L. J and D. Shi]

	Moments	Monic Orthogonal Polynomials
$X$	$m_n$	$P_{n+1}(y) = (y - s_n)P_n(y) - t_n P_{n-1}(y)$
$X + c$	$\sum_{k=0}^n \binom{n}{k} m_k c^{n-k}$	$Q_{n+1}(y) = (y - s_n - c)Q_n(y) - t_n Q_{n-1}(y)$
$CX$	$C^n m_n$	$R_{n+1}(y) = (y - Cs_n)R_n(y) - C^2 t_n R_{n-1}(y)$

Recall

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_n x^{n-k}.$$

# Conclusion

For Bernoulli polynomials  $B_n(x)$ , the corresponding orthogonal polynomials  $\theta_n(y)$  satisfy

$$\theta_{n+1}(y) = \left( y - \left( x - \frac{1}{2} \right) \right) \theta_n(y) - \omega_n \theta_{n-1}(y).$$



# Conclusion

For Bernoulli polynomials  $B_n(x)$ , the corresponding orthogonal polynomials  $\theta_n(y)$  satisfy

$$\theta_{n+1}(y) = \left( y - \left( x - \frac{1}{2} \right) \right) \theta_n(y) - \omega_n \theta_{n-1}(y).$$

Thus, we define the generalized Motzkin numbers by

$$M_{n+1,k} = M_{n,k-1} + \left( x - \frac{1}{2} \right) M_{n,k} + \omega_{k+1} M_{n,k+1},$$

# Conclusion

For Bernoulli polynomials  $B_n(x)$ , the corresponding orthogonal polynomials  $\theta_n(y)$  satisfy

$$\theta_{n+1}(y) = \left( y - \left( x - \frac{1}{2} \right) \right) \theta_n(y) - \omega_n \theta_{n-1}(y).$$

Thus, we define the generalized Motzkin numbers by

$$M_{n+1,k} = M_{n,k-1} + \left( x - \frac{1}{2} \right) M_{n,k} + \omega_{k+1} M_{n,k+1},$$

which implies  $M_{n,0} = B_n(x)$ . The matrix presentation follows from the lattice path interpretation.

# Euler Polynomials

Euler numbers  $(E_n)_{n=0}^{\infty}$  and Euler polynomials  $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

# Euler Polynomials

Euler numbers  $(E_n)_{n=0}^{\infty}$  and Euler polynomials  $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

In addition,

$$E_n(x) = \int_{\mathbb{R}} \left( x - \frac{1}{2} + it \right)^n \operatorname{sech}(\pi t) dt.$$

# Euler Polynomials

Euler numbers  $(E_n)_{n=0}^{\infty}$  and Euler polynomials  $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

In addition,

$$E_n(x) = \int_{\mathbb{R}} \left( x - \frac{1}{2} + it \right)^n \operatorname{sech}(\pi t) dt.$$

$$E_n = 2^n E_n\left(\frac{1}{2}\right)$$

# Euler Polynomials

Euler numbers  $(E_n)_{n=0}^{\infty}$  and Euler polynomials  $(E_n(x))_{n=0}^{\infty}$

$$\operatorname{sech}(t) = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

In addition,

$$E_n(x) = \int_{\mathbb{R}} \left( x - \frac{1}{2} + it \right)^n \operatorname{sech}(\pi t) dt.$$

$$E_n = 2^n E_n\left(\frac{1}{2}\right)$$


$$M_{n+1,k} = M_{n,k-1} + \left( x - \frac{1}{2} \right) M_{n,k} - \frac{(k+1)^2}{4} M_{n,k+1}$$

# Euler Polynomials

$$E_n = 2^n E_n \left( \frac{1}{2} \right) \Rightarrow M_{n+1,k} = M_{n,k-1} - \frac{(k+1)^2}{4} M_{n,k+1}$$

# Euler Polynomials


$$E_n = 2^n E_n \left( \frac{1}{2} \right) \Rightarrow M_{n+1,k} = M_{n,k-1} - \frac{(k+1)^2}{4} M_{n,k+1}$$





# Euler Polynomials


$$E_n = 2^n E_n \left( \frac{1}{2} \right) \Rightarrow M_{n+1,k} = M_{n,k-1} - \frac{(k+1)^2}{4} M_{n,k+1}$$



$n = 6, E_6$

# Euler Polynomials

$$E_n = 2^n E_n \left( \frac{1}{2} \right) \Rightarrow M_{n+1,k} = M_{n,k-1} - \frac{(k+1)^2}{4} M_{n,k+1}$$




$n = 6$ ,  $E_6$  Dyck paths

$$C_3 = \frac{1}{4} \binom{6}{3} = \frac{6 \cdot 5 \cdot 4}{4 \cdot 3 \cdot 2 \cdot 1} = 5.$$

# Euler Polynomials

$$E_n = 2^n E_n \left( \frac{1}{2} \right) \Rightarrow M_{n+1,k} = M_{n,k-1} - \frac{(k+1)^2}{4} M_{n,k+1}$$



$n = 6$ ,  $E_6$  Dyck paths

$$C_3 = \frac{1}{4} \binom{6}{3} = \frac{6 \cdot 5 \cdot 4}{4 \cdot 3 \cdot 2 \cdot 1} = 5.$$

$$-61 = E_6 = (-1)^3 (3^2 2^2 1^2 + 2^2 2^2 1^2 + 1^2 2^2 1^2 + 2^2 1^2 1^2 + 1^2 1^2 1^2)$$

# Current Work

- ▶ Dyck paths  $\Rightarrow (E_{2n}, C_n)$ ;

# Currenct Work

- ▶ Dyck paths  $\Rightarrow (E_{2n}, C_n)$ ;
- ▶ Nörlund polynomials :

$$\left(\frac{t}{e^t - 1}\right)^p e^{xt} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!};$$

# Current Work

- ▶ Dyck paths  $\Rightarrow (E_{2n}, C_n)$ ;
- ▶ Nörlund polynomials :

$$\left(\frac{t}{e^t - 1}\right)^p e^{xt} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!};$$

$$(R_4)^4 \begin{pmatrix} x^4 - 2x^3 + x^2 - \frac{1}{30} & -\frac{1}{3}x^3 + \frac{1}{2}x^2 - \frac{2}{15}x - \frac{1}{135} & -\frac{16}{15}x^3 + \frac{14}{15}x^2 - \frac{2}{3}x - \frac{1}{15} \\ 4x^3 - 6x^2 + \frac{8}{5}x + \frac{4}{45} & x^4 - 2x^3 - \frac{3}{5}x^2 + \frac{52}{45}x - \frac{37}{3780} & -\frac{16}{15}x^3 + \frac{14}{15}x^2 - \frac{2}{3}x - \frac{1}{15} \\ 6x^2 - \frac{13}{3}x - \frac{89}{1008} & 4x^3 - \frac{7}{2}x^2 - \frac{635}{252}x + \frac{1673}{1728} & -\frac{16}{15}x^3 + \frac{14}{15}x^2 - \frac{2}{3}x - \frac{1}{15} \\ 4x - \frac{19}{12} & 6x^2 - \frac{13}{3}x - \frac{89}{1008} & 4x^3 - \frac{7}{2}x^2 - \frac{635}{252}x + \frac{1673}{1728} \end{pmatrix}$$

End

Thank you