

# Research Statement

Lin JIU

My current research interests are Experiment Mathematics and Symbolic Computation, involving Special Functions, Combinatorics, Number Theory, Probability Theory, and Information Geometry. The following sections describe topics in details, where terms with ★ are obtained results while that with ● refer to future work. (Some items contain both.)

## 1 The Bernoulli symbol

### 1.1 Introduction

The Bernoulli symbol  $\mathcal{B}$  is a random variable  $\mathcal{B} = \iota L_B - \frac{1}{2}$ , where  $\iota^2 = -1$  and  $L_B$  has probability density function

$$p_{L_B}(t) = \frac{\pi}{2} \operatorname{sech}^2(\pi t) \text{ on } \mathbb{R}.$$

The Bernoulli numbers  $(B_n)_{n=0}^\infty$  are defined and also can be computed in terms of moments [11, Thm. 2.3, pp. 384] as follows

$$\sum_{n=0}^{\infty} B_n \frac{t^n}{n!} = \frac{t}{e^t - 1} \text{ and } B_n = \mathbb{E}[\mathcal{B}^n] = \frac{\pi}{2} \int_{\mathbb{R}} \left( \iota t - \frac{1}{2} \right)^2 \operatorname{sech}^2(\pi t) dt, \quad (1.1)$$

which interprets the evaluation (eval) of umbral symbol [32] as the expectation:  $\operatorname{eval}(\mathcal{B}^n) = B_n = \mathbb{E}[\mathcal{B}^n]$ . Therefore,

1. properties of umbral calculus are preserved, e.g., Bernoulli polynomials  $B_n(x)$  are defined and symbolically expressed as

$$\sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} = \frac{te^{xt}}{e^t - 1} \Rightarrow B_n(x) = \mathbb{E}[(\mathcal{B} + x)^n] = \frac{\pi}{2} \int_{\mathbb{R}} \left( x + \iota t - \frac{1}{2} \right)^2 \operatorname{sech}^2(\pi t) dt;$$

2. formulae can be simplified into more compact form:

$$\sum_{k=1}^n k^m = \frac{1}{m+1} \sum_{l=0}^m \binom{m+1}{l} B_l n^{m+1-l} = \frac{1}{m+1} \sum_{l=0}^m \binom{m+1}{l} \mathcal{B}^l n^{m+1-l} = \int_0^n (\mathcal{B} + x)^m dx;$$

3. formulae can also be visualized more directly, e.g., a direct calculus fact reveals the derivative of  $B_n(x)$  as

$$B'_n(x) = B_{n-1}(x) \Leftrightarrow [(\mathcal{B} + x)^n]' = n(\mathcal{B} + x)^{n-1}.$$

More than traditional umbral calculus, Bernoulli symbol admits probabilistic technique to further explore as follows.

1. Conjugate variables, e.g.,  $X$  and  $Y$ , are defined as

$$\mathbb{E}[(X + Y)^n] = \delta_{0,n}, \text{ or equivalently, } \mathbb{E}(e^{tX}) \cdot \mathbb{E}(e^{tY}) = 1.$$

Easily, the uniform random variable (or uniform symbol)  $\mathcal{U} \sim U[0, 1]$  and  $\mathcal{B}$  are conjugate, since

$$\mathbb{E}(e^{t\mathcal{U}}) = \int_0^1 e^{tu} \cdot 1 du = \frac{e^t - 1}{t} \text{ and } \mathbb{E}(e^{t\mathcal{B}}) = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} = \frac{t}{e^t - 1}.$$

Therefore, consider the case that  $f(x) = x^n$ . Since  $\mathcal{B}$  and  $\mathcal{U}$  cancel each other in powers,

$$f(x) = f(x + \mathcal{B} + \mathcal{U}) = \int_0^1 f(x + \mathcal{B} + u) du \Rightarrow B_n(x + 1) - B_n(x) = nx^{n-1}.$$

2. Moment generating functions of independent random variables are the product of individual ones, i.e.,

$$\mathbb{E}[e^{t(X_1 + \dots + X_n)}] = \mathbb{E}[e^{tX_1}] \dots \mathbb{E}[e^{tX_n}].$$

Thus, Nörlund polynomials  $B_n^{(p)}(x)$  are defined and symbolically expressed as

$$\sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} = e^{tx} \left( \frac{t}{e^t - 1} \right)^p \Rightarrow B_n^{(p)}(x) = (\mathcal{B}_1 + \dots + \mathcal{B}_p + x)^n,$$

where  $\{\mathcal{B}_k\}_{k=1}^p$  is a sequence of independent and identical distribution (i. i. d. s. ), with  $\mathcal{B}_k \sim \mathcal{B}$  (throughout this statement, where  $p$  may vary).

## 1.2 Main results

- ★ The Bernoulli-Barnes polynomials  $B_n(\mathbf{a}; x)$ , where  $\mathbf{a} = (a_1, \dots, a_k)$  satisfies  $|\mathbf{a}| = \prod_{l=1}^k a_l \neq 0$ , is defined and symbolically expressed [25, eq. 1.17, pp. 651] as, by letting  $\mathcal{B} = (\mathcal{B}_1, \dots, \mathcal{B}_k)$ :

$$e^{tx} \prod_{i=1}^k \frac{t}{e^{a_i t} - 1} = \sum_{n=0}^{\infty} B_n(\mathbf{a}; x) \frac{t^n}{n!} \Rightarrow B_n(\mathbf{a}; x) = \frac{1}{|\mathbf{a}|} (x + \mathbf{a} \cdot \mathcal{B})^n, \text{ where } \mathbf{a} \cdot \mathcal{B} = \sum_{l=1}^k a_l \mathcal{B}_l.$$

Based on this symbolic expression, we have,

1. for polynomial  $P$ , [25, Thm. 2.2, pp. 652]

$$P(x - \mathbf{a} \cdot \mathcal{B}) = \sum_{j=0}^n \sum_{|J|=j} |a|_{J^*} P^{(n-j)}(x + (\mathbf{a} \cdot \mathcal{B})_J), \text{ where } J \subset [n] := \{1, \dots, n\} \text{ and } J^* = [n] \setminus J,$$

which, by taking  $P(x) = \frac{x^m}{|\mathbf{a}|^m}$ , gives the difference formula [6, Thm. 5.1]:

$$(-1)^m B_m(-x; \mathbf{a}) - B_m(x; \mathbf{a}) = m! \sum_{j=0}^{n-1} \sum_{|J|=j} \frac{B_{m-n+j}(x; \mathbf{a}_J)}{(m-n+j)!};$$

2. for  $A := a_1 + \dots + a_n \neq 0$ , direct proofs of self-duality [25, Thm. 5.1, pp. 656] of sequence  $((-\frac{1}{A})^n B_n(\mathbf{a}))_{n=0}^{\infty}$  and the symmetry formula [25, Thm. 5.2, pp. 656]

$$(-1)^m \sum_{k=0}^m \binom{m}{k} A^{m-k} B_{l+k}(x; \mathbf{a}) = (-1)^l \sum_{k=0}^l \binom{l}{k} A^{l-k} B_{m+k}(-x; \mathbf{a}),$$

which Bayad and Beck [6] are looking for.

- ★ The multiple zeta values (MZV),

$$\zeta_r(n_1, \dots, n_r) = \sum_{k_1, \dots, k_r > 0} \frac{1}{k_1^{n_1} (k_1 + k_2)^{n_2} \dots (k_1 + \dots + k_r)^{n_r}}$$

have more than one analytic continuations at non-positive integers. For instance, Sadaoui [33, Thm. 1] used the Raabe's identity while Akiyama and Yanigawa [2, pp. 350] considered the Euler-Maclaurin summation. Since both results involve Bernoulli number, applying Bernoulli symbol reveals, to our surprise, that both analytic continuations coincide, symbolically expressed as [24, Thm. 2.1], for  $n_1, \dots, n_k \in \mathbb{Z}_- \cup \{0\}$ ,

$$\zeta_r(n_1, \dots, n_r) = \prod_{k=1}^r (-1)^{n_k} \mathcal{C}_{1, \dots, k}^{n_k+1}, \text{ where recursively } \mathcal{C}_1^n = \frac{(\mathcal{B}_1)^n}{n}, \mathcal{C}_{1, \dots, k+1} = \frac{(\mathcal{C}_{1, \dots, k} + \mathcal{B}_{k+1})^n}{n},$$

Recurrence [24, Thm. 3.1]

$$\zeta_r(-n_1, \dots, -n_r) = \frac{(-1)^{n_r}}{n_r + 1} \sum_{k=0}^{n_r+1} \binom{n_r+1}{k} (-1)^k \zeta_{r-1}(-n_1, \dots, -n_{r-2}, -n_{r-1} - k) B_{n_r+1-k},$$

contiguity identities [24, Thm. 4.1] and generating functions [24, Thm. 5.1] follow naturally.

## 1.3 Future work

- Explanation for the coincidence of two analytic continuations mentioned above;
- Application to other types of zeta functions, e.g., Witten-zeta function, where Bernoulli number are also involved;
- Further probabilistic study on Bernoulli symbol in the probabilistic approach;
- ★ Symbolic expressions and results on hypergeometric Bernoulli numbers [9, eq. 1.13, pp. 1763], polynomials [9, eq. 7.8, pp. 1778] and hypergeometric zeta functions [9, eq. 1.12, pp. 1763];
- ★ Analogue on Euler numbers, denoted by  $E_n$ , which similarly satisfies [26, Prop. 2.1, pp. 778]:

$$E_n = \int_{\mathbb{R}} \left( \iota t - \frac{1}{2} \right)^2 \operatorname{sech}(\pi t) dt. \quad (1.2)$$

## 2 Matrix presentation of harmonic sums and multiplicative nested sums

### 2.1 Introduction

Nested sums are given by  $\mathcal{S}(f; k; N) := \sum_{N \geq n_1 \geq \dots \geq n_k \geq 1} f(n_1, \dots, n_k)$ , and we focus on the special case that  $f(n_1, \dots, n_k) = f_1(n_1) \dots f_k(n_k)$ , called *multiplicative nested sums*. A large class of multiple nested sums is the harmonic sums [8, eq. 4, pp. 1]: for indices  $a_1, \dots, a_k \in \mathbb{R} \setminus \{0\}$ ,

$$S_{a_1, \dots, a_k}(N) = \sum_{N \geq n_1 \geq \dots \geq n_k \geq 1} \frac{\text{sign}(a_1)^{n_1}}{n_1^{|a_1|}} \times \dots \times \frac{\text{sign}(a_k)^{n_k}}{n_k^{|a_k|}}, \quad N \in \mathbb{N}, \quad (2.1)$$

which connect to a variety of topics, including combinatorics, quantum field theory, and naturally zeta functions, e.g., taking  $N \rightarrow \infty$ ,  $k = 1$  and  $a_1 > 0$ :  $S_{a_1}(\infty) = \zeta(a_1)$ . Instead of the recurrence [7, eq. 2.1, pp. 21] that inherited from the quasi-shuffle relation [21, eq. 1, pp. 51], which the `Mathematica` package `HarmonicSums.m`<sup>1</sup> [1, Chpt. 6] uses, we associate each factor  $f_l$ ,  $l = 1, \dots, k$ , an *index matrix*:

$$\mathcal{P}_{N|f_l} := \begin{pmatrix} f_l(1) & 0 & \dots & 0 \\ f_l(2) & f_l(2) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ f_l(N) & f_l(N) & \dots & f_l(N) \end{pmatrix}$$

to provide alternative computations and interpretations.

### 2.2 Main results

★ If additionally define  $f_0(x) = \frac{1}{x}$ , then [23, Thm. 15]:

$$\mathcal{S}(f; k; N) = N \cdot \left( \prod_{l=0}^k \mathcal{P}_{N|f_l} \right)_{N,1}, \quad (2.2)$$

the  $N^{\text{th}}$  row and  $1^{\text{st}}$  column entry of the product matrix. Taking  $f_l(n_l) = \text{sign}(a_l)^{n_l} / n_l^{|a_l|}$ , the harmonic sums admit the same result [23, Thm. 1], which is more direct than a recurrence.

- ★ For the special case of harmonic sums when  $a_1 = \dots = a_k = a \geq 1$  and  $N < \infty$ , a random walk over finite number of sites [23, Sec. 3] interprets the index matrix as the stochastic matrix and the harmonic sums as probabilities of certain paths.
- ★ When all  $f_l$ 's are identical, the product (2.2) becomes powers. In particular, if  $\{f_l(1), \dots, f_l(N)\}$  are all distinct, then  $\mathcal{P}_{N|f_l}$  is diagonalizable, so that

$$\mathcal{P}_{N|f_l} = \mathcal{Q}_{N|f_l} \text{diag}\{f_l(1), \dots, f_l(N)\} \mathcal{Q}_{N|f_l}^{-1} \Rightarrow \mathcal{P}_{N|f_l}^m = \mathcal{Q}_{N|f_l} \text{diag}\{f_l(1)^m, \dots, f_l(N)^m\} \mathcal{Q}_{N|f_l}^{-1},$$

the alternative computation for its powers lead to combinatorial identities:

1.  $f_1(x) = x$  and  $k = 1$  gives binomial-type identity:

$$\sum_{l=1}^N (-1)^{N-l} l^{N+1} \binom{N}{l} = \frac{N(N+1)!}{2};$$

2.  $f_1 \equiv \dots \equiv f_k = f$  with  $f(m) = a_m$  leads to [35, eq. 2, pp. 313]:

$$\sum_{N \geq n_1 \geq \dots \geq n_k \geq 1} a_{n_1} \dots a_{n_k} = \sum_{j=1}^N \left( \prod_{\substack{m=1 \\ m \neq j}}^N \frac{1}{1 - \frac{a_m}{a_j}} \right) a_j^k,$$

which, when  $a_j = \frac{a-bq^{j+i-1}}{c-zq^{j+i-1}}$  and  $N = n-i+1$ , “turns out to be a common source of several  $q$ -identities” [35, pp. 314];

3. special harmonic sum (2.1) when  $a_1 = \dots = a_k = 1$  recovers [10, Cor. 3, pp. 93]:

$$\sum_{l=1}^N (-1)^{l-1} \binom{N}{l} \frac{1}{l^k} = \sum_{N \geq n_1 \geq \dots \geq n_k \geq 1} \frac{1}{n_1 \dots n_k}.$$

### 2.3 Future work

- Matrix interpretation of the quasi-shuffle relation, i.e., the recurrence of harmonic sums;
- Modification of the index matrices to visualize the limit  $N \rightarrow \infty$ ;
- Systematic algorithms for finding and proving combinatorial identities through index matrices.

<sup>1</sup><http://www.risc.jku.at/research/combinat/software/HarmonicSums/index.php>

### 3 The Method of Brackets (MoB)

#### 3.1 Introduction

The *method of brackets* (MoB), developed by I. Gonzalez [16, 17], has its origin on the evaluation of definite integrals arising from the Schwinger parametrization of Feynman diagrams. Besides examples of its application appearing in [5, 12, 13, 14, 15], software implementation has been produced by K. Kohl in [27] using Sage with internal use of Mathematica. In concrete, MoB evaluates the integral  $\int_0^\infty f(x) dx$ , by the following rules:

Rule  $P_1$ :

$$f(x) = \sum_{n=0}^{\infty} a_n x^{\alpha n + \beta - 1} \Rightarrow \int_0^\infty f(x) dx = \sum_n a_n \langle \alpha n + \beta \rangle.$$

Rule  $P_2$ : For  $\alpha \notin \mathbb{N} \cup \{0\}$ ,

$$(a_1 + \dots + a_r)^\alpha = \sum_{m_1, \dots, m_r} \phi_{m_1, \dots, m_r} a_1^{m_1} \dots a_r^{m_r} \frac{\langle -\alpha + a_1 + \dots + a_r \rangle}{\Gamma(-\alpha)},$$

where  $\phi_{m_1, \dots, m_r} = \prod_{i=1}^r \phi_{m_i}$  is the product of *indicators* defined by  $\phi_n := (-1)^n / \Gamma(n+1)$ .

Rule  $E_1$ :

$$\sum_n \phi_n f(n) \langle \alpha n + \beta \rangle = \frac{1}{|\alpha|} f(n^*) \Gamma(-n^*), \text{ where } n^* \text{ solves } \alpha n + \beta = 0$$

Rule  $E_2$ :

$$\sum_{n_1, \dots, n_r} \phi_{n_1, \dots, n_r} \prod_{i=1}^r \langle a_{i1} n_1 + \dots + a_{ir} n_r + c_i \rangle = f(n_1^*, \dots, n_r^*) \prod_{i=1}^r \Gamma(-n_i^*).$$

where  $(n_1^*, \dots, n_r^*)$  solves  $A(n_1, \dots, n_r)^T + (c_1, \dots, c_r)^T = \vec{0}^T$  for non-singular matrix  $A = (a_{ij})_{r \times r}$ .

Rule  $E_3$ : Each representation of an integral by a bracket series has associated an index as

$$\text{index} = \text{number of sums} - \text{number of brackets}.$$

The value of a multi-dimensional bracket series of positive index is obtained by computing all the contributions of maximal rank by Rule  $E_2$ . These contributions to the integral appear as series in the free parameters. Series converging in a common region are added and divergent series are discarded. Any series producing a non-real contribution is discarded too.

Rule  $E_4$ : Fix  $k \in \mathbb{N}$ , and for  $m \in \mathbb{N}$ , Pochhammer symbols with negative index and negative integer base are evaluated as

$$(-km)_{-m} = \frac{k}{k+1} \cdot \frac{(-1)^m (km)!}{((k+1)m)!} \quad (3.1)$$

Basic results and examples can be found in [12, 13, 14, 15].

#### 3.2 Main results

- ★ The rule  $E_4$  was recently introduced [18, eq. 27, pp. 685] from the analytic continuation of Pochhammer symbol with negative integer index and negative integer base, while calculating entry **6.671.7** in [19].
- ★ We verify that this method does not depend on factorization of the integrand [22, Thm. 4.2.2, pp. 24], i.e., if  $f(x) = \prod_{i=1}^r g_i(x)$ , applying MoB to compute  $\int_0^\infty f(x) dx$  and  $\int_0^\infty \prod_{i=1}^r g_i(x) dx$  leads to the same results.
- ★ We compared MoB with the negative dimension integration method [22, Sec. 8.1], showing two methods are completely different; and also with the integrating by differentiating method [22, Sec. 8.2], revealing their formal connections.
- ★ A pure Mathematica package [22, Chpt. 9] is developing and current version recovers examples in [14, 15].

#### 3.3 Future work

- Further study on Ramanujan's master theorem [20, pp. 186, eq. B], which is the funding theorem of MoB;
- Application of the `Sigma.m`<sup>2</sup> package developed by Schneider [34] to the obtained series from MoB;
- Development of a purely Sage package.

---

<sup>2</sup><http://www.risc.jku.at/research/combinat/software/Sigma/index.php>

## 4 Differential Geometry and Information Geometry

### 4.1 Introduction

In order to apply differential geometry on probability and statistics, Amari [3, 4] initiated the theory of information geometry working over statistical manifold

$$S := \{p(x; \theta) \mid \theta \in \Theta \subseteq \mathbb{R}^n \text{ and } p \text{ is a probability density function.}\}$$

A series of concepts such as  $\alpha$ -connections, dual connections and Fisher metrics are consequently introduced and studied. In particular, with the help of the Fisher information matrix, which is defined as

$$g_{ij}(\theta) := \mathbb{E}[(\partial_i l)(\partial_j l)] = \int (\partial_i l)(\partial_j l) p(x; \theta) dx, \quad i, j = 1, 2, \dots, n, \text{ where } \partial_i := \frac{\partial}{\partial \theta_i} \text{ and } l := \log p(x; \theta),$$

the geometric structure is constructed and also highly connected to information theory. A remarkable result which motivated Amari is that the sectional curvature of the manifold consisting of normal distributions is  $-\frac{1}{2}$ , the hyperbolic geometry.

Currently, information geometry can be well applied to exponential family (including normal distributions), whose the probability density function  $p(x)$  can be expressed in terms of functions  $\{C, F_1, \dots, F_n\}$  and a convex function  $\phi$  on  $\Theta$  as:

$$p(x; \theta) = \exp \left\{ C(x) + \sum_i \theta_i F_i(x) - \phi(\theta) \right\}.$$

### 4.2 Main results

- ★ Both 1<sup>st</sup> [36, Thm. 1.1, pp. 1102] and 2<sup>nd</sup> [36, Thm. 1.2, 1.3, pp. 1102–1103] arc length variational formulae are given.
- ★ The  $\alpha$ -structure of frame bundles [30, Thm. 5.8] is obtained.
- ★ A list of possible holonomy groups [31, Thm. 5.3] of exponential family is calculated.
- ★ A natural gradient algorithm [37, Thm. 1, pp. 4342] is introduced to design the controller of an open-loop stochastic distribution control system (SDCS) of multi-input and single output.
- ★ An algorithm [29, Algorithm 1, pp. 6] for optimal control on  $SE(n)$  is studied and simulated.
- ★ While considering the geometric means of matrices for the trace class operator, the Rényi entropy [28, Thm. 1], Tsallis entropy [28, Cor. 1], and Shannon entropy [28, Thm. 2] uncertainty relations are studied.

### 4.3 Future work

Currently, I am mainly interested in

- exploring the application of information geometry technique to hyperbolic secant family, namely the probability density function of the form  $C(s) \operatorname{sech}^t(x)$ ,

since both Bernoulli symbol (1.1) and Euler symbol (1.2) belong to this family.

## References

- [1] J. Ablinger, *Computer Algebra Algorithms for Special Functions in Particle Physics*, PhD Thesis, Research Institute for Symbolic Computation, Johannes Kepler University, 2012.
- [2] S. Akiyama and Y. Tanigawa, Multiple zeta values at non-positive integers, *Ramanujan J.* **5** (2001), 327–351.
- [3] S. Amari and H. Nagaoka, *Methods on Information Geometry*, AMS, 2000.
- [4] S. Amari, *Differential-Geometrical Methods in statistics*, Springer Lectures Notes in Statistics 28, 1985.
- [5] T. Amdeberhan, O. Espinosa, I. Gonzalez, M. Harrison, V. H. Moll, and A. Straub, Ramanujan master theorem, *Ramanujan J.* **29** (2012), 379–422.
- [6] A. Bayad and M. Beck, Relations for Bernoulli-Barnes numbers and Barnes zeta functions, *Int. J. Number Theory* **10** (2014), 1321–1335.
- [7] J. Blümlein, Algebraic relations between harmonic sums and associated quantities. *Comput. Phys. Commun.* **159** (2004), 19–54.
- [8] J. Blümlein and S. Kurth, Harmonic sums and Mellin transforms up to two-loop order, *Phys. Rev. D* **60** (1999), Article 014018.
- [9] A. Byrnes, L. Jiu, V. H. Moll, and C. Vignat, Recursion rules for the hypergeometric zeta functions, *Int. J. Number Theory* **10** (2014), 1761–1782.
- [10] K. Dilcher, Some  $q$ -series identities related to divisor functions. *Discrete Math.* **145** (1995), 83–93.
- [11] A. Dixit, V. H. Moll and C. Vignat, The Zagier modification of Bernoulli numbers and a polynomial extension. Part I. *Ramanujan J.* **33** (2014), 379–422.
- [12] I. Gonzalez, Method of brackets and Feynman diagrams evaluations. *Nucl. Phys. B. Proc. Suppl.* **205** (2010), 141–146.
- [13] I. Gonzalez, K. Kohl, and V. H. Moll, Evaluation of entries in Gradshteyn and Ryzhik employing the method of brackets. *Scientia Series A: Mathematical Sciences* **25** (2014), 65–84.

- [14] I. Gonzalez and V. H. Moll, Definite integrals by the method of brackets. Part 1. *Adv. Appl. Math.* **45** (2010), 50–73.
- [15] I. Gonzalez, V. H. Moll, and A. Straub, The method of brackets. Part 2: examples and applications, in T. Amdeberhan, L. Medina, and V. H. Moll, eds., *Gems in Experimental Mathematics*, Contemporary Mathematics, Vol. 517, AMS, 2010, pp. 157–172.
- [16] I. Gonzalez and I. Schmidt. Optimized negative dimensional integration method (NDIM) and multiloop Feynman diagram calculation, *Nuclear Phys. B* **769** (2007), 124–173.
- [17] I. Gonzalez and I. Schmidt. Modular application of an integration by fractional expansion (IBFE) method to multiloop Feynman diagrams, *Phys. Rev. D* **78** (2008), Article 086003.
- [18] I. Gonzales, L. Jiu and V. H. Moll, Pochhammer symbol with negative indices. A new rule for the method of brackets, *Open Math.* **14** (2016) 681–686.
- [19] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, D. Zwillinger and V. H. Moll eds., Academic Press, 2015.
- [20] G. H. Hardy, *Ramanujan. Twelve Lectures on Subjects Suggested by His Life and Work*, Chelsea Publishing Company, 1978.
- [21] M. E. Hoffman, Quasi-shuffle product, *J. Algebraic Combin.* **11** (2000), 49–68.
- [22] L. Jiu, The method of brackets and the Bernoulli symbols, PhD Thesis, Tulane University, 2016.
- [23] L. Jiu, Matrix representation of harmonic sums, Submitted for Publication.
- [24] L. Jiu, V. H. Moll and C. Vignat, A symbolic approach to multiple zeta values at the negative integers, Submitted for Publication.
- [25] L. Jiu, V. H. Moll, and C. Vignat, A symbolic approach to some identities for Bernoulli-Barnes polynomials, *Int. J. Number Theory* **12** (2016), 649–662.
- [26] L. Jiu, V. H. Moll, and C. Vignat, Identities for generalized Euler polynomials, *Integral Transforms Spec. Funct.* **25** (2014), 777–789.
- [27] K. Kohl. *Algorithmic methods for definite integration*. PhD thesis, Tulane University, 2011.
- [28] Y. Li, B. Li, H. Sun, and L. Jiu, Matrix geometric means and uncertainty relation, Submitted for Publication.
- [29] C. Li, E. Zhang, L. Jiu, and H. Sun, Optimal control on special Euclidean group via natural gradient descent algorithm, *Sci. China Inf. Sci.* **59** (2016) Article: 112203
- [30] D. Li, H. Sun, C. Tao and L. Jiu, Principal Bundles over Statistical Manifolds, Submitted for Publication.
- [31] D. Li, H. Sun, C. Tao and L. Jiu, Riemannian Holonomy Groups of Statistical Manifolds, Submitted for Publication.
- [32] S. Roman, *The Umbral Calculus*, Academic Press Inc., 1984.
- [33] B. Sadaoui, Multiple zeta values at the non-positive integers, *C. R. Acad. Sci. Paris, Ser. I*, **12** (2014), 977–984.
- [34] C. Schneider, Symbolic summation assists combinatorics, *Sem. Lothar. Combin.* **56** (2007), Article B56b.
- [35] J. Zeng, On some  $q$ -identities related to divisor functions, *Adv. Appl. Math.* **34** (2005) 313–315.
- [36] F. Zhang, H. Sun, L. Jiu, and L. Peng, The arc length variational formula on the exponential manifold, *Math. Slovaca* **63** (2013), 1101–1112.
- [37] Z. Zhang, H. Sun, L. Jiu, and L. Peng, A natural gradient algorithm for stochastic distribution systems, *Entropy* **16** (2014), 4338–4352.