Final Exam

● Graded

Student

Eric Duma

Total Points

29 / 45 pts

Question 1

Overall Grades **29** / 45 pts

✔ **– 0 pts** See the details on each problem.

💬 **– 16 pts** Point adjustment

MATH 202–001 FALL 2026 SESSION 1 FINAL EXAM
OCTOBER 19TH, 15:30–18:30

Name (Please Print)	NetID	Signature
ERIC-STEFAN DUMA	ESD26	ES

INSTRUCTION

- This is a 180-minute test.
- The total number of points is 45.
- **No Cellphones.**
- **No Calculators.**
- **Do NOT write on the back of the test; if you do so, your answer(s) will NOT be graded.**
- There are 9 problems.
- Show your work. To receive full credit, your answers must be neatly written, logically organized, and clearly labeled. Writing just the answer will not give you credit for that.
- The backs and extra pages are offered as scratch paper.
- GOOD LUCK.

Duke Community Standard

Duke Kunshan University (DKU) is a community comprised of individuals from diverse cultures and backgrounds. We are dedicated to scholarship, leadership, and service and to the principles of honesty, fairness, respect, and accountability. Members of this community commit to reflecting upon and upholding these principles in all academic and non-academic endeavors and to protecting and promoting a culture of integrity and trust.

To uphold the DKU Community Standard

- I will hold myself to the highest standards for honesty, integrity, fairness, and responsibility in my academic and non-academic endeavors
- I will respect other cultures and embrace all forms of diversity
- I will uphold the standards if they are compromised.

Honor Pledge: I will neither give nor receive aid on this test. SIGNATURE: ES

IF YOU DO NOT SIGN THE HONOR PLEDGE, YOUR FINAL EXAM WILL BE RECORDED AS 0 POINTS.

Problems	1	2	3	4	5	6	7	8	9	Total
Scores	5	2	5	5	2	5	4	0.5	0.5	29

Problem 1. [5] Find the general solution of the linear system:

$$\begin{aligned} 2x - 3y &= 1 \\ -2y - 3z &= 1 \\ x + 2z &= 1 \end{aligned} \quad \sim \begin{pmatrix} 2 & -3 & 0 \\ 0 & -2 & -3 \\ 1 & 0 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

$$\Rightarrow \left(\begin{array}{ccc|c} 2 & -3 & 0 & 1 \\ 0 & -2 & -3 & 1 \\ 1 & 0 & 2 & 1 \end{array}\right) \overset{R_1 \leftrightarrow R_3}{\sim} \left(\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & -2 & -3 & 1 \\ 2 & -3 & 0 & 1 \end{array}\right) \overset{R_3 \leftarrow R_3 - 2R_1}{\sim}$$

$$\overset{R_2 \leftarrow R_2 - R_3}{\left(\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & -2 & -3 & 1 \\ 0 & -3 & -4 & -1 \end{array}\right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & 1 & 1 & 2 \\ 0 & -3 & -4 & -1 \end{array}\right)} \overset{R_3 \leftarrow R_3 + 3R_2}{\sim} \left(\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & -1 & 5 \end{array}\right) \sim$$

$$\overset{\substack{R_1 \leftarrow R_1 + 2R_3 \\ R_2 \leftarrow R_2 + R_3}}{\sim} \left(\begin{array}{ccc|c} 1 & 0 & 0 & 11 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & -1 & 5 \end{array}\right) \overset{R_3 \leftarrow -R_3}{\sim} \left(\begin{array}{ccc|c} 1 & 0 & 0 & 11 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 1 & -5 \end{array}\right)$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 11 \\ 7 \\ -5 \end{pmatrix}$$

2/5

Problem 2. [5'] Let $A = \begin{bmatrix} 2 & 4 & 2 \\ -5 & 0 & 5 \\ 0 & 1 & 1 \\ 1 & 3 & 2 \end{bmatrix}$. Find

(1) a basis of the column space of A,
(2) a basis of the row space of A,
(3) and a basis of the null space of A.

$\begin{pmatrix} 2 & 4 & 2 \\ -5 & 0 & 5 \\ 0 & 1 & 1 \\ 1 & 3 & 2 \end{pmatrix}$ $R_1 \leftarrow \frac{1}{2} R_1$, $R_2 \leftarrow \frac{1}{5} R_2$ $\sim$ $\begin{pmatrix} 1 & 2 & 1 \\ -1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 3 & 2 \end{pmatrix}$ $R_2 \leftarrow R_2 + R_1$, $R_3 \leftarrow R_3 - R_1$ $\sim$

$R_4 \leftarrow R_4 - R_3$ $\begin{pmatrix} 1 & 2 & 1 \\ 0 & 2 & 2 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ $R_1 \leftarrow R_1 - 2R_3$, $R_2 \leftarrow R_2 - 2R_3$ $\sim$ $\begin{pmatrix} 1 & 0 & -1 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$

$R_1 \leftarrow R_1 + R_2$, $R_3 \leftarrow R_3 - R_2$ $\sim$ $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ $R_2 \leftrightarrow R_3$ $\sim$ $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$

a)

~~$\Rightarrow$ basis of $col(A) = \left\{ \begin{pmatrix} 2 \\ -5 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 4 \\ 0 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 5 \\ 1 \\ 2 \end{pmatrix} \right\}$~~

b)

~~basis of row space $= \{(1\,0\,0), (0\,1\,0), (0\,0\,1)\}$~~

~~$\Rightarrow$ basis of null space of A~~

null space $null(A) = \{ x \mid Ax = 0 \}$

$\Rightarrow$ we do $(A|0) <=>$ $\left(\begin{array}{ccc|c} 2 & 4 & 2 & 0 \\ -5 & 0 & 5 & 0 \\ 0 & 1 & 1 & 0 \\ 1 & 3 & 2 & 0 \end{array}\right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right)$ (x y z)

~~we cannot switch rows~~ because it messes up go back t_2

$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ $R_2 \leftarrow R_2 + R_3$ $\sim$ $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ $R_3 \leftarrow R_3 + R_2$ $\sim$ $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{pmatrix}$ $R_2 \leftarrow R_2 - R_3$ $\sim$

$\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{pmatrix}$ $R_3 \leftarrow R_3 + 2R_2$ $\sim$ $\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$ $R_2 \leftarrow -R_2$ $\sim$ $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$

$\Rightarrow$ all columns are pivot columns

$\Rightarrow$ basis of col $A = \left\{ \begin{pmatrix} 2 \\ -5 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 4 \\ 0 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 5 \\ 1 \\ 2 \end{pmatrix} \right\}$

5/5

Problem 3. [5] Let $T : \mathbb{R}^3 \longrightarrow \mathbb{R}^2$ be a linear transformation such that

$$T\left(\begin{bmatrix}0\\1\\-1\end{bmatrix}\right) = \begin{bmatrix}1\\1\end{bmatrix}, \qquad T\left(\begin{bmatrix}1\\0\\-1\end{bmatrix}\right) = \begin{bmatrix}0\\2\end{bmatrix}, \qquad T\left(\begin{bmatrix}0\\0\\-1\end{bmatrix}\right) = \begin{bmatrix}2\\2\end{bmatrix}.$$

Find the matrix transformation of T, i.e., find a matrix A such that

$T(au+bv) = aT(v) + bT(u)$

$$T\left(\begin{bmatrix}x\\y\\z\end{bmatrix}\right) = A\begin{bmatrix}x\\y\\z\end{bmatrix}.$$

we need to find $T\begin{pmatrix}0\\1\\0\end{pmatrix}$, $T\begin{pmatrix}1\\0\\0\end{pmatrix}$, $T\begin{pmatrix}0\\0\\1\end{pmatrix}$

$$T\begin{pmatrix}0\\0\\1\end{pmatrix} = -T\begin{pmatrix}0\\0\\-1\end{pmatrix} = \begin{pmatrix}-2\\-2\end{pmatrix}$$

$$T\begin{pmatrix}0\\1\\0\end{pmatrix} = T\left(\begin{pmatrix}0\\1\\-1\end{pmatrix} + \begin{pmatrix}0\\0\\1\end{pmatrix}\right) = T\begin{pmatrix}0\\1\\0\end{pmatrix} = \begin{pmatrix}1\\1\end{pmatrix} + \begin{pmatrix}-2\\-2\end{pmatrix} = \begin{pmatrix}-1\\-1\end{pmatrix}$$

$$T\begin{pmatrix}1\\0\\0\end{pmatrix} = T\left(\begin{pmatrix}1\\0\\-1\end{pmatrix} + \begin{pmatrix}0\\0\\1\end{pmatrix}\right) = T\begin{pmatrix}1\\0\\0\end{pmatrix} = \begin{pmatrix}0\\2\end{pmatrix} + \begin{pmatrix}-2\\-2\end{pmatrix} = \begin{pmatrix}-2\\0\end{pmatrix}$$

$$\Rightarrow A = \underset{2\times 3}{\begin{pmatrix}-2 & -1 & -2\\0 & -1 & -2\end{pmatrix}}$$

$$T(V) = AV = \underset{2\times 3}{\begin{pmatrix}-2 & -1 & -2\\0 & -1 & -2\end{pmatrix}} \underset{3\times 1}{\begin{pmatrix}x\\y\\z\end{pmatrix}} = \underset{2\times 1}{\begin{pmatrix}-2x-y-2z\\-y-2z\end{pmatrix}}$$

$$T\begin{pmatrix}1\\0\\0\end{pmatrix} = \begin{pmatrix}-2-0-2\cdot 0\\-0-2\cdot 0\end{pmatrix} = \begin{pmatrix}-2\\0\end{pmatrix}$$

$$T\begin{pmatrix}0\\1\\0\end{pmatrix} = \begin{pmatrix}0-1\cdot 0\\-1-0\end{pmatrix} = \begin{pmatrix}-1\\-1\end{pmatrix}$$

$$T\begin{pmatrix}0\\0\\1\end{pmatrix} = \begin{pmatrix}0+0-2\cdot 1\\0-2\cdot 1\end{pmatrix} = \begin{pmatrix}-2\\-2\end{pmatrix}$$

$$\Rightarrow A = \begin{pmatrix}-2 & -1 & -2\\0 & -1 & -2\end{pmatrix}$$

Problem 4. [5'] Find the Jordan canonical form of

$$A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 2 & 0 \\ -2 & -2 & 1 \end{bmatrix}$$

$\det(A-\lambda I) = \begin{vmatrix} 1-\lambda & 2 & 0 \\ 0 & 2-\lambda & 0 \\ -2 & -2 & 1-\lambda \end{vmatrix}$

$\det(A-\lambda I) = (1-\lambda)^2(2-\lambda) + 2(2-\lambda) + 0$

$\det(A-\lambda I) = (2-\lambda)\left[(1-\lambda)^2 + 2\right]$

$\det(A-\lambda I) = (2-\lambda)(\lambda^2 - 2\lambda + 1 + 2)$

$\det(A-\lambda I) = (2-\lambda)(\lambda^2 - 2\lambda + 3)$

$\det(A-\lambda I) = (2-\lambda)(\lambda^2 + \lambda + \cdots)$

$\det(A-\lambda I) = (1-\lambda)^2(2-\lambda)$

$\Rightarrow \lambda_1 = 1 \quad \lambda_2 = 1 \quad \lambda_3 = 2$

$\lambda_{1,2} = 1 \Rightarrow \left(\begin{array}{ccc|c} 1-1 & 2 & 0 & 0 \\ 0 & 2-1 & 0 & 0 \\ -2 & -2 & 1-1 & 0 \end{array}\right) \sim \left(\begin{array}{ccc|c} 0 & 2 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -2 & -2 & 0 & 0 \end{array}\right) \overset{\substack{R_1 \leftarrow \frac{1}{2}R_1 \\ R_3 \leftarrow -\frac{1}{2}R_3}}{\sim}$

$\sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix} \overset{R_3 \leftrightarrow R_1}{\sim} \left(\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{array}\right) \overset{\substack{R_1 \leftarrow R_1 - R_2 \\ R_3 \leftarrow R_3 - R_2}}{\sim} \left(\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right)$

$\Rightarrow z = t \Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = t\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

$\lambda_1 = 1 = \lambda_2 \Rightarrow$ for $\lambda = 1$ algebraic multiplicity $= 2$

but since $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = t\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$ geometric multiplicity $= 1$

$\Rightarrow J = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$ we need to find P so as $A = PJP^{-1} \Leftrightarrow J = P^{-1}AP$

We use $AV_1 = V_1 \Leftrightarrow (A-\lambda I) = 0$ we already did that for $\lambda_1 = 1 \Rightarrow V_1 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

next for $\lambda_2 = 1$ we do

$(A - \lambda_2 I)V_2 = V_1$

$\Leftrightarrow (A-I)V_2 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

2/5

Problem 5. [5'] Diagonalize the quadratic form

$$f(x,y,z) = 3x^2 + 3y^2 + 4z^2 + 2xz - 2yz.$$

$f(x;y;z) = (x\ y\ z)^T A \begin{pmatrix} x \\ y \\ z \end{pmatrix}$, $A = \begin{pmatrix} 3 & 0 & 1 \\ 0 & 3 & -1 \\ 1 & -1 & 4 \end{pmatrix}$

$\det(A - \lambda I) = \begin{vmatrix} 3-\lambda & 0 & 1 \\ 0 & 3-\lambda & -1 \\ 1 & -1 & 4-\lambda \end{vmatrix} =$

$= (3-\lambda)^2(4-\lambda) + 0 + 0 - 3 + \lambda - 3 + \lambda + 0 =$

$= (3-\lambda)^2(4-\lambda) - 2(3-\lambda)$

$= (3-\lambda)[(3-\lambda)(4-\lambda) - 2]$

$= (3-\lambda)[12 - 3\lambda - 4\lambda + \lambda^2 - 2)$

$= (3-\lambda)(\lambda^2 - 7\lambda + 10)$

$= (3-\lambda)[\lambda^2 - 2\lambda - 5\lambda + 10]$

$= (3-\lambda)[\lambda(\lambda-2) - 5(\lambda-2)]$

$= (3-\lambda)(\lambda-2)(\lambda-5)$

$\det(A - \lambda I) = 0 \Rightarrow \lambda_1 = 2,\ \lambda_2 = 3,\ \lambda_3 = 5$

$\Rightarrow D = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{pmatrix}$

let $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = P_{orth} \begin{pmatrix} u \\ v \\ w \end{pmatrix} \Rightarrow (x\ y\ z)^T = (u\ v\ w)^T P_{orth}^T$

$\Rightarrow f(x;y;z) = (u\ v\ w)^T P_{orth}^T A P_{orth} \begin{pmatrix} u \\ v \\ w \end{pmatrix}$

since P_{orth} - orthogonal $\Rightarrow P_{orth}^T = P_{orth}^{-1}$

$\Rightarrow P_{orth}^T A P_{orth} = P_{orth}^{-1} A P_{orth} = D$

$\Rightarrow f(x;y;z) = f(u;v;w) = (u;v;w) D \begin{pmatrix} u \\ v \\ w \end{pmatrix} =$

$= \underset{1\times 3}{(u\ v\ w)} \underset{3\times 3}{\begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{pmatrix}} \underset{1\times 3}{\begin{pmatrix} u \\ v \\ w \end{pmatrix}} = (2u\ 3v\ 5w) \begin{pmatrix} u \\ v \\ w \end{pmatrix} =$

$= 2u^2 + 3v^2 + 5w^2$

Problem 6. [5'] Find a, b, and c such that $\mathbf{w} = a\mathbf{v}_1 + b\mathbf{v}_2 + c\mathbf{v}_3$, where

$$\mathbf{v}_1 = \begin{bmatrix} 0 \\ 2 \\ -2 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{w} = \begin{bmatrix} -3 \\ 8 \\ -8 \end{bmatrix}.$$

$W = aV_1 + bV_2 + cV_3$

$$\Rightarrow \left(\begin{array}{ccc|c} 0 & 2 & -1 & -3 \\ 2 & -1 & 1 & 8 \\ -2 & 2 & 0 & -8 \end{array}\right) \xrightarrow{R_3 \leftarrow R_3 + R_2} \left(\begin{array}{ccc|c} 0 & 2 & -1 & -3 \\ 2 & -1 & 1 & 8 \\ 0 & 1 & 1 & 0 \end{array}\right)$$

$R_1 \leftarrow R_1 - 2R_3$

$R_2 \leftarrow R_2 + R_3$

$$\sim \left(\begin{array}{ccc|c} 0 & 0 & -3 & -3 \\ 2 & 0 & 2 & 8 \\ 0 & 1 & 1 & 0 \end{array}\right) \xrightarrow{R_1 \leftarrow -\frac{1}{3}R_1} \sim \left(\begin{array}{ccc|c} 0 & 0 & 1 & 1 \\ 2 & 0 & 2 & 8 \\ 0 & 1 & 1 & 0 \end{array}\right)$$

$R_2 \leftarrow \frac{1}{2}R_2$

$$\sim \left(\begin{array}{ccc|c} 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 4 \\ 0 & 1 & 1 & 0 \end{array}\right) \xrightarrow[R_3 \leftarrow R_3 - R_1]{R_2 \leftarrow R_2 - R_1} \sim \left(\begin{array}{ccc|c} 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & -1 \end{array}\right)$$

$R_1 \leftrightarrow R_2$

$$\sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & 3 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & -1 \end{array}\right) \xrightarrow{R_2 \leftrightarrow R_3} \left(\begin{array}{ccc|c} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{array}\right)$$

$\Rightarrow \quad a = 3 \qquad b = -1 \qquad c = 1$

$$3\begin{pmatrix} 0 \\ 2 \\ -2 \end{pmatrix} + (-1)\begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix} + \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 14 \\ -14 \end{pmatrix} + \begin{pmatrix} -2 \\ 1 \\ -2 \end{pmatrix} + \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} =$$

$$= \begin{pmatrix} -2-1 \\ \text{WRONG} \end{pmatrix}$$

We go back to $\left(\begin{array}{ccc|c} 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 4 \\ 0 & 1 & 1 & 0 \end{array}\right)$ $\xrightarrow[R_3 \leftarrow R_3 - R_1]{R_2 \leftarrow R_2 - R_1}$

$$\sim \left(\begin{array}{ccc|c} 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & -1 \end{array}\right) \xrightarrow{R_2 \leftarrow R_1} \left(\begin{array}{ccc|c} 1 & 0 & 0 & 3 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & -1 \end{array}\right)$$

$R_2 \leftrightarrow R_3$

$$\sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{array}\right) \Rightarrow \boxed{\begin{array}{l} a = 3 \\ b = -1 \\ c = 1 \end{array}}$$

$$\begin{pmatrix} 3 \cdot 0 - 2 - 1 \\ 3 \cdot 2 + 1 + 1 \\ 3(-2) - 2 + 0 \end{pmatrix} = 3V_1 - 1V_2 + 1V_3 = \begin{pmatrix} -3 \\ 8 \\ -8 \end{pmatrix} = W$$

4/5

Problem 7. [5'] Let V be a subspace of $\mathbb{R}^4$, defined by

$$V = span\left\{\begin{bmatrix}0\\1\\1\\0\end{bmatrix}, \begin{bmatrix}3\\0\\2\\-1\end{bmatrix}\right\}.$$

Find an orthonormal basis for V. we use $e_1 = \begin{pmatrix}1\\0\\0\\0\end{pmatrix}$, $e_2 = \begin{pmatrix}0\\1\\0\\0\end{pmatrix}$, $e_3 = \begin{pmatrix}0\\0\\1\\0\end{pmatrix}$ and $e_4 = \begin{pmatrix}0\\0\\0\\1\end{pmatrix}$

NOT R4

$$\Rightarrow \begin{pmatrix}0&3&1&0&0&0\\1&0&0&1&0&0\\1&2&0&0&1&0\\0&-1&0&0&0&1\end{pmatrix} \overset{R_2 \leftrightarrow R_1}{\sim} \begin{pmatrix}1&0&0&1&0&0\\0&3&1&0&0&0\\1&2&0&0&1&0\\0&-1&0&0&0&1\end{pmatrix} \overset{R_2 \leftrightarrow R_3}{\sim}$$

$$\begin{pmatrix}1&0&0&1&0&0\\1&2&0&0&1&0\\0&3&1&0&0&0\\0&-1&0&0&0&1\end{pmatrix} \overset{R_2 \leftarrow R_2 - R_1}{\sim} \begin{pmatrix}1&0&0&1&0&0\\0&2&0&-1&1&0\\0&3&1&0&0&0\\0&-1&0&0&0&1\end{pmatrix} \overset{\substack{R_2 \leftarrow R_2 + 2R_4\\ R_3 \leftarrow R_3 + 3R_4\\ R_1 \leftarrow R_1 + R_4}}{\sim}$$

$$\begin{pmatrix}1&0&0&1&0&0\\0&0&0&-1&0&2\\0&0&1&0&0&3\\0&-1&0&0&0&1\end{pmatrix} \overset{R_4 \leftrightarrow R_2}{\sim} \begin{pmatrix}1&0&0&1&0&0\\0&-1&0&0&0&1\\0&0&1&0&0&3\\0&0&0&-1&0&2\end{pmatrix} \sim$$

$$\begin{pmatrix}1&0&0&0&0&2\\0&-1&0&0&0&1\\0&0&1&0&0&3\\0&0&0&-1&0&2\end{pmatrix} \overset{\substack{R_2 \leftarrow -R_2\\ R_4 \leftarrow -R_4}}{\sim} \begin{pmatrix}1&0&0&0&0&2\\0&1&0&0&0&-1\\0&0&1&0&0&3\\0&0&0&1&0&-2\end{pmatrix}$$

$$\Rightarrow V = \left\{ \begin{pmatrix}0\\1\\1\\0\end{pmatrix}_{=v_1}, \begin{pmatrix}3\\0\\2\\-1\end{pmatrix}_{=v_2}, \begin{pmatrix}1\\0\\0\\0\end{pmatrix}_{=v_3}, \begin{pmatrix}0\\1\\0\\0\end{pmatrix}_{=v_4} \right\} \text{ bases}$$

bases $\{v_1, v_2, v_3, v_4\} \rightarrow \{u_1, u_2, u_3, u_4\}$ ~~orthogo~~ orthonormal bases

let $u_1 = v_1 = \begin{pmatrix}0\\1\\1\\0\end{pmatrix}$

we use the formula $u_K = v_K - \frac{(u_1; v_K)}{(u_1; u_1)} u_1 - \frac{(u_2; v_K)}{(u_2; u_2)} u_2 - \dots - \frac{(u_{K-1}; v_K)}{(u_{K-1}; u_{K-1})} \cdot u_{K-1}$

$$\Rightarrow u_2 = v_2 - \frac{(u_1; v_2)}{(u_1; u_1)} u_1$$

$(u_1; v_2) = u_1 \cdot v_2 = 0 \cdot 3 + 1 \cdot 0 + 1 \cdot 2 + 0 \cdot (-1) = 2$

0.5/5

Problem 8. [5'] Let A be an $n \times n$ square matrix satisfying $A^T = -A$. Show that if n is odd, then $\det A = 0$.

~~Let $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$~~ let A be 3×3 for this example.

let $A = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}$ $A^T = \begin{pmatrix} a & d & g \\ b & e & h \\ c & f & i \end{pmatrix}$

$A^T = -A$ $\Rightarrow$ $a = -a$, $b = -d$, $c = -g$, $d = -b$, $e = -e$, $f = -h$, $g = -c$, $h = -f$, $i = -i$

$\Rightarrow A = \begin{pmatrix} a & b & c \\ -b & e & f \\ -c & -f & i \end{pmatrix}$ $\Rightarrow \det A = aei + bfc - cbf + cec + fbf + bbi$

$\det A = aei + c^2e + f^2a + b^2i$

but let's look at $a = -a \Rightarrow 2a = 0 \Rightarrow a = 0$

$e = -e \Rightarrow 2e = 0 \Rightarrow e = 0$

$i = -i \Rightarrow 2i = 0 \Rightarrow i = 0$

$\Rightarrow \det A = 0 + 0 + 0 + 0 = 0$

what if we have a bigger matrix?

$A = \begin{pmatrix} a & b & c & v_{14} & v_{15} & \cdots & v_{1n} \\ d & e & f & v_{24} & v_{25} & \cdots & v_{2n} \\ g & h & i & v_{34} & v_{35} & \cdots & v_{3n} \\ v_{41} & v_{42} & v_{43} & v_{44} & v_{55} & \cdots & v_{3n} \\ v_{n1} & v_{n2} & v_{n3} & v_{n4} & v_{n5} & \cdots & v_{nn} \end{pmatrix}$, n odd

$\det(A) =$ Sum of cofactors $(-1)^{i+j} M_{ij} = 0$

0.5/5

Problem 9. [5'] For two vectors $\vec{u}$ and $\vec{v}$, suppose they satisfy that

$$\|\vec{u}+\vec{v}\| = \|\vec{u}-\vec{v}\|.$$

Show that $\vec{u}$ and $\vec{v}$ are perpendicular to each other.

$\|\bar{u}+\bar{v}\| \leq \|\bar{u}\| + \|\bar{v}\|$ Triangle inequality.

$\|\bar{u}+\bar{v}\| = \|\bar{u}\| + \|\bar{v}\|$

$|\bar{u}\cdot\bar{v}| \leq \|u\| \cdot \|v\|$ Cauchy-Schwarz Inequality.

$|\bar{u}\cdot\bar{v}| = \|u\|\cdot\|v\|$

ok let's break the ice

Suppose $\bar{u} = \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix}$ $\bar{v} = \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}$

$\|\bar{u}+\bar{v}\| = \|\bar{u}-\bar{v}\|$

$\sqrt{(u_1+v_1)^2 + (u_2+v_2)^2 + \dots + (u_n+v_n)^2} =$

$= \sqrt{(u_1-v_1)^2 + (u_2-v_2)^2 + \dots + (u_n-v_n)^2}$

$(u_1+v_1)^2 + (u_2+v_2)^2 + \dots + (u_n+v_n)^2 =$

$= (u_1-v_1)^2 + (u_2-v_2)^2 + \dots + (u_n-v_n)^2$

for $n=1 \Rightarrow (u_1+v_1)^2 = (u_1-v_1)^2$

$(u_1+v_1)^2 - (u_1-v_1)^2 = 0$ why: $a^2-b^2 = (a+b)(a-b)$

$(u_1+v_1+u_1-v_1)(u_1+v_1-u_1+v_1) = 0$

$(2u_1)(2v_1) = 0$

$4u_1v_1 = 0$

$u_1v_1 = 0$

we do the same for the others $u_2v_2 = 0$, $u_3v_3 = 0$, $\dots$, $u_nv_n = 0$

$\Rightarrow \bar{u}\cdot\bar{v} = u_1v_1 + u_2v_2 + \dots + u_nv_n$

$\bar{u}\cdot\bar{v} = 0+0+\dots+0$

$\bar{u}\cdot\bar{v} = 0$

$\Rightarrow \bar{u} \perp \bar{v}$