Final Exam

Graded

Student

Xiyun Zhang

Total Points

45 / 45 pts

Question 1

Overall Grades **45** / 45 pts

✔ **– 0 pts** See the details on each problem.

MATH 202–001 FALL 2026 SESSION 1 FINAL EXAM
OCTOBER 19TH, 15:30–18:30

Name (Please Print)	NetID	Signature
Xiyun Zhang	xz475	Xiyun Zhang

INSTRUCTION

- This is a 180-minute test.
- The total number of points is **45**.
- **No Cellphones.**
- **No Calculators.**
- **Do NOT write on the back of the test; if you do so, your answer(s) will NOT be graded.**
- There are 9 problems.
- Show your work. To receive full credit, your answers must be neatly written, logically organized, and clearly labeled. Writing just the answer will not give you credit for that.
- The backs and extra pages are offered as scratch paper.
- GOOD LUCK.

Duke Community Standard

Duke Kunshan University (DKU) is a community comprised of individuals from diverse cultures and backgrounds. We are dedicated to scholarship, leadership, and service and to the principles of honesty, fairness, respect, and accountability. Members of this community commit to reflecting upon and upholding these principles in all academic and non-academic endeavors and to protecting and promoting a culture of integrity and trust.

To uphold the DKU Community Standard

- I will hold myself to the highest standards for honesty, integrity, fairness, and responsibility in my academic and non-academic endeavors
- I will respect other cultures and embrace all forms of diversity
- I will uphold the standards if they are compromised.

Honor Pledge: I will neither give nor receive aid on this test. SIGNATURE: Xiyun Zhang

IF YOU DO NOT SIGN THE HONOR PLEDGE, YOUR FINAL EXAM WILL BE RECORDED AS 0 POINTS.

Problems	1	2	3	4	5	6	7	8	9	Total
Scores	5	5	5	5	5	5	5	5	5	45

Problem 1. [5] Find the general solution of the linear system:

$$\begin{aligned} 2x - 3y &= 1 \\ -2y - 3z &= 1 \\ x + 2z &= 1 \end{aligned}$$

$$\left[\begin{array}{ccc|c} 2 & -3 & 0 & 1 \\ 0 & -2 & -3 & 1 \\ 1 & 0 & 2 & 1 \end{array}\right] \xrightarrow{R_3 \to R_3 - \frac{1}{2}R_1} \left[\begin{array}{ccc|c} 2 & -3 & 0 & 1 \\ 0 & -2 & -3 & 1 \\ 0 & \frac{3}{2} & 2 & \frac{1}{2} \end{array}\right]$$

$$\xrightarrow{R_3 \to R_3 + \frac{3}{4}R_2} \left[\begin{array}{ccc|c} 2 & -3 & 0 & 1 \\ 0 & -2 & -3 & 1 \\ 0 & 0 & -\frac{1}{4} & \frac{5}{4} \end{array}\right] \xrightarrow{R_3 \to -4R_3} \left[\begin{array}{ccc|c} 2 & -3 & 0 & 1 \\ 0 & -2 & -3 & 1 \\ 0 & 0 & 1 & -5 \end{array}\right]$$

$$\xrightarrow{R_2 \to R_2 + 3R_3} \left[\begin{array}{ccc|c} 2 & -3 & 0 & 1 \\ 0 & -2 & 0 & -14 \\ 0 & 0 & 1 & -5 \end{array}\right] \xrightarrow{R_2 \to \frac{1}{2}R_2} \left[\begin{array}{ccc|c} 2 & -3 & 0 & 1 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 1 & -5 \end{array}\right]$$

$$\xrightarrow{R_1 \to R_1 + 3R_2} \left[\begin{array}{ccc|c} 2 & 0 & 0 & 22 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 1 & -5 \end{array}\right] \xrightarrow{R_1 \to \frac{1}{2}R_1} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 11 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 1 & -5 \end{array}\right]$$

So the general solution of this linear system is:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 11 \\ 7 \\ -5 \end{bmatrix}$$

Problem 2. [5'] Let $A = \begin{bmatrix} 2 & 4 & 2 \\ -5 & 0 & 5 \\ 0 & 1 & 1 \\ 1 & 3 & 2 \end{bmatrix}$. Find

(1) a basis of the column space of A,
(2) a basis of the row space of A,
(3) and a basis of the null space of A.

由. $A \begin{bmatrix} 2 & 4 & 2 \\ -5 & 0 & 5 \\ 0 & 1 & 1 \\ 1 & 3 & 2 \end{bmatrix} \xrightarrow{R_1 \to -\frac{1}{2}R_1,\ R_2 \to -\frac{1}{5}R_2} \begin{bmatrix} 1 & 2 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 1 \\ 1 & 3 & 2 \end{bmatrix} \xrightarrow{R_4 \to R_4 - R_1,\ R_4 \to R_4 - R_3} \begin{bmatrix} 1 & 2 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_2 \to R_2 - R_1} \begin{bmatrix} 1 & 2 & 1 \\ 0 & -2 & -2 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_3 \to 2R_3 + \frac{1}{2}R_2} \begin{bmatrix} 1 & 2 & 1 \\ 0 & -2 & -2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$.

So 1 and -2 are the pivot entries of A.

(1). basis of column space of A: $\begin{bmatrix} 2 \\ -5 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 4 \\ 0 \\ 1 \\ 3 \end{bmatrix}$.

(2) basis of row space of A: $[2\ 4\ 2], [-5\ 0\ 5]$

(3). $\begin{bmatrix} 1 & 2 & 1 & 0 \\ 0 & -2 & -2 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_2 \to -\frac{1}{2}R_2} \begin{bmatrix} 1 & 2 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 \to R_1 - 2R_2} \begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

~~null~~ ~~solution~~. so the null space is $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$.

Problem 3. [5] Let $T : \mathbb{R}^3 \longrightarrow \mathbb{R}^2$ be a linear transformation such that

$$T\left(\begin{bmatrix}0\\1\\-1\end{bmatrix}\right) = \begin{bmatrix}1\\1\end{bmatrix}, \qquad T\left(\begin{bmatrix}1\\0\\-1\end{bmatrix}\right) = \begin{bmatrix}0\\2\end{bmatrix}, \qquad T\left(\begin{bmatrix}0\\0\\-1\end{bmatrix}\right) = \begin{bmatrix}2\\2\end{bmatrix}.$$

Find the matrix transformation of T, i.e., find a matrix A such that

$$T\left(\begin{bmatrix}x\\y\\z\end{bmatrix}\right) = A\begin{bmatrix}x\\y\\z\end{bmatrix}.$$

$$\left[\begin{array}{ccc|ccc} 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ -1 & -1 & -1 & 0 & 0 & 1 \end{array}\right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ -1 & -1 & -1 & 0 & 0 & 1 \end{array}\right]$$

$$\xrightarrow{R_3 \to R_3 + R_1,\ R_3 \to R_3 + R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 1 & 1 \end{array}\right] \xrightarrow{R_3 \to -R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & -1 & -1 \end{array}\right]$$

Set ~~A~~ $M = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ -1 & -1 & -1 \end{bmatrix}$ so $M^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ -1 & -1 & -1 \end{bmatrix}$.

Set $S = \begin{bmatrix} 1 & 0 & 2 \\ 1 & 2 & 2 \end{bmatrix}$

$AM = S$.

$A = SM^{-1}$

$$= \begin{bmatrix} 1 & 0 & 2 \\ 1 & 2 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ -1 & -1 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} -2 & -1 & -2 \\ 0 & -1 & -2 \end{bmatrix}$$

Problem 4. [5'] Find the Jordan canonical form of

$$\begin{bmatrix} 1 & 2 & 0 \\ 0 & 2 & 0 \\ -2 & -2 & 1 \end{bmatrix}$$

$$\det \begin{bmatrix} 1-\lambda & 2 & 0 \\ 0 & 2-\lambda & 0 \\ -2 & -2 & 1-\lambda \end{bmatrix} = (2-\lambda)\cdot \det \begin{bmatrix} 1-\lambda & 0 \\ -2 & 1-\lambda \end{bmatrix}$$

$$= (2-\lambda)\left[(1-\lambda)^2 + 0\right] = (2-\lambda)(1-\lambda)^2$$

~~$= (2-\lambda)\cdot(\lambda^2 - 2\lambda + 3)$~~

~~$= (2-\lambda)(\lambda-1)(\lambda$~~

when $\lambda = 2$:

$$\left[\begin{array}{ccc|c} -1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -2 & -2 & -1 & 0 \end{array}\right] \xrightarrow{R_2 \leftrightarrow R_3} \left[\begin{array}{ccc|c} -1 & 2 & 0 & 0 \\ -2 & -2 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right]$$

$R_2 \to R_2 - 2R_1$

$$\left[\begin{array}{ccc|c} -1 & 2 & 0 & 0 \\ 0 & -6 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \xrightarrow{R_2 \to -\frac{1}{6}R_2,\ R_1 \to -R_1} \left[\begin{array}{ccc|c} 1 & -2 & 0 & 0 \\ 0 & 1 & \frac{1}{6} & 0 \\ 0 & 0 & 0 & 0 \end{array}\right]$$

$R_1 \to R_1 + 2R_2$

$$\left[\begin{array}{ccc|c} 1 & 0 & \frac{1}{3} & 0 \\ 0 & 1 & \frac{1}{6} & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \qquad v = \begin{bmatrix} -2 \\ -1 \\ 6 \end{bmatrix}$$

when $\lambda = 1$:

$$\left[\begin{array}{ccc|c} 0 & 2 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -2 & -2 & 0 & 0 \end{array}\right] \xrightarrow{R_3 \leftrightarrow R_1} \left[\begin{array}{ccc|c} -2 & -2 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 2 & 0 & 0 \end{array}\right] \xrightarrow{R_3 \to R_3 - 2R_2} \left[\begin{array}{ccc|c} -2 & -2 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right]$$

$R_1 \to R_1 - \frac{1}{2}R_1$

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \xrightarrow{R_1 \to R_1 - R_2} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \qquad v = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$J = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

Problem 5. [5] Diagonalize the quadratic form

$$f(x,y,z) = 3x^2 + 3y^2 + 4z^2 + 2xz - 2yz.$$

$$A = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 3 & -1 \\ 1 & -1 & 4 \end{bmatrix}$$

$$\det \begin{bmatrix} 3-\lambda & 0 & 1 \\ 0 & 3-\lambda & -1 \\ 1 & -1 & 4-\lambda \end{bmatrix} = (3-\lambda)\det \begin{bmatrix} 4-\lambda & -1 \\ -1 & 4-\lambda \end{bmatrix} + \det \begin{bmatrix} 0 & 4-\lambda \\ 1 & -1 \end{bmatrix}$$

$R_1 \to R_1 + R_2$

$$\sim \begin{bmatrix} 3-\lambda & 3-\lambda & 0 \\ 0 & 3-\lambda & -1 \\ 1 & -1 & 4-\lambda \end{bmatrix} = (3-\lambda)\left[(4-\lambda)-1\right] + (\lambda-4)$$

$$= (\lambda-4)\left[(3-\lambda)(\lambda^2-8\lambda+15)+1\right] = (\lambda-4)$$

$$\det \begin{bmatrix} 3-\lambda & 0 & 1 \\ 0 & 3-\lambda & -1 \\ 1 & -1 & 4-\lambda \end{bmatrix} = -\lambda^3 + 8\lambda^2 + 3\lambda^2 = 15\lambda - 24\lambda + \lambda + 45 - 4$$

$$= -\lambda^3 + 11\lambda^2 - 38\lambda + 41$$

$$= (3-\lambda)\det \begin{bmatrix} 3-\lambda & -1 \\ -1 & 4-\lambda \end{bmatrix} - (3-\lambda)\det \begin{bmatrix} 0 & -1 \\ 1 & 4-\lambda \end{bmatrix}$$

$$= (3-\lambda)\left[(3-\lambda)(4-\lambda)-1\right] - (3-\lambda)(0+1)$$

$$= (3-\lambda)(\lambda^2 - 7\lambda + 12 - 2) = (3-\lambda)(\lambda-2)(\lambda-5)$$

when $\lambda = 2$:

$$\left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 1 & -1 & 2 & 0 \end{array}\right] \sim \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \quad \text{so } v = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix} \quad v_{(\text{unit})} = \begin{bmatrix} -\frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{bmatrix}$$

when $\lambda = 3$

$$\left[\begin{array}{ccc|c} 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 \\ 1 & -1 & 1 & 0 \end{array}\right] \sim \left[\begin{array}{ccc|c} 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \sim \left[\begin{array}{ccc|c} 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \quad \text{so } v = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad v_{(\text{unit})} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \end{bmatrix}$$

when $\lambda = 5$

$$\left[\begin{array}{ccc|c} -2 & 0 & 1 & 0 \\ 0 & -2 & -1 & 0 \\ 1 & -1 & -1 & 0 \end{array}\right] \sim \left[\begin{array}{ccc|c} 1 & 0 & -\frac{1}{2} & 0 \\ 0 & 1 & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \quad v = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \quad v_{(\text{unit})} = \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix}$$

$$\text{so } D = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{bmatrix} \quad P = \begin{bmatrix} -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & \frac{2}{\sqrt{6}} \end{bmatrix} \quad \text{since eigenvectors are orthonormal: } P^{-1} = P^T = \begin{bmatrix} -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} \end{bmatrix}$$

$$\text{so } A = PDP^{-1} = \begin{bmatrix} -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & \frac{2}{\sqrt{6}} \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} \end{bmatrix}$$

5/5

Problem 6. [5] Find a, b, and c such that $\mathbf{w} = a\mathbf{v}_1 + b\mathbf{v}_2 + c\mathbf{v}_3$, where

$$\mathbf{v}_1 = \begin{bmatrix} 0 \\ 2 \\ -2 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{w} = \begin{bmatrix} -3 \\ 8 \\ -8 \end{bmatrix}.$$

since ~~w=~~



since $\vec{w} = a\vec{v_1} + b\vec{v_2} + c\vec{v_3}$

$$\vec{w} = \begin{bmatrix} \vec{v_1} & \vec{v_2} & \vec{v_3} \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

$$\left[\begin{array}{ccc|c} 0 & 2 & -1 & -3 \\ 2 & -1 & 1 & 8 \\ -2 & 2 & 0 & -8 \end{array}\right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|c} 2 & -1 & 1 & 8 \\ 0 & 2 & -1 & -3 \\ -2 & 2 & 0 & -8 \end{array}\right] \xrightarrow{R_3 \to R_3 + R_1} \left[\begin{array}{ccc|c} 2 & -1 & 1 & 8 \\ 0 & 2 & -1 & -3 \\ 0 & 1 & 1 & 0 \end{array}\right] \xrightarrow{R_2 \leftrightarrow R_3} \left[\begin{array}{ccc|c} 2 & -1 & 1 & 8 \\ 0 & 1 & 1 & 0 \\ 0 & 2 & -1 & -3 \end{array}\right]$$

$$\xrightarrow{R_3 \to R_3 - 2R_2} \left[\begin{array}{ccc|c} 2 & -1 & 1 & 8 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & -3 & -3 \end{array}\right] \xrightarrow{R_3 \to -\frac{1}{3}R_3} \left[\begin{array}{ccc|c} 2 & -1 & 1 & 8 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{array}\right] \xrightarrow{R_2 \to R_2 - R_3} \left[\begin{array}{ccc|c} 2 & -1 & 1 & 8 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{array}\right]$$

$$\xrightarrow{R_1 \to R_1 - R_3,\ R_1 \to R_1 + R_2} \left[\begin{array}{ccc|c} 2 & 0 & 0 & 6 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{array}\right] \xrightarrow{R_1 \to \frac{1}{2}R_1} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{array}\right]$$

So $\begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}$

Problem 7. [5] Let V be a subspace of $\mathbb{R}^4$, defined by

$$V = span\left\{\begin{bmatrix}0\\1\\1\\0\end{bmatrix}, \begin{bmatrix}3\\0\\2\\-1\end{bmatrix}\right\}.$$

Find an orthonormal basis for V.

$\vec{u_1} = \vec{v_1} = \begin{bmatrix}0\\1\\1\\0\end{bmatrix}$

$\vec{u_2} = \vec{v_2} - \frac{\vec{u_1}\cdot\vec{v_2}}{\|\vec{u_1}\|^2}\cdot\vec{u_1}$

$= \begin{bmatrix}3\\0\\2\\-1\end{bmatrix} - \frac{2}{2}\begin{bmatrix}0\\1\\1\\0\end{bmatrix}$

$= \begin{bmatrix}3\\-1\\1\\-1\end{bmatrix}$

orthogonal basis: $\left\{\begin{bmatrix}0\\1\\1\\0\end{bmatrix}, \begin{bmatrix}3\\-1\\1\\-1\end{bmatrix}\right\}$

~~orthogonal~~ orthonormal basis: $\left\{\begin{bmatrix}0\\\frac{1}{\sqrt{2}}\\\frac{1}{\sqrt{2}}\\0\end{bmatrix}, \begin{bmatrix}\frac{3}{\sqrt{13}}\\-\frac{1}{\sqrt{13}}\\\frac{1}{\sqrt{13}}\\-\frac{1}{\sqrt{13}}\end{bmatrix}\right\}$


Problem 8. [5] Let A be an $n \times n$ square matrix satisfying $A^T = -A$. Show that if n is odd, then $\det A = 0$.

since $A^T = -A$.

$$A = \begin{bmatrix} 0 & b & c \\ -b & 0 & \\ & & 0 \end{bmatrix}$$

Set $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$, $A^T = \begin{bmatrix} a & d & g \\ b & e & h \\ c & f & i \end{bmatrix} = \begin{bmatrix} -a & -b & -c \\ -d & -e & -f \\ -g & -h & -i \end{bmatrix}$

~~det A =~~ So $A = \begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix}$

So $A =$ 

since $A^T = -A$.

$\det(A^T) = \det(-A)$

since $\det(A^T) = \det(A)$.

$\det(A) = \det(-A)$.

Since: $\det(kA) = k^n \det A$.

So when n is even:

$\det(kA) = \det A$

So $\det(A) = \det(A)$

when n is odd:

$\det(kA) = -\det A$.

$2\det(A) = 0$

$\det(A) = 0$.

5/5

Problem 6. [5'] For two vectors $\vec{u}$ and $\vec{v}$, suppose they satisfy that

$$\|\vec{u}+\vec{v}\| = \|\vec{u}-\vec{v}\|.$$

Show that $\vec{u}$ and $\vec{v}$ are perpendicular to each other.

$\|\vec{u}+\vec{v}\| = \|\vec{u}-\vec{v}\|$

$\|\vec{u}+\vec{v}\|^2 = \|\vec{u}-\vec{v}\|^2$

$(\vec{u}+\vec{v})\cdot(\vec{u}+\vec{v}) = (\vec{u}-\vec{v})(\vec{u}-\vec{v})$

$\|\vec{u}\|^2 + 2\vec{u}\cdot\vec{v} + \|\vec{v}\|^2 = \|\vec{u}\|^2 + \|\vec{v}\|^2 - 2\vec{u}\cdot\vec{v}$

$4\vec{u}\cdot\vec{v} = 0$

$\vec{u}\cdot\vec{v} = 0$

so $\vec{u}$ and $\vec{v}$ are perpendicular to each other