

YONSEI UNIVERSITY
Department of Computer Science

CSI3109 Automata and Formal Languages, SPRING 2017

Homework No.2

DUE: 2017.04.17 13:00:00pm

Student ID: 2014117007

Student Name: Chung, Ji-Wan

1. Given a DFA $A = (Q, \Sigma, \delta, s, F)$, where

$$Q = \{1, 2, 3, 4, 5, 6\}, \Sigma = \{a, b\}, s = 1, F = \{2, 5, 6\}$$

and δ is defined as follows:


	a	b
1	2	3
2	2	4
3	4	5
4	2	6
5	5	4
6	6	6

Run the TF (Table Filling) algorithm and draw a minimal DFA for $L(A)$.

2	X	-	-	-	-
3	X	X	-	-	-
4	X	X	X	-	-
5	X		X	X	-
6	X	X	X	X	X
/	1	2	3	4	5

Equivalence Class = (1), (3), (4), (6), (2, 5)

→ drawing on the next page



2. Prove that the following languages are not regular. You may use the pumping lemma or the closure properties of regular languages under union, intersection and complement.
- a) $L = \{0^n 1^m 0^n \mid m, n \geq 0\}$.

Answer:

- i. Assume that L is regular. Then, there is a pumping constant $n \geq 1$ strings with length greater than or equal to n can be 'pumped' according to pumping lemma.
- ii. For an arbitrary natural number $l \geq n$, we take a string $w = 0^l 1^l 0^l \in L$.
- iii. Following pumping lemma, since $|w| > l \geq n$ w can be divided into $w = xyz$ of which $|xy| \leq n \leq l$ and $|y| > 0$ holds.
- iv. Since $|xy| \leq l$, we can safely deduct the substrings x, y are composed of the symbol 0 only. Likewise we define $x = 0^i, y = 0^{j-i}$ for some $0 \leq i < j \leq n \leq l$.
- v. Then we consider another string $w' = xy^2z = 0^i 0^{2(j-i)} 0^{l-j} 1^l 0^l = 0^{l+(j-i)} 1^l 0^l$ which can be pumped from w . Since $|y| > 0, j - i > 0$. Then $l + (j - i) > l$, rendering $w' \notin L$. For that contradicts our assumption that the pumping lemma holds for L , L is not regular.

- b) $L = \{w \mid w \in \{0,1\}^* \text{ is not a palindrome}\}$.
(A palindrome is a string that reads the same forward and backward. e.g.: racecar)

Answer:

- i. Assume L is regular. Then, $P = \overline{L} = \{w \mid w \in \{0,1\}^* \text{ is a palindrome}\}$ is also regular since a regular language is closed under complement.
- ii. We take contrapositive of i). Then, if we can prove the nonregularity of P we prove the nonregularity of L automatically.
- iii. Now, define a Language $L' = L(0^*1^*0^*)$. We can see L' is regular by its definition relying on a regular expression. Then we recognize $L' \cap P$ equals the language of $a)$, which is already proven unregular. If P is regular, the language of $a)$ should be regular since a regular language is closed under intersection, which is not the case. So P is nonregular.
- iv. Combining ii) and iii), we can deduct L is also a nonregular language.

3. Show that the regular languages are closed under the following operations:

a)

$$\text{DROPOUT}(L) = \{xz \mid xyz \in L, \text{ where } x, z \in \Sigma^*, y \in \Sigma\}.$$

Namely, $\text{DROPOUT}(L)$ is the language containing all strings that can be obtained by removing one symbol from a string in L .

For example, if $L = \{012\}$, then $\text{DROPOUT}(L) = \{12, 02, 01\}$.

Answer:

- i. If L is regular, there is at least one corresponding regular expression E recognizing L .
- ii. Now we construct a regular expression E' recognizing the language of $\text{DROPOUT}(L)$. Beforehand, we define a function f which accepts a RegEx E'' and returns a set formed by processing DROPOUT on the language accepted by that expression.
- iii. For $E'' = \emptyset$, $f(\emptyset) = \emptyset$ by definition.
- iv. For $E'' = \{\epsilon\}$, $f(\{\epsilon\}) = \emptyset$ by definition.
- v. For $E'' = \{\sigma\}$, $f(\{\sigma\}) = \{\epsilon\}$ by definition.
- vi. For union operation, $f(E_1 + E_2) = f(L(E_1) \cup L(E_2))$. Note that for a string w , $w \in L(E_1) \cup L(E_2)$ implies either $w \in L(E_1)$ or $w \in L(E_2)$ holds. For the former, $f(\{w\}) \subset f(L(E_1))$ and $f(L(E_1) \cup L(E_2))$ respectively. For the latter, $f(\{w\}) \subset f(L(E_2))$ and $f(L(E_1) \cup L(E_2))$. So every w is in $f(L(E_1) \cup L(E_2))$ if and only if it is in $f(L(E_1)) \cup f(L(E_2))$. Hence $f(E_1 + E_2) = f(E_1) + f(E_2)$.
- vii. For catenation, a string w' in $f(E_1 \cdot E_2)$ has a symbol deleted from w in $E_1 \cdot E_2$. Since that deletion happened either in E_1 or E_2 , w is either in $f(E_1) \cdot E_2$ or $E_2 \cdot f(E_2)$. Hence $f(E_1 \cdot E_2) = f(E_1) \cdot E_1 + E_1 \cdot f(E_2)$.
- viii. For Kleene Star, we can divide the case of R^* into ϵ and R^+ . The epsilon case was dealt with in *iv*). For the latter, consider R^* can be divided into an arbitrary sequence of R^* . With R^+ , it can be followed that $R^+ = R^* \cdot R \cdot R^*$ for R^+ contains at least one R . Now, $f(R^+) = f(R^* \cdot R \cdot R^*)$ for any arbitrary choice of the single R from the sequence. Since DROPOUT has to remove a symbol from an occurrence of R , let that be our R . Then, $f(R^+) = f(R^* \cdot R \cdot R^*) = R^* \cdot f(R) \cdot R^*$.
- ix. Now that for any RegEx E we can construct $E' = f(E)$, we can conclude a regular language is closed under DROPOUT .

b)

$$\text{INIT}(L) = \{w \mid w \text{ for some } x, wx \in L\}.$$

For example, if $L = \{01, 110\}$, then $\text{INIT}(L) = \{0, 01, 1, 11, 110\}$.

(*HINT*: Start with a DFA A for L and describe how to construct an FA for $\text{INIT}(L)$ using A . We assume that A has no sink states.)

Answer:

- i. Let a DFA accepting L be $A = (Q, \Sigma, \delta, s, F)$. Then, we construct a DFA A' accepting $\text{INIT}(L)$.
- ii. Let A' be $A' = (Q, \Sigma, \delta, s, F')$. The single difference between A and A' is the final states F and F' . We define $F' = Q/\{s\}$. Namely, we let every state of A' except for the start state be final state.
- iii. Pick an arbitrary string $w = \sigma^1 \sigma^2 \dots \sigma^n \in L$. Note $|w| = n$. Now we prove $\sigma^1 \in L(A'), \sigma^1 \sigma^2 \in L(A'), \dots w \in L(A')$.
- iv. For w to be accepted in A , there has to be a sequence $s \xrightarrow{\sigma^1_A} q_1 \xrightarrow{\sigma^2_A} \dots f$. Since A' has the same set of transition functions as A , it follows $s \xrightarrow{\sigma^1_{A'}} q_1 \xrightarrow{\sigma^2_{A'}} \dots f$.
- v. Consider $w_1 = \sigma_1$. With A , processing w_1 takes one transition which takes the current counter to the state q_1 . The situation is almost alike with A' except that in A' , $q_1 \in F$. So A' accepts w_1 .
- vi. Assume A' accepts $w_n = \sigma_1 \sigma_2 \dots \sigma_n$. Then we consider whether A' accepts $w_{n+1} = \sigma_1 \sigma_2 \dots \sigma_n \sigma_{n+1}$.
- vii. Since $q_n \xrightarrow{\sigma^{n+1}_A} q_{n+1}$, $q_n \xrightarrow{\sigma^{n+1}_{A'}} q_{n+1}$. However, q_{n+1} is a final state of A' . So A' accepts w_{n+1} as well.
- viii. With mathematical induction from *vi*) and *vii*), we can deduct A' accepts the language of $\text{INIT}(L)$.
- ix. Since there is a DFA accepting $\text{INIT}(L)$ for every regular language L , we conclude regular languages are closed under the operation of INIT .

4. Given two NFAs $A_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$ and $A_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$, suggest an NFA construction for $L(A_1) \cap L(A_2)$ and justify the construction (in other words, prove the correctness of your construction.)

Answer: We define an NFA $A = (Q, \Sigma, \delta, s, F)$ where $Q = Q_1 \times Q_2$, $s = (s_1, s_2)$, $F = F_1 \times F_2$ and the transition function δ is defined by $\delta((q_{11}, q_{21}), \sigma) = Q'$ where $Q' = \{(q_{12}, q_{22}) \mid \delta_1(q_{11}, \sigma) = q_{12} \wedge \delta_2(q_{21}, \sigma) = q_{22}\}$. We suppose A accepts the language of $L(A_1) \cap L(A_2)$.

Then we prove that it indeed does.

For a string w of which $|w| = n$ holds, $w \in L(A_1) \cap L(A_2)$ means $w \in L(A_1)$ and $w \in L(A_2)$, which then implies both A_1 and A_2 accepts w . That says there are two sequences $s_1 \mapsto_{A_1}^{\sigma_1} q_{11} \mapsto_{A_1}^{\sigma_2} q_{12} \cdots \mapsto_{A_1}^{\sigma_n} f_1, f_1 \in F_1$ and $s_2 \mapsto_{A_2}^{\sigma_1} q_{21} \mapsto_{A_2}^{\sigma_2} q_{22} \cdots \mapsto_{A_2}^{\sigma_n} f_2, f_2 \in F_2$ respectively.

Now we do the mathematical induction on the length of a string $w \in L(A_1) \cap L(A_2)$. If $|w| = 0$, $w = \epsilon$. We call this case vacuously true since if $s_1 \in F_1$ and $s_2 \in F_2$, $(s_1, s_2) \in F_1 \times F_2$.

For $|w| = 1$, we can denote $w = \sigma$ for a $\sigma \in \Sigma$. Since $s_1 \mapsto_{A_1}^{sigma} Q_1$ and $s_2 \mapsto_{A_2}^{\sigma} Q_2$ results in $(s_1, s_2) \mapsto_A^{\sigma} Q, Q = Q_1 \times Q_2$ by definition, $w \in L(A)$.

We assume A accepts all w of which length is $|w| \leq n$. Then we prove A accepts w' of which length is $|w'| = n + 1$. Since w' is accepted by both A_1 and A_2 , there are two respective sequences from s_1 and s_2 to q_{1n+1} and q_{2n+1} . That implies $s_1 \mapsto_{A_1}^* q_{1n} \mapsto_{A_1}^{\sigma_{n+1}} q_{1n+1}$ and $s_2 \mapsto_{A_2}^* q_{2n} \mapsto_{A_2}^{\sigma_{n+1}} q_{2n+1}$. Here, from our assumption $(s_1, s_2) \mapsto_A^* (q_{1n}, q_{2n})$. By definition of A we can deduct $(q_{1n}, q_{2n}) \mapsto_A^{\sigma_{n+1}} (q_{1n+1}, q_{2n+1})$. Hence, there is a sequence from (s_1, s_2) to (q_{1n+1}, q_{2n+1}) .

From the induction, we can claim that A accepts any length of $w \in L(A_1) \cap L(A_2)$.

5. Consider the following two languages:

- $L_1 = \{w \mid w \text{ has the same number of } a\text{'s and } b\text{'s}\}.$
 - $L_2 = \{w \mid w \text{ has the same number of the substrings } ab \text{ and } ba.\}.$
- a) Is L_1 regular? Justify your answer—If L_1 is regular, show a regular expression or an FA. If not, prove it.

Answer:


If L_1 is regular, $L_1 \cap L_a (L_a = L(a^*b^*))$ is regular since a regular language is closed under intersection. However $L_1 \cap L_a = \{a^i b^i \mid i \geq 0\}$, which is a nonregular language. Hence our assumption is wrong, so L_1 is not regular.

b) Is L_2 regular? Justify your answer.

Answer:

We construct a DFA $A = (Q, \Sigma, \delta, s, F)$ in which $Q = \{q_0, q_1, q_2, q_3, q_4\}$, $\Sigma = \{a, b\}$, $s = q_0$, $F = \{q_0, q_1, q_2\}$.

The transition functions are defined as following: $\delta(q_0, a) = q_1$, $\delta(q_1, a) = q_1$, $\delta(q_1, b) = q_4$, $\delta(q_4, b) = q_4$, $\delta(q_4, a) = q_1$, $\delta(q_0, b) = q_2$, $\delta(q_2, b) = q_2$, $\delta(q_2, a) = q_3$, $\delta(q_3, a) = q_3$, $\delta(q_3, b) = q_2$.



First we should notice that the difference between $|w|_{ab}$ and $|w|_{ba}$ is at most 1. Because $s \in F$, $\epsilon \in L(A)$. Additionally, $s \mapsto_A^{a^*} q_1$ and $s \mapsto_A^{b^*} q_2$ makes $L(A)$ accept a^* and b^* since $q_1, q_2 \in F$.

Now we assume for a string w for which $w \in L_2 \wedge 0 < |w| \leq n$, A accepts w . Then, we prove the same holds for w' that $w' \in L_2 \wedge |w'| = n + 1$.

For $w' = v\sigma$, A can distinguish the string v , ($|v| \geq 1$).

- If A accepts v , A would be at state either q_1 or q_2 .
 - If A is at state q_1 after reading v , the last symbol of v is a .
 - * If σ is a , w' has the same occurrence of ab, ba with v , so $w' \in L_2$. Since A terminates in q_1 after reading w' , $w' \in L(A)$ too.
 - * If σ is b , w' has one more occurrence of ab than v , so ab occurs one more time than ba in w' and thus $w' \notin L_2$. Since A terminates in q_4 after reading w' , $w' \notin L(A)$ too.
 - If A is at state q_2 after reading v , the last symbol of v is b .
 - * If σ is b , w' has the same occurrence of ab, ba with v , so $w' \in L_2$. Since A terminates in q_2 after reading w' , $w' \in L(A)$ too.

- * If σ is a , w' has one more occurrence of ba than v , so ba occurs one more time than ab in w' and thus $w' \notin L_2$. Since A terminates in q_3 after reading w' , $w' \notin L(A)$ too.
- If A does not accept v , A would be at state either q_4 or q_3 .
 - If A is at state q_4 after reading v , the last symbol of v is b .
 - * If σ is a , w' has one more occurrence of ba than v , so ba occurs the same times as ab in w' and thus $w' \in L_2$. Since A terminates in q_1 after reading w' , $w' \in L(A)$ too.
 - * If σ is b , w' has the same occurrence of ab, ba with v , so $w' \notin L_2$. Since A terminates in q_4 after reading w' , $w' \notin L(A)$ too.
 - If A is at state q_3 after reading v , the last symbol of v is a .
 - * If σ is b , w' has one more occurrence of ab than v , so ab occurs the same times as ba in w' and thus $w' \in L_2$. Since A terminates in q_2 after reading w' , $w' \in L(A)$ too.
 - * If σ is a , w' has the same occurrence of ab, ba with v , so $w' \notin L_2$. Since A terminates in q_3 after reading w' , $w' \notin L(A)$ too.

There we proved $L(A) = L_2$ for any length of string $w \in L_2$.