

# 1 Introduction

The global focus on sustainability in agronomic industries has changed the way in which these enterprises do business. When strategies for a sustainable industry are assessed, there is a trade-off between balancing the amount of resources invested and the resultant quality verses quantity of crops produced. This dilemma exists across agriculture, sharing fundamental considerations such as water and nitrogen input [11, 14, 37]. Within viticulture (the cultivation of grapes for wine production) quality is driven by its integration within the wine industry; with a wine’s potential quality being initially defined through the chemical makeup of the grapes used in its production. The consideration of sustainability is further complicated by environmental and socio-demographic pressures. In the Australian context, these include biosecurity, climate and international market demands.

In this analysis we observe relationships between quantity and quality through the use of linear models. Quality can be defined in a variety of ways for grapes, such as . For the purpose of this study quality was defined by the financial value of grapes per tonne; this was due to varying differences in defining quality between grape varieties.

An extensive amount of research into a variety of factor’s effect on grape quality and yield exists. Due to the lack of long-term and in-depth data sources, individual effects are often studied in isolation [2]. The lack of consolidated datasets also restricts the ability to gain statistical insight at a large scale across multiple regions [15, 16]. The dataset used for this analysis includes data collected for the past 10 years from a multitude of vineyards located over a diverse range of Australian winegrowing regions.

# 2 Methods

We created four linear models to explore the relationships between resource use and vineyard outputs (see table 1). The models looked at the value and quantity of grapes produced by vineyards, as a total and as a scale of area harvested. We compared values to observe how economies of scale affect the use of resources.

Table 1: Summary of models; their predictors, covariates and variable interactions.

|         | Response  | Predictors                    | Covariates               |
|---------|---|-------------------------------|--------------------------|
| Model 1 | Yield   | Water Used, Scope 1 Emissions | Area Harvested, Year, GI |
| Model 2 | $\frac{\text{Yield}}{\text{Area Harvested}}$                                  | Water Used, Scope 1 Emissions | Area Harvested, Year, GI |
| Model 3 | Yield $\times$ Average Sale Price   | Water Used, Scope 1 Emissions | Area Harvested, Year, GI |
| Model 4 | $\frac{\text{Yield} \times \text{Average Sale Price}}{\text{Area Harvested}}$ | Water Used, Scope 1 Emissions | Area Harvested, Year, GI |

## 2.1 Data

The data used in this analysis was provided by Sustainable Winegrowing Australia, Australia’s national wine industry sustainability program. Sustainable winegrowing Australia aims to facilitate grape-growers and winemakers in demonstrating and improving their sustainability [25]. Data recorded by Sustainable Winegrowing Australia is entered manually by winegrowers using a web based interface; with some fields being optional. Two subsets of this data were defined by vineyards that recorded values for average price of grape sales per tonne, and vineyards who did not. Both subsets contained: region, harvest year, yield, area harvested, water used and fuel used (in litres for diesel, petrol, biodiesel and LPG). To enable comparisons, total fuel was converted to amount of carbon emissions in metric tons.

The first subset of data was used for Model 1 and Model 2 (see Table 1). This subset contained 5298 samples spanning the period from 2012 to 2022, covering 57 GI Regions and 1432 separate vineyards.

The second subset of data, was limited to vineyards that recorded a value for their average sale price of grapes per tonne. This subset was used for Model 3 and Model 4 (see Table 1); and contained 2878 samples spanning the period from 2015 to 2022, covering 51 GI Regions and 944 separate vineyards.

Data was limited to samples that had recorded values for variables used (see Table 1). After reviewing correlation coefficients the data was logarithmically transformed, centred and scaled by standard deviation. Two values for average sale price were removed from the dataset, due to a recording of \$1. Unreported values for average prices per tonne were filled in using regional averages taken from Wine Australia’s annual reports, where they were

available [26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36].

Other variables including the use of renewable energy, contractors; and pressures such as frost, fire and disease were also explored. Variables that did not significantly contribute to the prediction of a response variable were excluded.

## 2.2 Total Emissions

Emissions were calculated from the total diesel, petrol, bio-diesel and LPG used for irrigation and activities within the vineyard. The equations given from the Australian National Greenhouse Accounts Factors, shown as

$$tCO_2e = \frac{Q \times EC \times EF1 + EF3}{1000}, \quad (1)$$

was used to convert the quantity of fuel in litres,  $Q$ , using a prescribed Energy Content,  $EC$ , and emission factors of scope one,  $EF1$ , and scope three,  $EF3$ , to tonnes of Carbon Dioxide equivalent,  $tCO_2e$  [6].

The variables were reviewed for correlations by using a Pearson’s Correlation Coefficient (see Tables 1, 2 and 3). This was undertaken for data on the original scale (see Table 1) and for data as a logarithmic transform (see Table 2). All P-values were found to be significant ( $p < 2.200E-16$ ), except the non-transformed values for water used (see Table 3). The logarithmic transforms performed the best due to a skew likely caused by a greater number of smaller vineyards within the dataset (see Table 4).

## 2.3 Region

The site of a vineyard predetermines several physical parameters such as climate, geology and soil; making location a widely considered key determinant of grape yield and quality [2, 3, 7]. Differences in vineyard locations were captured through the use of Geographical Indicator Regions (GI Regions). Each GI Region has its own unique mixture of climatic and geophysical properties that describes a unique winegrowing region within Australia; these regions were predefined by Wine Australia [10, 20, 22].

The climatic properties of a GI Region are summarised in the Sustainable Winegrowing Australia user manual [24]. The user manual describes climates by rainfall and temperature, creating supersets of Regions of similar climatic

properties. The climatic groups were used to illustrate similarities and differences occurring in areas larger than GI regions.

## **2.4 Analysis**

General Linear Models were used as they offered the ability to produce statistical models that were explicit in the relationships between predictors and response variables. They also allowed the exploration of interactions between predictors and easily comparable differences in the influence and magnitude of relationships.

Data preprocessing, such as logarithmic transforms, was done using the Python programming language [8]. Linear models were created using the R statistical programming language [21]. These models were created iteratively to explore a variety of variable interactions and approaches to modelling the data. Not all explored approaches yielded improvements or accurate models. Alternate approaches included the use of Splines, hierarchical regression, Additive and Generalised Linear Models. Other variables were also explored but not used due to low reporting values such as fertiliser, tractor and contractor use. The use of only scope one emissions was due to the same reason where scope 2 sources were recorded sporadically at best.

## **2.5 Model Validation**

Models were validated using K-fold cross validation calculated through the R Caret Package [17]. K-fold cross validation works by removing a subset of data from the sample used to train models and then predicts those variables to determine how sensitive the model is to changes in the sample data. For this analysis each model was validated using 10 folds, repeated 100 times.

# **3 Results**

## **3.1 Exploratory Analysis**

Simple linear relationships between variables were explored using Pearson Correlation Coefficients. This was undertaken for data on the original scale (see Table 2) and for data as a logarithmic transform (see Table 3). Strong

Table 2: Summary of models, their predictors, covariates and variable interactions.

| Variable                               | Yield      | Area       | Water Used | Scope One Emissions | $\frac{Yield}{Area}$ |
|--|------------|------------|------------|---------------------|----------------------|
| Yield                                  | 1.000E+00  | 7.440E-01  | -4.309E-03 | 7.290E-01           | 3.500E-01            |
| Area                                   | 7.440E-01  | 1.000E+00  | -5.331E-03 | 8.921E-01           | 7.854E-02            |
| Water Used                             | -4.309E-03 | -5.331E-03 | 1.000E+00  | -1.929E-03          | -5.600E-03           |
| Scope One Emissions                    | 7.290E-01  | 8.921E-01  | -1.929E-03 | 1.000E+00           | 9.357E-02            |
| $\frac{Yield}{Area}$                   | 3.500E-01  | 7.854E-02  | -5.600E-03 | 9.357E-02           | 1.000E+00            |
| Average Price Per Tonne                | -2.262E-01 | -1.178E-01 | -3.562E-02 | -9.422E-02          | -4.844E-02           |
| $\frac{Average Price per tonne}{Area}$ | -1.644E-01 | -2.042E-01 | -2.669E-02 | -1.933E-01          | -1.698E-01           |

Table 3: Pearson correlation coefficients for each logarithmically transformed variable.

| Variable                               | Yield      | Area       | Water Used | Scope One Emissions | $\frac{Yield}{Area}$ |
|--|------------|------------|------------|---------------------|----------------------|
| Yield                                  | 1.000E+00  | 8.822E-01  | 8.245E-01  | 7.617E-01           | 9.353E-01            |
| Area                                   | 8.822E-01  | 1.000E+00  | 7.750E-01  | 8.311E-01           | 6.742E-01            |
| Water Used                             | 8.245E-01  | 7.750E-01  | 1.000E+00  | 6.668E-01           | 7.292E-01            |
| Scope One Emissions                    | 7.617E-01  | 8.311E-01  | 6.668E-01  | 1.000E+00           | 6.086E-01            |
| $\frac{Yield}{Area}$                   | 9.353E-01  | 6.742E-01  | 7.292E-01  | 6.086E-01           | 1.000E+00            |
| Average Price Per Tonne                | -4.591E-01 | -1.911E-01 | -4.881E-01 | -1.559E-01          | -5.625E-01           |
| $\frac{Average Price per tonne}{Area}$ | -8.918E-01 | -8.474E-01 | -8.300E-01 | -7.063E-01          | -8.070E-01           |

relationships were found to be present, as all P-values were considered significant ( $p < 2.200E-16$ , see Tables 2 and 3), except for the non-transformed values for water used (see Table 4). The logarithmic transforms showed the strongest correlations, this was likely due to a skew caused by a greater number of smaller vineyards within the dataset (see Table 5).

### 3.2 General Linear Models

Models 1 and 2 showed significant relationships between each of the predictors and their response variable (see Tables 6 and 7). Variables in models 3 and 4 reported similar significance; except for scope 1 emissions (see Tables 8 and 9). Scope one emissions was included in all models to directly compare the response variables as ratios of vineyard size to raw values. Even though not significant within models 3 and 4, when using the Pearson Correlation Coefficients, scope one emissions was strongly correlated to every Model's re-

Table 4: P-values for the non-transformed water used variable's Pearson correlation coefficients.

| Variable   | Water Used |
|--|------------|
| Yield  | 7.538E-01  |
| Area   | 6.981E-01  |
| Scope One Emissions                                  | 8.883E-01  |
| $\frac{\text{Yield}}{\text{Area}}$                   | 6.836E-01  |
| Average Price Per Tonne                              | 5.600E-02  |
| $\frac{\text{Average Price per tonne}}{\text{Area}}$ | 1.522E-01  |

Table 5: Summary statistics for each variable on the original scale..

| Variable   | Yield      | Area       | Water Used | Scope One Emissions | $\frac{\text{Yield}}{\text{Area}}$ |
|--|------------|------------|------------|---------------------|------------------------------------|
| Yield  | 1.000E+00  | 8.822E-01  | 8.245E-01  | 7.617E-01           | 9.353E-01                          |
| Area   | 8.822E-01  | 1.000E+00  | 7.750E-01  | 8.311E-01           | 6.742E-01                          |
| Water Used   | 8.245E-01  | 7.750E-01  | 1.000E+00  | 6.668E-01           | 7.292E-01                          |
| Scope One Emissions                                  | 7.617E-01  | 8.311E-01  | 6.668E-01  | 1.000E+00           | 6.086E-01                          |
| $\frac{\text{Yield}}{\text{Area}}$                   | 9.353E-01  | 6.742E-01  | 7.292E-01  | 6.086E-01           | 1.000E+00                          |
| Average Price Per Tonne                              | -4.591E-01 | -1.911E-01 | -4.881E-01 | -1.559E-01          | -5.625E-01                         |
| $\frac{\text{Average Price per tonne}}{\text{Area}}$ | -8.918E-01 | -8.474E-01 | -8.300E-01 | -7.063E-01          | -8.070E-01                         |

Table 6: Model 1 ANOVA summarising variable significance at the .5 level.

| Variable            | Df | Sum Sq    | Mean Sq   | F Value   | Pr(>F)        |
|---------------------|----|-----------|-----------|-----------|---------------|
| Year                | 9  | 7.060E+01 | 7.800E+00 | 8.353E+01 | <2.20E-16 *** |
| GI Region           | 54 | 1.507E+03 | 2.790E+01 | 2.972E+02 | <2.20E-16 *** |
| Area Harvested      | 1  | 3.211E+03 | 3.211E+03 | 3.419E+04 | <2.20E-16 *** |
| Water Used          | 1  | 1.040E+01 | 1.040E+01 | 1.103E+02 | <2.20E-16 *** |
| Scope One Emissions | 1  | 6.600E+00 | 6.600E+00 | 7.056E+01 | <2.20E-16 *** |

Table 7: Model 2 ANOVA summarising variable significance at the .5 level.

| Variable                           | Df  | Sum Sq    | Mean Sq   | F Value   | Pr(>F)      |
|------------------------------------|-----|-----------|-----------|-----------|-------------|
| Area Harvested                     | 1   | 2.407E+03 | 2.407E+03 | 1.080E+04 | <2.20E-16   |
| Scope One Emissions                | 1   | 3.989E+01 | 3.989E+01 | 1.789E+02 | <2.20E-16   |
| Water Used                         | 1   | 5.500E+02 | 5.500E+02 | 2.467E+03 | <2.20E-16   |
| Area Harvested*Scope One Emissions | 1   | 6.921E+01 | 6.921E+01 | 3.104E+02 | <2.20E-16   |
| Area Harvested * Water Used        | 1   | 1.040E+00 | 1.040E+00 | 4.686E+00 | 3.045E-02 * |
| Year * GI Region                   | 424 | 1.144E+03 | 2.700E+00 | 1.210E+01 | <2.20E-16   |

sponse variable; this was especially so for Model 1 and 4 (Yield and average price per tonne as a ratio to area harvested, respectively).

The comparison of models performance shows that the average price per tonne of grapes describes a great deal of the relationship between predictor and response when comparing model 2 to model 4 (see Table 10). This relationship between yield and average price was also illustrated in the correlation values between them (see Table 2).

Limitations included overestimating yield for models 1 and 2, (see Figures 1 and 2) and underestimating crop value in models 3 and 4 (see Figures 3 and 4). Reviewing the data to uncover reasons for this included the use of binary

Table 8: Model 3 ANOVA summarising variable significance at the .5 level.

| Variable            | Df | Sum Sq    | Mean Sq   | F Value   | Pr(>F)        |
|---------------------|----|-----------|-----------|-----------|---------------|
| Year                | 6  | 1.324E+01 | 2.210E+00 | 8.748E+01 | <2.20E-16 *** |
| GI Region           | 50 | 6.498E+02 | 1.300E+01 | 5.151E+02 | <2.20E-16 *** |
| Area Harvested      | 1  | 2.142E+03 | 2.142E+03 | 8.491E+04 | <2.20E-16 *** |
| Water Used          | 1  | 3.200E-01 | 3.200E-01 | 1.259E+01 | 3.947E-04 **  |
| Scope One Emissions | 1  | 4.000E-02 | 4.000E-02 | 1.492E+00 | 2.221E-01     |

Table 9: Model 4 ANOVA summarising variable significance at the .5 level.

| Variable                           | Df  | Sum Sq    | Mean Sq   | F Value   | Pr(>F)    |
|------------------------------------|-----|-----------|-----------|-----------|-----------|
| Area Harvested                     | 1   | 2.066E+03 | 2.066E+03 | 5.700E+04 | <2.20E-16 |
| Scope One Emissions                | 1   | 6.000E-02 | 6.000E-02 | 1.569E+00 | 2.105E-01 |
| Water Used                         | 1   | 2.014E+02 | 2.014E+02 | 5.557E+03 | <2.20E-16 |
| Area Harvested*Scope One Emissions | 1   | 5.246E+01 | 5.246E+01 | 1.448E+03 | <2.20E-16 |
| Area Harvested * Water Used        | 1   | 7.270E+00 | 7.270E+00 | 2.005E+02 | <2.20E-16 |
| Year * GI Region                   | 243 | 4.546E+02 | 1.870E+00 | 5.162E+01 | <2.20E-16 |

Table 10: Comparison of Model Residuals

|         | Df   | Sum Sq    | Mean Sq   |
|---------|------|-----------|-----------|
| Model 1 | 5231 | 4.913E+02 | 1.000E-01 |
| Model 2 | 4868 | 1.085E+03 | 2.200E-01 |
| Model 3 | 2818 | 7.111E+01 | 3.000E-02 |
| Model 4 | 2629 | 9.528E+01 | 4.000E-02 |

Table 11: Comparison of Model performance.

|         | RSE       | R2        | Adjusted R2 | F-statistic | P-Value  |
|---------|-----------|-----------|-------------|-------------|----------|
| Model 1 | 3.065E-01 | 9.072E-01 | 9.061E-01   | 7.753E+02   | <2.2e-16 |
| Model 2 | 4.722E-01 | 7.951E-01 | 7.770E-01   | 4.403E+01   | <2.2e-16 |
| Model 3 | 1.589E-01 | 9.753E-01 | 9.748E-01   | 1.885E+03   | <2.2e-16 |
| Model 4 | 1.904E-01 | 9.669E-01 | 9.638E-01   | 3.095E+02   | <2.2e-16 |



variables such as the utilisation of renewable energy, contractors, and the occurrence of disease, fire and frost; however none of these variables were able to explain why some vineyards produced less, or why other vineyards sold at higher prices than predicted. A wide variety of these influences were likely already explained within the use of year and GI Region, or the interaction of both variables. The change between some regions was dramatic, with particularly warmer and drier regions producing much higher volumes of grapes at lower prices (See Figures 5 and 6). The use of other variables and methods, specifically splines, were able to create a more normally distributed set of residuals but at a drastically reduced accuracy when comparing R2 and RSE. The introduction of known average prices per tonne also helped increase R2 values a small amount; it is important to not that it is common practice for wineries to purchase grapes at a regional average rate, likely resulting in much less variance within a region.

The correlation between average sales price and yield was a negative trend (see table 2); the contributing factors to yield and average sales price was ???. Correlation values showed that water and emissions increased with yield but decreased with average sale price (see Table 4). In alternative attempts at models it was found that without the incorporation of GI Region or year the predictions greatly under performed. The possible reason behind this effect was that different strategies are likely employed between different regions, where some regions target the mass production of cheaper grapes over quality. This is most notable when grouping regions by climate, especially when considering GI Regions in the 'Hot Very Dry' climate (see Figure 7). The effect of climate in the models was not more significant than the more granular use of GI regions. The interaction between year and GI Region likely accounted for localised events such as bushfires, which would be impactful, but only at a local level in both time and space.

### 3.3 Model Validation

To validate the performance of these models k-fold cross validation was used. This was done using 10 folds, k=10, repeated 100 times. The models performed similarly to their original counter parts (see Table 11).

Table 12: Model validation using k-fold cross validation, for 10 folds repeated 100 times.

|         | RMSE      | R2        | MAE       |
|---------|-----------|-----------|-----------|
| Model 1 | 3.087E-01 | 9.045E-01 | 2.165E-01 |
| Model 2 | 5.104E-01 | 7.409E-01 | 3.493E-01 |
| Model 3 | 1.652E-01 | 9.723E-01 | 1.008E-01 |
| Model 4 | 2.235E-01 | 9.500E-01 | 1.279E-01 |

## 4 Discussion

This study investigated the general relationships between input resources of a vineyard, including fuel and water, and the outputs including yield and value. Some regions appeared to produce many low quality grapes at scale compared to attempting to produce fewer higher quality grapes. This behaviour can be observed when reviewing Wine Australia’s annual reports, where it is apparent that warm inland regions such as the Riverland are known to only produce large amounts of lower graded grapes [29, 35]. Comparatively, regions such as Tasmania only produce A grade grapes but in much smaller quantities than the Riverland. Knowing that the difference in pricing per tonne can exceed a magnitude of 10 between grades E and A, the operations in regions that target different grades would have varied priorities. However, some regions such as the Yarra Valley produce a Variety of different grades of grapes, from C to A, highlighting that vineyard priorities, although may be somewhat present within regional classifications, are not necessarily aligned within a given region.

The opportunity to target different grades of grapes may not always be available, with some regions being more renowned than others, and likely to be sought after regardless [10]. The Barossa is an example of this, known for its quality could also lend itself to a bias in purchasers not considering other regions that may be capable of similar quality. This effect could stifle the potential for market opportunities within these lesser known regions. A further possibility is that there may be regional upper limits with the relationship between resource input and the value gained becoming no longer proportional due to diminishing returns. Climate was considered to be a large determinant of the ability to grow a larger quantity of grapes, as well as a determinant in grape quality [3]; however there were vineyards in similar regions that were able to produce exceptionally better results than others

(See Figure 7).

The issue of model 1 and 2 over predicting yield, may have been due to preventative measures brought on by regional pressures such as fire, frost and disease. Where, more resources were required to prevent these issues from spreading within a region, thus disproportionately effecting some vineyards compared to others locally. This type of maintenance is not well captured especially when considering that some regions, those in warmer areas are not as prone to disease as cooler climates and could potentially have lower operating costs per hectare. This could create a discrepancy in vineyards that utilise preventative measures in wetter regions, as opposed to those who do not, and thus expend less fuel and energy but risk disease. When reviewing the differences between regions it is important to consider that vineyards in Hot Very Dry areas can be hundreds of times the size of those in other regions. It is interesting that while area, although significantly correlated to the ratio of yield to area, was still lower than water and about the same as emissions. This points to economies of scale playing a role but still being only one consideration alongside the potential resources that can be used. The negative trend between size and average sales price could also be a side effect of mass supply verse demand, especially when looking at the level of difference in production of some vineyards (see Table 4). The relationships between yield, value and area are not simply about efficiently producing the most grapes; sales price and by association grape quality, are integral to the profitability, and this is strongly linked to resource use and thus the longevity and sustainability of a vineyard.

Literature shows that there are many on-the-ground decisions that influence both quality and yield. Where these decisions are governed by complex physical and social forces such as international market demands, disease pressures and natural disasters [1, 5, 9, 12, 13, 20, 23]. Many of these occurrences being highlighted throughout the past decades vintage reports [26, 28, 29, 31, 32, 33, 34, 35, 36]. It is also important to consider that these reports show that the warm inland regions have seen a decline in profit during this period, as they were often compared to other regions that focused more on quality than quantity. This is an important consideration, as the size of some of these vineyards when considering their ratio of value to area would only require a marginal increase to out compete other regions. There are also differences when comparing winegrowers to other agricultural industries as they are vertically integrated within the wine industry, tying them to secondary and tertiary industries, such as wine production, packaging,

transport and sales. This results in unique issues and considerations for each vineyard, where these on-the-ground decisions may be influenced by other wine industry’s choices, such as the use of sustainable practices in vineyards as a requirement for sale in overseas markets; notably these interactions are further complicated by some winegrowers being totally integrated into wine companies, while others are not (Knight et al., 2019). Incorporating such decisions into the model could help describe the contributing factors to regional differences beyond resource consumption and regional differences.

Having more data for each region would also be an improvement, allowing greater comparison between regions. More variables may also help to discern vineyards that can produce larger volumes of grapes at higher prices. The use of semi transparent tools such as random forests and decision trees alongside more variables and data may help to uncover the reasons for values that were under or over estimated. These differences could be caused by the use of alternative sustainable practices in the field. While there is evidence to suggest that environmentally sustainable practices can reduce costs, increase efficiency, whilst improving the quality of grapes, more research is needed to link these benefits across different regions and climates [4, 18, 19].

The relationship between scope one emissions and the response variables that included average sales price

It is possible that the relationships between scope one emissions and the response variables were closely tied to a vineyards area. This possibility could be explained through the emissions

Noting that irrigation systems use fuel and that the application of water was a significant variable in each model scope one emissions’ lack of significance and contribution given its F-statistics (See Tables 7 and 8), indicated that it is possible other vineyard activities requiring fuel are not as determining factors for a vineyards grape quality.

## References

- [1] Javier Abad et al. ‘Cover Crops in Viticulture. A Systematic Review (1): <br>Implications on Soil Characteristics and Biodiversity in Vineyard’. In: *OENO One* 55.1 (15 Mar. 2021), pp. 295–312. DOI: 10 . 20870/oeno-one . 2021 . 55 . 1 . 3599. URL: <https://oeno-one.eu/article/view/3599> (visited on 14/02/2023).

- [2] Philippe Abbal et al. ‘Decision Support System for Vine Growers Based on a Bayesian Network’. In: *Journal of agricultural, biological, and environmental statistics* 21.1 (2016), pp. 131–151. ISSN: 1085-7117. DOI: 10.1007/s13253-015-0233-2.
- [3] Eduardo Agosta, Pablo Canziani and Martín Cavagnaro. ‘Regional Climate Variability Impacts on the Annual Grape Yield in Mendoza, Argentina’. In: *Journal of Applied Meteorology and Climatology* 51.6 (2012), pp. 993–1009.
- [4] Antonietta Baiano. ‘An Overview on Sustainability in the Wine Production Chain’. In: *Beverages* 7.1 (2021). ISSN: 2306-5710. DOI: 10.3390/beverages7010015.
- [5] Paulo Cortez et al. ‘Using Data Mining for Wine Quality Assessment’. In: *Discovery Science: 12th International Conference, DS 2009, Porto, Portugal, October 3-5, 2009 12*. Springer, 2009, pp. 66–79. ISBN: 3-642-04746-7.
- [6] Department of Climate Change, Energy, the Environment and Water. *Australian National Greenhouse Accounts Factors*. Nov. 2022. URL: <https://www.dcceew.gov.au/sites/default/files/documents/national-greenhouse-accounts-factors-2022.pdf>.
- [7] Helder Fraga, Ricardo Costa and João A. Santos. ‘Multivariate Clustering of Viticultural Terroirs in the Douro Winemaking Region’. In: *Ciência Téc. Vitiv.* 32.2 (2017), pp. 142–153. URL: <https://doi.org/10.1051/ctv/20173202142>.
- [8] G. van Rossum. *Python Tutorial, Technical Report CS-R9526*. Version 3.10.6. Amsterdam, The Netherlands: Centrum voor Wiskunde en Informatica (CWI), 1995.
- [9] Andrew Hall et al. ‘Within-Season Temporal Variation in Correlations between Vineyard Canopy and Winegrape Composition and Yield’. In: *Precision Agriculture* 12 (2011), pp. 103–117.
- [10] J. C. (James C.) Halliday. *Australian Wine Encyclopedia*. VIC: Hardie Grant Books, 2009. ISBN: 978-1-74273-245-9.
- [11] Silke Hemming et al. ‘Cherry Tomato Production in Intelligent Greenhouses-Sensors and Ai for Control of Climate, Irrigation, Crop Yield, and Quality’. In: *Sensors (Basel, Switzerland)* 20.22 (2020), pp. 1–30. ISSN: 1424-8220. DOI: 10.3390/s20226430.

- [12] I. Goodwin et al. *Managing Soil and Water to Target Quality and Reduce Environmental Impact*. 2009. URL: <https://www.wineaustralia.com/getmedia/6dd0cbc3-1db4-43cf-a908-9c9d2a1070cc/DPI-04-04-smaller>.
- [13] Aikaterini Kasimati et al. ‘Predicting Grape Sugar Content under Quality Attributes Using Normalized Difference Vegetation Index Data and Automated Machine Learning’. In: *Sensors* 22.9 (2022). ISSN: 1424-8220. DOI: 10.3390/s22093249.
- [14] Kentaro Kawasaki and Shinsuke Uchida. ‘Quality Matters More Than Quantity: Asymmetric Temperature Effects on Crop Yield and Quality Grade’. In: *American journal of agricultural economics* 98.4 (2016), pp. 1195–1209. ISSN: 0002-9092. DOI: 10.1093/ajae/aaw036.
- [15] Keith Jones. *Australian Wine Industry Environment Strategy*. 27 Sept. 2002.
- [16] Hanne Knight et al. ‘Firm Resources and the Development of Environmental Sustainability among Small and Medium-Sized Enterprises: Evidence from the Australian Wine Industry’. In: *Business Strategy and the Environment* 28.1 (1 Jan. 2019), pp. 25–39. ISSN: 0964-4733. DOI: 10.1002/bse.2178. URL: <https://doi.org/10.1002/bse.2178> (visited on 13/02/2023).
- [17] Max Kuhn. ‘Building Predictive Models in R Using the Caret Package’. In: *Journal of Statistical Software, Articles* 28.5 (2008), pp. 1–26. ISSN: 1548-7660. DOI: 10.18637/jss.v028.i05. URL: <https://www.jstatsoft.org/v028/i05>.
- [18] Angela Mariani and Antonella Vastola. ‘Sustainable Winegrowing: Current Perspectives’. In: *International Journal of Wine Research* 7.1 (2015), pp. 37–48.
- [19] Johnny V. Montalvo-Falcón et al. ‘Sustainability Research in the Wine Industry: A Bibliometric Approach’. In: *Agronomy* 13.3 (2023). ISSN: 2073-4395. DOI: 10.3390/agronomy13030871.
- [20] D.P. Oliver et al. ‘Review: Soil Physical and Chemical Properties as Indicators of Soil Quality in Australian Viticulture’. In: *Australian Journal of Grape and Wine Research* 19.2 (1 June 2013), pp. 129–139. ISSN: 1322-7130. DOI: 10.1111/ajgw.12016. URL: <https://doi.org/10.1111/ajgw.12016> (visited on 13/02/2023).

- [21] R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing, 2021. URL: <https://www.R-project.org/>.
- [22] C.J SOAR, V.O SADRAS and P.R PETRIE. ‘Climate Drivers of Red Wine Quality in Four Contrasting Australian Wine Regions’. In: *Australian journal of grape and wine research* 14.2 (2008), pp. 78–90. ISSN: 1322-7130. DOI: 10.1111/j.1755-0238.2008.00011.x.
- [23] Satyam Srivastava and Shashikant Sadistap. ‘Non-Destructive Sensing Methods for Quality Assessment of on-Tree Fruits: A Review’. In: *Journal of Food Measurement and Characterization* 12.1 (2018), pp. 497–526.
- [24] Sustainable Winegrowing Australia SWA. *Sustainable Winegrowing Australia User Manual*. Aug. 2021. URL: <https://www.awri.com.au/wp-content/uploads/2019/07/SWA-user-manual.pdf>.
- [25] Sustainable Winegrowing Australia SWA. *Sustainable Winegrowing Australia*. Case Studies. 2022. URL: <https://sustainablewinegrowing.com.au/case-studies/> (visited on 05/11/2022).
- [26] Wine Australia. ‘National Vintage Report 2019’. In: (2019). URL: [https://www.wineaustralia.com/getmedia/807bf053-3692-448a-9ed5-c0084a47e1bb/Vintage-report-2019\\_full-version.pdf](https://www.wineaustralia.com/getmedia/807bf053-3692-448a-9ed5-c0084a47e1bb/Vintage-report-2019_full-version.pdf).
- [27] Wine Australia. ‘National Vintage Report 2020’. In: (2020). URL: [https://www.wineaustralia.com/getmedia/7dcd66a7-86b5-4606-9ab7-416086077099/MI\\_VintageReport2020.pdf](https://www.wineaustralia.com/getmedia/7dcd66a7-86b5-4606-9ab7-416086077099/MI_VintageReport2020.pdf).
- [28] Wine Australia. ‘National Vintage Report 2021’. In: (2021). URL: [https://www.wineaustralia.com/getmedia/100ffcfcd-5672-49eb-8b5e-ac31cd2f57b3/MI\\_VintageReport2021\\_full-\(1\)\\_1.pdf](https://www.wineaustralia.com/getmedia/100ffcfcd-5672-49eb-8b5e-ac31cd2f57b3/MI_VintageReport2021_full-(1)_1.pdf).
- [29] Wine Australia. ‘National Vintage Report 2022’. In: (2022). URL: [https://www.wineaustralia.com/getmedia/b61c7a0e-9e17-4142-a98b-945e3f18873e/MI\\_VintageReport2022\\_full.pdf](https://www.wineaustralia.com/getmedia/b61c7a0e-9e17-4142-a98b-945e3f18873e/MI_VintageReport2022_full.pdf).
- [30] Winemakers’ Federation of Australia. ‘National Vintage Report 2012’. In: (2012). URL: <https://www.agw.org.au/assets/vintage-reports/PDFs/WFA-Vintage-Report-2012.pdf>.

- [31] Winemakers' Federation of Australia. 'National Vintage Report 2013'. In: (2013). URL: <https://www.agw.org.au/assets/vintage-reports/PDFs/WFA-Vintage-Report-2013.pdf>.
- [32] Winemakers' Federation of Australia. 'National Vintage Report 2014'. In: (2014). URL: <https://www.agw.org.au/assets/vintage-reports/Vintage-Report-and-Production-Profitability-2014.pdf>.
- [33] Winemakers' Federation of Australia. 'National Vintage Report 2015'. In: (2015). URL: <https://www.agw.org.au/assets/vintage-reports/Vintage-Report-and-Production-Profitability-2015.pdf>.
- [34] Winemakers' Federation of Australia. 'National Vintage Report 2016'. In: (2016). URL: <https://www.agw.org.au/assets/vintage-reports/Vintage-Report-2016.pdf>.
- [35] Winemakers' Federation of Australia. 'National Vintage Report 2017'. In: (2017). URL: <https://www.agw.org.au/assets/vintage-reports/NationalVintageReport-2017.pdf>.
- [36] Winemakers' Federation of Australia. 'National Vintage Report 2018'. In: (2018). URL: [https://www.wineaustralia.com/getmedia/fce5bf18-468b-4e1b-a478-d6ca231d40f6/VintageReport2018\\_full.pdf](https://www.wineaustralia.com/getmedia/fce5bf18-468b-4e1b-a478-d6ca231d40f6/VintageReport2018_full.pdf).
- [37] Da-wei ZHU et al. 'Effects of Nitrogen Level on Yield and Quality of Japonica Soft Super Rice'. In: *Journal of Integrative Agriculture* 16.5 (2017), pp. 1018–1027. ISSN: 2095-3119. DOI: 10.1016/S2095-3119(16)61577-0.