

The influence of resource use on yield versus sale price trade-off in Australian vineyards

Bryce Boyd^{1,2,3,*}, Kate Helmstedt¹, Mardi Longbottom², Kerrie Mengersen¹

1 Affiliation Faculty of Mathematical Sciences, Queensland University of Technology, Brisbane, QLD, Australia

2 Affiliation Australian Wine Research Institute, Adelaide, SA, Australia

3 Affiliation Food Agility CRC, Sydney, NSW, Australia

* Bryce.polley@awri.com.au

Abstract

Strategies for achieving sustainability in the winegrowing industry requires balancing resource investment against the economic outcomes of resultant yields and sale price of the produce. Although there has been much research into the forecasting of outcomes and resource use, little has been done to illustrate the effects they have on one another and the consequential economic outcomes. In this analysis we observe relationships between resource use, yield and sale price through the use of statistical models. The dataset used for this analysis includes data collected for the past 10 years from 1261 vineyards located over a diverse range of Australian winegrowing regions. Yield and sale price were evaluated with respect to the resource use factors water use and Green House Gas (GHG) emissions. The analysis confirmed a strong relationship between area and resource use, with the overall area of a vineyard and its access to resources greatly determining the upper limit of yield. However, area was also negatively related to the average sale price of grapes; we find that higher average sale prices were connected to high resource inputs per area, rather than to the overall expenditure of resources. Regional and temporal effects on vineyard yield and average sales price were also identified. Overall, the analysis highlighted the importance of considering a vineyard's business goal, region, external pressures and economies of scale, when considering whether to pursue higher yields or higher average sales prices.

Introduction

The focus on sustainability in agronomic industries has changed the way in which these enterprises do business. The diverse nature of vineyard operations results in complicated interactions with both environmental and economic outcomes. The interactions between different factors can put an environmentally advantageous decision at odds with an economic one, and conversely economic decisions can be detrimental to the environment [1]. As a result many studies are including more diverse ranges of data and scales. However, studies tend to focus on specific aspects, such as reducing chemical use or improving soil health, often focusing on singular relationships or variable outcomes [2–4]. Other analyses, look at life cycles or cost benefit analyses, reflecting a specific management decision's outcome [5–7] or their cost benefits by utilising different scenarios to assess specific practices [8,9]. It has been noted that some studies have relied too heavily on single points of engagement when relaying the idea of sustainability

and vineyard outcomes, especially when the inclusion of more variables or data could have better described a situation as a whole [10, 11]. The changing nature of sustainability also calls for measures to be more than simple snapshots of data [10–12]. Given recent studies there is still growing interest from the winegrowing industry in further studies, with several recent reviews calling for more research to demonstrate the potential trade-offs between different operational approaches across different climates and regions [10, 12, 13].

A dilemma exists across agriculture through shared fundamental considerations of resource use such as water and fuel, and the resultant yield and crop value that are produced [14–16]. The average price of grapes for wine production is driven through its integration within the wine industry. An important connection between grapes and their price is the grape's perceived quality, which initially defines a wine's potential through the grape's chemical makeup [17, 18]. Grapes of higher perceived quality or grapes from well known regions are likely to have higher prices [19]. Grape quality is connected to the market value of winegrapes, with Wine Australia explicitly defining grape quality through the use of discrete price brackets in their annual reports [20]. It is important to note that the generalisation made to reflect quality through using average price assumes a due diligence of those who purchase the grapes [21]. The economic sustainability of a vineyard is tied to this market culture, driven by the wine industry. The consideration of sustainability within viticulture is also subject to environmental and sociodemographic pressures [11]. In the Australian context, these pressures include biosecurity, climate and international market changes [22–24].

There is an extensive amount of research into the varied effects of factors on grape quality and yield [3, 25, 26]. Due to the lack of long-term and in-depth data there is a lack of research on grape sale price and its driving factors. Furthermore, individual factors are often studied in isolation with yield and sales price not appearing together [27]. The lack of consolidated datasets restricts the ability to gain statistical insights at large scales and across multiple regions. As a result, broader studies are lacking [28, 29]. The dataset used for this analysis includes data spanning 10 years from a multitude of vineyards located over a diverse range of Australian winegrowing regions. We use this dataset to describe the relationship of resources related to water and fuel use with the output yield and average sale price of the resultant product, taking into account the size and location of the vineyard. In a practical sense, this aim is a baseline for comparison: given a vineyard within Australia, one could estimate the comparative efficiency with regard to the trade-off between invested resources, yield and average sale price. This is the first time that such a trade-off has been confirmed explicitly across such varying regions, scales and climates in the Australian winegrowing industry.

Methods

Data

Data used in these analyses were obtained from Sustainable Winegrowing Australia and Wine Australia. Sustainable Winegrowing Australia is Australia's national wine industry sustainability program, which aims to facilitate grape growers and winemakers in demonstrating and improving their sustainability [30]. Wine Australia is an Australian Government statutory authority governed by the Wine Australia Act [31]. Data collected by Wine Australia are publicly available. The response (predicted) variables in this analysis were yield, defined as the total tonnes of grapes harvested, and average sale price of grapes in Australian dollars per tonne. Both response variables were examined as totals and also relative to area harvested (see Table 1). Values were compared in this manner to observe how economies of scale affect the use of resources.

Table 1. Summary of models; their predictors, covariates and variable interactions.

	Response	Predictors	Covariates	Interactions
Model 1	Yield	Water Used scope one Emissions	Area Harvested Year GI Region	N/A
Model 2	$\frac{\text{Yield}}{\text{Area Harvested}}$	Water Used scope one Emissions	Area Harvested Year GI Region	Area Harvested * scope one Emissions Area Harvested * Water Use Year * Region
Model 3	$\text{Yield} \times \text{Average Sale Price}$	Water Used Scope One Emissions	Area Harvested Year GI Region	N/A
Model 4	Average Sale Price	Water Used Scope One Emissions	Area Harvested Year GI Region	Area Harvested * Scope One Emissions Area Harvested * Water Use Year * Region
Model 5	Average Sale Price	Water Used Scope One Emissions	Year GI Region	Year * Region

Data recorded by Sustainable Winegrowing Australia were entered manually by winegrowers voluntarily using a web based interface. For each model, vineyards were only included if they recorded all the variables used for the corresponding model (see Table 1). Each vineyard had at least recorded region, harvest year, yield and area harvested. Other variables that were used but not present for every single vineyard were average sale price, water used and fuel used (diesel, petrol, biodiesel and LPG). To enable direct comparisons between fuels, fuel use was converted to tonnes of carbon dioxide equivalent and collectively referenced to as emissions. All variables were continuous except for harvest year and region, which were categorical variables (Table 1).

As data from Sustainable Wine Australia were voluntarily given, missing values were improved using regional average prices from the Wine Australia data. Data obtained from Wine Australia were collected via phone surveys and included: total tonnes purchased, average price per tonne and yearly change in price for region and grape varietal; with the data being publicly available.

The dataset was split into two distinct subsets that could be used across the different models. The first subset contained all vineyards and was used for two models (Model 1 and Model 2, see Table 1). The second subset contained vineyards which either recorded a value for average price of sale per tonne through Sustainable Winegrowing Australia, or were within a region with an average price of sale recorded by Wine Australia; this subset was used for three further models (Models 3, 4 and 5, see Table 1). These subsets meant that the data would be limited to samples which had recorded values for the response variables (see Table 1), as every sample/vineyard had a recorded value for yield but not average price of sale per tonne.

The first subset of data (used for Model 1 and Model 2, see Table 1) contained 5298 samples spanning the period from 2012 to 2022, covering 55 GI Regions and 1261 discrete vineyards.

The second subset of data (used for Model 3, Model 4 and Model 5, see Table 1) contained 2878 samples spanning the period from 2015 to 2022, covering 51 GI Regions and 944 separate vineyards. Average price of sale per tonne was extracted from both Wine Australia (1842 values) and Sustainable Winegrowing Australia (remaining 1036 values).

Additional variables were considered for analysis but were excluded due to being either underreported or had insignificant contributions to model accuracies. Variables explored but not used due to low reporting values included fertiliser (kg), electricity (Kw/h), and scope two emissions (TCO² equivalent). Variables considered but




Fig 1. Map of each GI Regions' (outlined by Wine Australia) average income for a vineyard of that region (average grape sale price per tonne \times total tonnes yielded).

ultimately removed due to a lack of significant contributions to models, included the use of renewable energy, use of contractors, and pressures such as frost, fire and disease.

Data preprocessing was conducted prior to analysis using the Python programming language [32]. Preprocessing included the conversion of fuel to scope one emissions and prior calculations for all continuous variables which included logarithmic transformations, centring and scaling by standard deviation. We converted multiple emission sources into scope one emissions using the equation given from the Australian National Greenhouse Accounts Factors [33]. The calculation conducted using

$$tCO_2e = \frac{Q \times EC \times EF1 + EF3}{1000}, \quad (1)$$

where emissions was the product of the quantity of fuel in litres, Q , a prescribed Energy Content, EC , and emission factors (as given by the Australian National Greenhouse Accounts factors) of scope one, $EF1$, and scope three, $EF3$, to tonnes of Carbon Dioxide Emission equivalent, tCO_2e [34].

Differences in vineyard locations were captured through the use of Geographical Indicator Regions defined by Wine Australia [24, 35, 36]. Although vineyards generally differed in income, there were pronounced differences between regions as shown in

Figure 1. These differences were likely due to the site of a vineyard being important as it predetermines several physical parameters such as climate, geology and soil, making location a widely considered key determinant of grape yield and average sale price [27, 37, 38]. Each GI Region has its own unique mixture of climatic and geophysical properties that describes a unique winegrowing region within Australia and is a protected trademark under the Wine Australia act [31]. Both Wine Australia and Sustainable Winegrowing Australia used the same GI Region categorical variable format to describe location.

The climatic properties of each GI Region were summarised by using predefined classifications as per the [39] user manual. The user manual describes climates by rainfall and temperature, creating supersets of Regions of similar climatic properties. The climatic groups were used to illustrate similarities and differences occurring in areas larger than GI Regions. These classifications were literal descriptors used by industry members to discern between different weather types within winegrowing regions. Other climatic descriptors were explored, such as the Köppen climate classification, however this did not offer a granular enough index to highlight any trends or clustering within the data.

Analysis

Pairwise Pearson Correlation Coefficients (PPCC) were calculated to assess the potential existence of linear relationships between the input and predicted variables. To determine if a coefficient was indicative of a strong relationship, both the magnitude of the value and the associated confidence intervals were evaluated. P-values reflected the significance of a given correlation coefficient with statistical significance being declared when the corresponding P-value was lower than 0.05. PPCC analyses were undertaken for data on the original scale and for data as a logarithmic transform. Transforming data prior to calculating the coefficients changes several things. The logarithmic transform of the data alters the interpretation of the coefficients to percentage change; a coefficient will be indicative of the change in percentage of one variable compared to the other, scaling by standard deviation also changes this interpretation to be a percentage of that variables standard deviation. When considering the logarithmically transformed variables, a coefficient of 1 would indicate that the change of one variable by one percentage of its standard deviation would correlate to the other variable changing by one percent of its own standard deviation. The importance of this is the dimensionless nature of these relationships and that it can be translated directly to any vineyard's case that has a well known distribution.

Five general linear models (GLMs) were created (see Table 1). GLMs were chosen as they offer the ability to produce statistical models that are explicit in the relationships between predictors and response variables. General Linear Models also allowed the exploration of interactions between predictors and allow for easily comparable differences in the influence and magnitude of relationships. Model fit was measured in terms of R^2 and adjusted R^2 as well as F statistics, and t-tests were used to determine if predictors significantly contributed to the models when accounting for other variables, showing which specific years and areas contributed significantly. Both the PPCC and GLM analyses were created using the R statistical programming language [40] with the Caret package [41].

A variety of alternate methods were also explored, including splines, hierarchical regression, general additive models, and generalised linear models. These alternative approaches were not used as final models due to offering little further insights or improvements in accuracy.

Table 2. Summary statistics of each continuous variable.

Variable	Mean	Standard Deviation	Minimum	Maximum
Yield (tonnes)	7.76E+02	2.18E+03	1.00E+00	7.23E+04
Area Harvested (ha)	6.67E+01	1.34E+02	7.00E-02	2.44E+03
Water Used (ML)	7.47E+06	5.65E+08	1.00E+00	4.27E+10
Scope One Emissions (tCO_2e)	4.17E+04	8.57E+04	6.76E+00	2.11E+06
$\frac{\text{Yield (tonnes)}}{\text{Area harvested (ha)}}$	1.01E+01	8.13E+00	4.00E-02	8.63E+01
Average Sale Price (AUD/tonne)	1.48E+03	9.22E+02	1.60E+02	2.60E+04
$\frac{\text{Average Sale Price (AUD/tonne)}}{\text{Area Harvested (ha)}}$	1.35E+02	5.71E+02	1.75E-01	2.98E+04

Model Validation

Models were validated using K-fold cross validation. This was performed by removing a subset of data from the sample used to train models and then predicting those variables to determine how sensitive the model is to changes in the sample data. For this analysis, each model was validated using 10 folds, repeated 100 times.

Results

Summary Statistics

Table 2 shows the summary statistics of each variable in its original units. The range of these values shows the level of difference between some vineyards, with operations differing by orders of magnitude in size, yield and average price of sale (See Table 1).

PPCCs values for the transformed, centred and scaled variables are shown in Table 3. The process of centring and scaling variables changes them so that their mean is centred on 0 and all values are scaled to the variables standard deviation, such that 1 is equal to the variables standard deviation. All correlations were found to be statistically significant ($p < 2.200E-16$), and except for Average Sale Price all variables were positively correlated. With water use, area harvested and emissions being positively correlated to yield, it can be considered that more resources and area are likely to lead to greater yields. The negative correlations between Average Sale Price and Yield, Water Use, Area and Scope One Emissions each indicated that Area Harvested and Fuel Use separately were not the determining factor for average sale price. The negative correlations are associative, not causal relationships (i.e using more water does not cause lower sale prices).

General Linear Models

Each model had a high R^2 value, indicating that most of the variance within the data was described by the models (see Table 4) and statistical significance of F-tests ($p < 2.200E-16$). Aside from 3 variables, all regression parameters were also statistically significant from zero, F-tests across each model's variables were ($p < 0.05$). The three

Table 3. Pairwise Pearson correlation coefficients for logarithmically transformed values.

	Yield	Area Harvested	Water Used	Scope One Emissions	Yield by Area	Average Price	Average Price by Area
Yield	1.00	0.88	0.82	0.76	0.96	-0.46	-0.88
Area Harvested	0.88	1.00	0.78	0.83	0.73	-0.19	-0.81
Water Used	0.82	0.78	1.00	0.67	0.76	-0.49	-0.82
Scope One Emissions	0.76	0.83	0.67	1.00	0.65	-0.16	-0.67
Yield by Area	0.96	0.73	0.76	0.65	1.00	-0.54	-0.84
Average Price	-0.46	-0.19	-0.49	-0.16	-0.54	1.00	0.72
Average Price by Area	-0.88	-0.81	-0.82	-0.67	-0.84	0.72	1.00

Table 4. Summary of models; their performance, F-statistics and Residual error.

	R ²	Adjusted R ²	F-Statistic	P-Value	Residual Standard Error	Residual Sum of Squares	Residual Mean of Squares
Model 1	0.9072	0.9061	775.3	2.200e-16	0.3065	491.3	0.1
Model 2	0.8291	0.8141	55.07	2.200e-16	0.4312	905.03	0.19
Model 3	0.9753	0.9748	1885	2.200e-16	0.1589	71.11	0.03
Model 4	0.9091	0.9006	106.1	2.200e-16	0.3153	261.41	0.10
Model 5	0.9089	0.9004	107.2	2.200e-16	0.3155	262.04	0.10

exceptions were: scope one emissions in Model 3 ($p=0.22$) and Model 4 ($p=0.0.39$), and the interaction between area harvested and water used in model 2 ($p=0.22$). Note that scope one emissions was included in all models to directly compare the response variables as ratios of vineyard size to raw values and because it was strongly correlated to the response variable in every model (except model 5) especially for Models 1 and 4 (Table 3).

Coefficients related to continuous variables are summarised in Table 5. In Model 1 all coefficients except for the intercept significantly contributed to the model ($p < 0.05$), and in Model 2 all coefficients were statistically significant ($p < 0.05$). However, scope one emissions did not significantly contribute to models 3, 4 and 5. Model 4 only had statistically significant contributions from the intercept and water use. Although the coefficient for water use was statistically significant for each model, it did not have the highest value; instead, area harvested, being an order of magnitude greater, dominated the models. Model 5 achieved a similar R^2 to Model 4 without area harvested, having stronger influences from water use and scope one emissions.

Table 5. Summary of each Model's coefficients for continuous variables

	Intercept	Area Harvested	Water Used	Scope One Emissions	Area Harvested	Area Harvested
					** Scope One Emissions	** Water Used
Model 1	Coefficient	-3.32E-02	7.42E-01	8.66E-02	6.73E-02	
	Std Error	1.96E-02	1.00E-02	8.90E-03	8.00E-03	
Model 2	Coefficient	1.70E-01	5.77E-01	1.08E-01	8.50E-02	-4.97E-02
	Std Error	5.91E-02	1.48E-02	1.31E-02	1.17E-02	8.10E-03
Model 3	Coefficient	1.81E-02	9.71E-01	-2.31E-02	-7.00E-03	
	Std Error	1.30E-02	7.20E-03	6.90E-03	5.70E-03	
Model 4	Coefficient	1.45E-01	2.40E-03	-4.66E-02	-1.70E-02	1.15E-02
	Std Error	5.28E-02	1.50E-02	1.43E-02	1.18E-02	7.90E-03
Model 5	Coefficient	1.52E-01		-4.04E-02	-1.71E-02	1.40E-03
	Std Error	5.27E-02		1.13E-02	9.70E-03	8.30E-03

Table 6. Model validation using k-fold cross validation, for 10 folds repeated 100 times.

	Residual Squared Error	Mean R2	Average Error
Model 1	.309	.905	.2165
Model 2	.457	.7921	.313
Model 3	.165	.972	.101
Model 4	.348	.878	.182
Model 5	.348	.878	.183

The regression coefficients for the Year variable of each model is depicted in Figure 2. For each model, the first year for a model's was used as the baseline: 2012 for Models 1 and 2, and 2015 for Models 3, 4 and 5. Adelaide Hills is used as the regional baseline with the interaction between year and region using the first year and the Adelaide Hills as the baseline. Region and year contributed, in some but not all cases, more than the other variables. However, the coefficients varied substantially over the years, and were not significantly different from zero in some years. Models 4 and 5 are very similar in Figure 2, indicating that the exclusion of area does not greatly affect the contribution from yearly influence. Models 4 and 5 have the most prominent trends, showing an increase in yearly effects over time, with Model 3 also increasing from 2016 to 2018, but plateauing afterwards. Model 1 and 2 do not show a clear trend, but do drop during 2017 and 2018 after increasing in the first 3 years.

Regional differences by temperature and rainfall are summarised in Figure 3. The most notable difference is between vineyards within 'Hot' and 'Very Dry' regions (warm inland regions), where high average sale prices are historically low, and yield is high. Water Use changes dramatically between these regions as well, with water being a driving force for yield but not necessarily average sale price. The warmer and drier regions tend to also have larger vineyards. These regional differences are further shown in Figure 4, as a ratio of vineyard; where there is a pronounced difference in 'Hot' and 'Very Dry' regions producing more per area with lower average sale prices per area than other regions.

Figure 4 further shows the emphasis that 'Hot' areas have on high yields, with low average sale price compared with other regions. Scaling average price and yield by area shows a strong negative trend, trading quantity for higher sales prices.

Table ?? shows the validation results of each of the models. The R^2 measures of fit show similar results to the initial models, with a slight decrease as expected, indicating that the models are robust and consistent.

Discussion

There was an expected strong relationship between size and resource use, with the overall area of a vineyard and its access to resources greatly determining the upper limit of potential yield. However, size was also inversely related to the potential average sale price. Higher average sales prices were also related to high resource inputs per area instead of the overall expenditure of resources. Vineyard yields and sales price changed greatly by region and year. Even given regional and yearly changes, there was a strong connection between smaller vineyards and higher sales prices. This could have been due to more attention available when managing smaller properties.

The lack of significance of scope one emissions and its contribution to models, given its F-statistics, could be indicative that other vineyard activities requiring fuel are not leading factors for a vineyards average sale price. The relationship between yield, value




Fig 2. Model Coefficient values for Year, with standard error bars. The coefficients represent the influence that each year had on the model outcomes (see Table 1 for model outcomes and their predictors). Each years influence is comparative to the first year in the models, 2012.

t




Fig 3. Radar plot of climatic profile's resource use, yield and average sale price. Each point shows the overall contribution of a variable to the vineyards outcomes for that climate. The left reflects vineyards in different climatic temperatures. The right reflects vineyards in different rainfall climates.




Fig 4. Scatter plot of vineyard yield against the average sale price as ratios to area harvested. The axes are in standard deviations with points coloured by climate.

and area was not simply about efficiently producing the most grapes. It is possible that
244
the relationship of scope one emissions between yield and sale price was closely tied to a
245
vineyard's area due to requiring more fuel to address more issues over greater distances.
246
It is difficult to discern the connection of scope one emissions directly, as fuel can be
247
used for a broad category of activities.
248

There are important considerations unique to winegrowing compared to other
249
agricultural industries. The vertical integration of winegrowing within the wine industry
250
ties winegrowers to secondary and tertiary industries, such as wine production,
251
packaging, transport and sales. This results in unique issues and considerations for each
252
vineyard, where on-the-ground decisions are influenced by other wine industry's choices,
253
such as the use of sustainable practices in vineyards as a requirement for sale in overseas
254
markets; notably these interactions can be further complicated by some winegrowers
255
being completely integrated into a wine company, while others are not [29].
256
Incorporating decisions into the model could help describe the contributing factors to
257
regional differences beyond resource consumption, motivating the call for more granular
258
data and more sophisticated modelling.
259

There are many on-the-ground decisions that influence both sales price and yield.
260
The decision to prioritise average sale price over quantity, is governed by complex
261
physical and social forces, for example international market demands, disease pressures
262
and natural disasters [24, 42–47], with many of these occurrences being highlighted
263
throughout the reports from Wine Australia over the past decade [19, 20, 48–54].
264
However, the changes in the coefficients (see Figure 2) are not reflective of many known
265
occurrences, such as the 2020 bush fires, which had higher values for coefficients than
266
prior years; During the 2020 bush fires 40,000 tonnes of grapes were lost across 18
267
different wine regions due to bush fires and smoke taint. In comparison to countrywide
268
pressures such as drought, this damage made up only 3% of the total amount of grapes
269
for that year; although acknowledged as a considerable loss on an individual basis, it
270
was deemed to be only a minor national concern by Wine Australia when compared to
271
other environmental pressures such as drought [55]
272

Climatic pressures are an important consideration for growers, especially those in
273
warmer and drier regions. The Wine Australia reports also show that warm inland
274
regions have seen a decline in profit over the past decade, whereas regions with lower
275
average sales prices did not [19, 20, 48, 50–55]. Vineyards in warm inland regions also
276
tend to be larger, making up for lower sale prices with larger yields. Considering the
277
negative correlation of average price to area, for this strategy to work, economies of
278
scale become an important factor. Given the large quantities of grapes that can be
279
produced by some vineyards, even at low margins there is the potential to be profitable.
280
However, the increasing climatic pressures mixed with the requirement for larger
281
volumes of water, make the sustainability of some vineyards come into question.
282
Furthermore, intensive farming in general is known to jeopardise the sustainability of an
283
operation through the degradation of soil and waterways [56–58]. There are established
284
methods that can help to mitigate these effects, such as the use of cover crops, midrow
285
crop rotation and efficient irrigation.
286

Some regions appeared to produce grapes of lower average sale price at scale whilst
287
others achieved higher priced grapes in lower volumes. This empirical finding is
288
consistent with Wine Australia's annual reports, which shows that some GI regions are
289
known for producing large amounts of lower grade (low value per tonne) grapes [49, 54].
290
Comparatively, other regions only produce grapes of higher sales price but in smaller
291
quantities. The difference in pricing per tonne between the lowest and highest regional
292
average sales prices was almost a hundred times, showing that region had a profound
293
influence. Some regions also had a mixture of high and low average sales price and yield
294
showing regional variability in pricing which may be explained by varieties produced. A
295

further possibility is the existence of regional upper limits on potential sales price, or
296
that there are diminishing returns in some regions when pursuing higher sales prices or
297
quantity; however these types of relationships may be obfuscated by knowledgeable
298
winegrowers who avoid such pitfalls.
299

Due to regional differences, different strategies are employed, such as some regions
300
targeting mass production over higher sales price. This is most notable when grouping
301
regions by climate, especially when considering GI Regions in the 'Hot Very Dry'
302
climate (see Figure 4). Figure 3 also shows that comparatively 'Warm' and 'Dry'
303
Regions manage their resources incredibly efficiently having generally larger areas using
304
similar resources to those in 'Cool' regions but having areas comparable to regions in
305
'Hot' climates. The coefficients for Model 4 also show a greater benefit of resource use
306
per area when producing grapes at higher average sale prices, showing higher resource
307
use per area reflective of higher average sales price (see Table 2). Although not chosen
308
over GI region, climate was considered to be a large determinant of the ability to
309
produce larger quantities of grapes, as well as a determinant in grape sale price [37].
310
The more granular GI Region likely explained a broader mix of geographical
311
phenomenon, such as soil, geology and access to water resources [27, 59]. The interaction
312
between year and GI Region likely accounted for events such as bushfires, which would
313
be impactful, but only at a local level, both in time and space.
314

We identified two main limitations to our linear modelling. First, model 1 and 2
315
over-predicting yield may have been due to preventative measures brought on by
316
regional pressures such as fire, frost and disease. More fuel and water was likely used to
317
prevent these issues from spreading within a region, thus disproportionately affecting
318
some vineyards compared to others locally. This type of maintenance is not well
319
captured in the models, especially when considering that some regions, especially those
320
in warmer areas, are not as prone to disease as cooler climates and could potentially
321
have lower fuel and water use per hectare. This could create a discrepancy in vineyards
322
that utilised preventative measures in wetter regions, as opposed to those that did not,
323
thus expending less fuel and energy but risking disease. When reviewing the differences
324
between regions, it is important to consider that vineyards in 'Hot Very Dry' areas can
325
be hundreds of times the size of those in other regions. This limitation could be
326
overcome by incorporating the profitability of vineyards, comparing the financial success
327
of working at different operational scales.
328

The second limitations was the lack of further explanatory variables to help link
329
models to causal affects. Variables such as the utilisation of renewable energy,
330
contractors, and the occurrence of disease, fire and frost were originally explored to
331
capture the discrepancies between similar vineyards that produced different yields and
332
crop values. However, none of these variables was significantly correlated with the
333
response variables, and did not add to model accuracy, even when considered as
334
interactions. Allowance for nonlinear relationships, specifically through splines, resulted
335
in more normally distributed residuals but at a drastically reduced overall accuracy
336
when comparing R^2 and Residual Square Error. Attempts to fully explain small
337
variations was always overshadowed by the dramatic differences in regional trends.
338
Having more data for each region would also be beneficial, allowing greater comparison
339
between regions.
340

The use of other models such as random forests and decision trees alongside more
341
variables and data may help to uncover the reasons for under or overestimation. These
342
differences could be caused by the use of alternative sustainable practices in the field.
343
Moreover, while there is evidence to suggest that environmentally sustainable practices
344
can reduce costs, and increase efficiency whilst improving the quality of grapes; more
345
research is needed to link these benefits across different regions and climates [10, 12, 13].
346

Conclusion

This study delved into the relationships between resource use, grape sales price and yield. The findings underscore the multifaceted nature of vineyard management, where the interplay of size, resource allocation, climate, and regional influences collectively shape both the expected sale price and the quantity of grape yields. Average sales price of grapes was not solely tied to the overall expenditure of resources, but rather to the efficient allocation of resources per area. This emphasises that factors beyond sheer scale contribute significantly to the final sale price of grapes produced. Moreover, regional and yearly variations exhibited substantial effects on vineyard outputs, impacting both sales price and quantity. The connection observed between smaller vineyards and higher grape sale price suggests that the management of smaller properties might be more efficient and profitable.

References

1. Beaumelle L, Giffard B, Tolle P, Winter S, Entling MH, Benítez E, et al. Biodiversity Conservation, Ecosystem Services and Organic Viticulture: A Glass Half-Full. *Agriculture, Ecosystems and Environment*. 2023;351. doi:10.1016/j.agee.2023.108474.
2. Barriguinha A, de Castro Neto M, Gil A. Vineyard Yield Estimation, Prediction, and Forecasting: A Systematic Literature Review. *Agronomy*. 2021;11(9). doi:10.3390/agronomy11091789.
3. He L, Fang W, Zhao G, Wu Z, Fu L, Li R, et al. Fruit Yield Prediction and Estimation in Orchards: A State-of-the-Art Comprehensive Review for Both Direct and Indirect Methods. *Computers and Electronics in Agriculture*. 2022;195:106812. doi:10.1016/j.compag.2022.106812.
4. Laurent C, Oger B, Taylor JA, Scholasch T, Metay A, Tisseyre B. A Review of the Issues, Methods and Perspectives for Yield Estimation, Prediction and Forecasting in Viticulture. *European Journal of Agronomy*. 2021;130:126339. doi:10.1016/j.eja.2021.126339.
5. Abbott T, Longbottom M, Wilkes E, Johnson D. AWRI: Assessing the Environmental Credentials of Australian Wine. *Wine & viticulture journal*. 2016;31(1):35–37.
6. Beauchet S, Rouault A, Thiollet-Scholtus M, Renouf M, Jourjon F, Renaud-Gentie C. Inter-Annual Variability in the Environmental Performance of Viticulture Technical Management Routes—a Case Study in the Middle Loire Valley (France). *The International Journal of Life Cycle Assessment*. 2019;24(2):253–265. doi:10.1007/s11367-018-1516-y.
7. Ferrara C, De Feo G. Life Cycle Assessment Application to the Wine Sector: A Critical Review. *Sustainability*. 2018;10(2). doi:10.3390/su10020395.
8. Finco A, Bentivoglio D, Chiaraluce G, Alberi M, Chiarelli E, Maino A, et al. Combining Precision Viticulture Technologies and Economic Indices to Sustainable Water Use Management. *Water*. 2022;14(9). doi:10.3390/w14091493.
9. Singh AP, Yerudkar A, Mariani V, Iannelli L, Glielmo L. A Bibliometric Review of the Use of Unmanned Aerial Vehicles in Precision Agriculture and Precision Viticulture for Sensing Applications. *Remote Sensing*. 2022;14(7). doi:10.3390/rs14071604.

10. Baiano A. An Overview on Sustainability in the Wine Production Chain. *Beverages*. 2021;7(1). doi:10.3390/beverages7010015.
11. Santiago-Brown I, Metcalfe A, Jerram C, Collins C. Sustainability Assessment in Wine-Grape Growing in the New World: Economic, Environmental, and Social Indicators for Agricultural Businesses. *Sustainability*. 2015;7(7):8178–8204. doi:10.3390/su7078178.
12. Montalvo-Falcón JV, Sánchez-García E, Marco-Lajara B, Martínez-Falcó J. Sustainability Research in the Wine Industry: A Bibliometric Approach. *Agronomy*. 2023;13(3). doi:10.3390/agronomy13030871.
13. Mariani A, Vastola A. Sustainable Winegrowing: Current Perspectives. *International Journal of Wine Research*. 2015;7(1):37–48.
14. Hemming S, de Zwart F, Elings A, Petropoulou A, Righini I. Cherry Tomato Production in Intelligent Greenhouses-Sensors and Ai for Control of Climate, Irrigation, Crop Yield, and Quality. *Sensors* (Basel, Switzerland). 2020;20(22):1–30. doi:10.3390/s20226430.
15. Kawasaki K, Uchida S. Quality Matters More Than Quantity: Asymmetric Temperature Effects on Crop Yield and Quality Grade. *American journal of agricultural economics*. 2016;98(4):1195–1209. doi:10.1093/ajae/aaw036.
16. ZHU Dw, ZHANG Hc, GUO Bw, XU K, DAI Qg, WEI Hy, et al. Effects of Nitrogen Level on Yield and Quality of Japonica Soft Super Rice. *Journal of Integrative Agriculture*. 2017;16(5):1018–1027. doi:10.1016/S2095-3119(16)61577-0.
17. Black CA, Parker M, Siebert TE, Capone DL, Francis IL. Terpenoids and Their Role in Wine Flavour: Recent Advances. *Australian Journal of Grape and Wine Research*. 2015;21(S1):582–600. doi:10.1111/ajgw.12186.
18. Schreier P, Jennings WG. Flavor Composition of Wines: A Review. *C R C Critical Reviews in Food Science and Nutrition*. 1979;12(1):59–111. doi:10.1080/10408397909527273.
19. Wine Australia. National Vintage Report 2021. 2021;.
20. Winemakers' Federation of Australia. National Vintage Report 2018. 2018;.
21. Yegge JM. Influence of Sensory and Non-Sensory Attributes of Chardonnay Wine on Acceptance and Purchase Intent. *ProQuest Dissertations Publishing*; 2001.
22. Canadell JG, Meyer CPM, Cook GD, Dowdy A, Briggs PR, Knauer J, et al. Multi-Decadal Increase of Forest Burned Area in Australia Is Linked to Climate Change. *Nature Communications*. 2021;12(1):6921. doi:10.1038/s41467-021-27225-4.
23. Longbottom ML, Petrie PR. Role of Vineyard Practices in Generating and Mitigating Greenhouse Gas Emissions. *Australian Journal of Grape and Wine Research*. 2015;21(S1):522–536. doi:10.1111/ajgw.12197.
24. Oliver DP, Bramley RGV, Riches D, Porter I, Edwards J. Review: Soil Physical and Chemical Properties as Indicators of Soil Quality in Australian Viticulture. *Australian Journal of Grape and Wine Research*. 2013;19(2):129–139. doi:10.1111/ajgw.12016.

25. Laurent C, Le Moguédec G, Taylor J, Scholasch T, Tisseyre B, Metay A. Local Influence of Climate on Grapevine: An Analytical Process Involving a Functional and Bayesian Exploration of Farm Data Time Series Synchronised with an eGDD Thermal Index. *OENO one*. 2022;56(2):301–317. doi:10.20870/oeno-one.2022.56.2.5443.
26. Liakos KG, Busato P, Moshou D, Pearson S, Bochtis D. Machine Learning in Agriculture: A Review. *Sensors*. 2018;18(8). doi:10.3390/s18082674.
27. Abbal P, Sablayrolles JM, Matzner-Lober É, Boursiquot JM, Baudrit C, Carboneau A. Decision Support System for Vine Growers Based on a Bayesian Network. *Journal of agricultural, biological, and environmental statistics*. 2016;21(1):131–151. doi:10.1007/s13253-015-0233-2.
28. Keith Jones. Australian Wine Industry Environment Strategy; 2002.
29. Knight H, Megicks P, Agarwal S, Leenders MAAM. Firm Resources and the Development of Environmental Sustainability among Small and Medium-Sized Enterprises: Evidence from the Australian Wine Industry. *Business Strategy and the Environment*. 2019;28(1):25–39. doi:10.1002/bse.2178.
30. SWA SWA. Sustainable Wingrowing Australia; 2022. <https://sustainablewingrowing.com.au/case-studies/>.
31. Attorney-General's Department. Wine Australia Corporation Act 1980; 2010.
32. G van Rossum. Python Tutorial, Technical Report CS-R9526; 1995. Centrum voor Wiskunde en Informatica (CWI),.
33. AGDEE AGDotEaE. National Greenhouse Accounts Factors: 2021; 2021.
34. Department of Climate Change, Energy, the Environment and Water. Australian National Greenhouse Accounts Factors; 2022.
35. Halliday JCJC. Australian Wine Encyclopedia. VIC: Hardie Grant Books; 2009.
36. SOAR CJ, SADRAS VO, PETRIE PR. Climate Drivers of Red Wine Quality in Four Contrasting Australian Wine Regions. *Australian journal of grape and wine research*. 2008;14(2):78–90. doi:10.1111/j.1755-0238.2008.00011.x.
37. Agosta E, Canziani P, Cavagnaro M. Regional Climate Variability Impacts on the Annual Grape Yield in Mendoza, Argentina. *Journal of Applied Meteorology and Climatology*. 2012;51(6):993–1009.
38. Fraga H, Costa R, Santos JA. Multivariate Clustering of Viticultural Terroirs in the Douro Winemaking Region. *Ciência Téc Vitiv.* 2017;32(2):142–153.
39. Sustainable Winegrowing Australia SWA. Sustainable Winegrowing Australia User Manual; 2021.
40. R Core Team. R: A Language and Environment for Statistical Computing; 2021. R Foundation for Statistical Computing.
41. Kuhn M. Building Predictive Models in R Using the Caret Package. *Journal of Statistical Software, Articles*. 2008;28(5):1–26. doi:10.18637/jss.v028.i05.
42. Abad J, Hermoso de Mendoza I, Marín D, Orcaray L, Santesteban LG. Cover Crops in Viticulture. A Systematic Review (1):
Implications on Soil Characteristics and Biodiversity in Vineyard. *OENO One*. 2021;55(1):295–312. doi:10.20870/oeno-one.2021.55.1.3599.

43. Cortez P, Teixeira J, Cerdeira A, Almeida F, Matos T, Reis J. Using Data Mining for Wine Quality Assessment. In: Discovery Science: 12th International Conference, DS 2009, Porto, Portugal, October 3-5, 2009 12. Springer; 2009. p. 66–79.
44. Hall A, Lamb DW, Holzapfel BP, Louis JP. Within-Season Temporal Variation in Correlations between Vineyard Canopy and Winegrape Composition and Yield. *Precision Agriculture*. 2011;12:103–117.
45. I Goodwin,, L McClymont,, D Lanyon, A Zerihun, J Hornbuckle, M Gibberd, et al.. Managing Soil and Water to Target Quality and Reduce Environmental Impact.; 2009.
46. Kasimati A, Espejo-García B, Darra N, Fountas S. Predicting Grape Sugar Content under Quality Attributes Using Normalized Difference Vegetation Index Data and Automated Machine Learning. *Sensors*. 2022;22(9). doi:10.3390/s22093249.
47. Srivastava S, Sadistap S. Non-Destructive Sensing Methods for Quality Assessment of on-Tree Fruits: A Review. *Journal of Food Measurement and Characterization*. 2018;12(1):497–526.
48. Wine Australia. National Vintage Report 2019. 2019;;
49. Wine Australia. National Vintage Report 2022. 2022;;
50. Winemakers' Federation of Australia. National Vintage Report 2013. 2013;;
51. Winemakers' Federation of Australia. National Vintage Report 2014. 2014;;
52. Winemakers' Federation of Australia. National Vintage Report 2015. 2015;;
53. Winemakers' Federation of Australia. National Vintage Report 2016. 2016;;
54. Winemakers' Federation of Australia. National Vintage Report 2017. 2017;;
55. Wine Australia. National Vintage Report 2020. 2020;;
56. Capello G, Biddoccu M, Ferraris S, Cavallo E. Effects of Tractor Passes on Hydrological and Soil Erosion Processes in Tilled and Grassed Vineyards. *Water*. 2019;11(10). doi:10.3390/w11102118.
57. Lin H. Hydropedology : Synergistic Integration of Soil Science and Hydrology. San Diego, NETHERLANDS, THE: Elsevier Science & Technology; 2012.
58. Pisciotta A, Cusimano G, Favara R. Groundwater Nitrate Risk Assessment Using Intrinsic Vulnerability Methods: A Comparative Study of Environmental Impact by Intensive Farming in the Mediterranean Region of Sicily, Italy. *Journal of geochemical exploration*. 2015;156:89–100. doi:10.1016/j.gexplo.2015.05.002.
59. Carmona G, Varela-Ortega C, Bromley J. The Use of Participatory Object-Oriented Bayesian Networks and Agro-Economic Models for Groundwater Management in Spain. *Water resources management*. 2011;25(5):1509–1524. doi:10.1007/s11269-010-9757-y.

Appendix

Table 7. P-values for the non-transformed water used variable's Pearson correlation coefficients.

Variable	Water Used
Yield	7.538E-01
Area	6.981E-01
Scope One Emissions	8.883E-01
$\frac{\text{Yield}}{\text{Area}}$	6.836E-01
Average Price Per Tonne	5.600E-02
$\frac{\text{Average Price per tonne}}{\text{Area}}$	1.522E-01




Fig 5. QQ-plot of Model 1.

Model 1 – Residuals vs fitted




Fig 6. Residuals vs fitted values for Model 1.

Model 2 – Normal QQ




Fig 7. QQ-plot of Model 2.

Model 2 – Residuals vs fitted




Fig 8. Residuals vs fitted values for Model 2.




Fig 9. QQ-plot of Model 3.




Fig 10. Residuals vs fitted values for Model 3.

Model 4 – Normal QQ




Fig 11. QQ-plot of Model 4.

Model 4 – Residuals vs fitted




Fig 12. Residuals vs fitted values for Model 4.

Model 5 – Normal QQ




Fig 13. QQ-plot of Model 5.

Model 5 – Residuals vs fitted




Fig 14. Residuals vs fitted values for Model 5.