

## **L6. Bayesian Decision Analysis**

# Bayesian Regression



## Linear regression:

$$y_i = \alpha + \beta x_i + \epsilon$$

## Probabilistic reformulation:

$$y_i \sim N(\alpha + \beta x_i, \sigma^2)$$

$$\epsilon \sim N(0, \sigma^2)$$



$$\alpha \sim N(\mu_\alpha = 0, \sigma_\alpha = 20^2)$$

$$\beta \sim N(\mu_\beta = 0, \sigma_\beta = 20^2)$$

$$\sigma \sim U(0, 20)$$

Jupyter Demo Simulation

## Hierarchical Bayesian Regression

$$\begin{array}{ll} \text{Group 1} & (x_{1,1}, y_{1,1}), (x_{2,1}, y_{2,1}), \dots, (x_{i,1}, y_{i,1}), \dots, (x_{n_1,1}, y_{n_1,1}) \\ \text{Group 2} & (x_{1,2}, y_{1,2}), (x_{2,2}, y_{2,2}), \dots, (x_{i,2}, y_{i,2}), \dots, (x_{n_2,2}, y_{n_2,2}) \\ & \vdots \\ \text{Group } j & (x_{1,j}, y_{1,j}), (x_{2,j}, y_{2,j}), \dots, (x_{i,j}, y_{i,j}), \dots, (x_{n_j,j}, y_{n_j,j}) \\ & \vdots \\ \text{Group } J & (x_{1,J}, y_{1,J}), (x_{2,J}, y_{2,J}), \dots, (x_{i,J}, y_{i,J}), \dots, (x_{n_J,J}, y_{n_J,J}) \end{array}$$

# Hierarchical Bayesian Regression


**Linear regression:**

$$y_{i,j} = \alpha_j + \beta_j x_{i,j} + \epsilon$$

**Probabilistic reformulation:**

$$y_{i,j} \sim N(\alpha_j + \beta_j x_{i,j}, \sigma_j^2)$$

$$\epsilon \sim N(0, \sigma_j^2)$$



$$y_j = (y_{1,j}, y_{2,j}, \dots, y_{n_j,j})$$

$$x_j = (x_{1,j}, x_{2,j}, \dots, x_{n_j,j})$$


**Linear regression:**

$$y_{i,j} = \alpha + \beta x_{i,j} + \epsilon$$

**Probabilistic reformulation:**

$$y_{i,j} \sim N(\alpha + \beta x_{i,j}, \sigma^2)$$

$$\epsilon \sim N(0, \sigma^2)$$



$$y_j = (y_{1,j}, y_{2,j}, \dots, y_{n_j,j})$$

$$x_j = (x_{1,j}, x_{2,j}, \dots, x_{n_j,j})$$

# Hierarchical Bayesian Regression


**Linear regression:**

$$y_{i,j} = \alpha_j + \beta_j x_{i,j} + \epsilon$$

**Probabilistic reformulation:**

$$y_{i,j} \sim N(\alpha_j + \beta_j x_{i,j}, \sigma_j^2)$$

$$\epsilon \sim N(0, \sigma_j^2)$$



$$y_j = (y_{1,j}, y_{2,j}, \dots, y_{n_j,j})$$

$$x_j = (x_{1,j}, x_{2,j}, \dots, x_{n_j,j})$$

# Hierarchical Bayesian Regression


**Linear regression:**

$$y_{i,j} = \alpha_j + \beta_j x_{i,j} + \epsilon$$

**Probabilistic reformulation:**

$$y_{i,j} \sim N(\alpha_j + \beta_j x_{i,j}, \sigma_j^2)$$


$$\epsilon \sim N(0, \sigma_j^2)$$



$$y_j = (y_{1,j}, y_{2,j}, \dots, y_{n_j,j})$$

$$x_j = (x_{1,j}, x_{2,j}, \dots, x_{n_j,j})$$

Hierarchical



$$y_j = (y_{1,j}, y_{2,j}, \dots, y_{n_j,j})$$

$$x_j = (x_{1,j}, x_{2,j}, \dots, x_{n_j,j})$$

# Hierarchical Bayesian Regression


## Bayesian Regression

$$y_{i,j} = \alpha + \beta x_{i,j} + \epsilon$$

$$y_{i,j} \sim N(\alpha + \beta x_{i,j}, \sigma^2)$$

$$\epsilon \sim N(0, \sigma^2)$$

Fixed hyper parameters




## Hierarchical Bayesian Regression

$$y_{i,j} = \alpha_j + \beta_j x_{i,j} + \epsilon$$

$$y_{i,j} \sim N(\alpha_j + \beta_j x_{i,j}, \sigma_j^2)$$

$$\epsilon \sim N(0, \sigma_j^2)$$

Fixed hyper parameters





Jupyter Demo Simulation

<http://pymc-devs.github.io/pymc3/notebooks/GLM-hierarchical.html>

## Bayesian decision theory


- ✓ Model has been built
- ✓ The inferences are conducted
- ✓ Model has been checked

What else?


- Two ways of using *Bayesian inference in decision making*

### (1) Measure uncertainties regarding predictive quantiles

Bayesian inference



### (2) Within decision analysis (Multi-stages decision problems)



## Elements of Bayesian decision analysis

Decision analysis is inherently more complicated than statistical inference because it involves *optimization over decision* as well *averaging over uncertainties*

### Elements of Bayesian decision analysis

1. Enumerate the space of all possible decisions (actions)  $a$  and outcome  $o$ 
  - The vector of outcomes  $o$  can include observables (predicted values  $\tilde{y}$ ) and parameters  $\theta$
2. Determine  $p(o|a)$ , a *conditional posterior probability distribution* of outcomes  $o$  for each decision option  $a$ 
  - Outcome  $o$  is random variable, while decision  $a$  is deterministic (set by user)
3. Define a utility function  $U(o)$  mapping outcomes  $o$  onto the real numbers
  - If multiple attributes (vector)  $o$  is considered, the utility function must trade off different goods
4. Compute the  $E =$  expected utility  $E(U(o)|a)$  as a function of the decision  $a$ , and choose the decision with highest expected utility

## Examples that will discussed in the lecture

1. Survey incentives
  - ✓ Conduct only step 1 and 2 to estimate the expected effect depending on the option that can be taken
2. Medical test decision
  - ✓ Conduct only step 1 and 2 in a sequential decision making and computed value of information
3. Risk analysis of Radon exposure and making prevention strategies
  - ✓ Conduct the full Bayesian analysis

## Example 1: Survey incentives




- Do the benefits of incentives outweigh the cost?
- If an incentive is given,
  - ✓ how and when should it be offered
  - ✓ whom should it be offered to
  - ✓ what form should it take
  - ✓ how large should its value be?

## Example 1: Survey incentives

### Data descriptions:

The New York City Social Indicators Survey, a telephone study conducted every two years that has had a response rate below 50%



- In total 39 surveys including 101 experiment conditions are conducted:
- $y_i$  is the observed response rate for observation  $i = 1, \dots, 101$
- All experiments has different conditions  $x_i$  on:
  - The **value of the incentive** (in tens of 1999dollars)
  - The **timing of the incentive** payment (given before the survey or after)
  - The **form of the incentive** (cash or gift)
  - The **mode of the survey** (face-to-face or telephone)
  - The **burden or effort**, required to survey respondents (high burden or low burden)
- Use the differences,  $z_i = y_i - y_i^0$ ,  
where  $y_i^0$  corresponds to the lowest valued incentive condition

## Example 1: Survey incentives

### A simple linear regression approach

Response rate :  $y_i = X_i\beta + \beta_0$

$$y_i = \frac{n_i}{N_i} = \frac{\text{repondents}}{\text{Trial}}$$

Predictor variables :  $X_i = (X_{1,i}, X_{2,i}, X_{3,i}, X_{4,i}, X_{5,i})$

$X_{1,i}$  :The **value of the incentive** (in tens of 1999dollars)

$X_{2,i}$  :The **timing of the incentive** payment (given before the survey or after)

$X_{3,i}$  :The **form of the incentive** (cash or gift)

$X_{4,i}$  :The **mode of the survey** (face-to-face or telephone)


$X_{5,i}$  :The **burden or effort**, required to survey respondents (high burden or low burden)

**Limitation**s in employing a classical regression model relating  $y_i$  to the predictor variables:

- Unable to model interactions
- Unable to reflect the hierarchical structure of the data
- Difficulty in dealing with unequal sample size

## Example 1: Survey incentives

- **Step1** : perform a meta-analysis to estimate the effects of incentives on response rate, as a function of the amount of the incentive and the way it is implemented



$\alpha_{j(i)}$  : random effect for the survey  $j = 1, \dots, 39$

(address the hierarchical structure)

$X_i \beta$  : linear predictor for conditioned on data point  $i$

$\pi_i$  : population response probability  $i = 1, \dots, 101$


$N_i$  : number of persons contacted

$n_i$  : number of respondents ( $y_i = n_i/N_i$ )



## Example 1: Survey incentives


- Step1** : perform a meta-analysis to estimate the effects of incentives on response rate, as a function of the amount of the incentive and the way it is implemented



$$y_{1,1} = \frac{n_{1,1}}{N_{1,1}} \quad \text{Response rate}$$

## Example 1: Survey incentives

- **Step2** : use the inference to estimate the costs and benefits of incentives



- Prepaid
- After paid

$$y_i = X_i\hat{\beta} + \beta_0$$

### Some of the findings are :

- an extra \$10 in incentive is expected to increase the response rate by 3–4 percentage points
- cash incentives increase the response rate by about 1 percentage point relative to noncash
- prepaid incentives increase the response rate by 1–2 percentage points relative to postpaid
- incentives have a bigger impact (by about 5 percentage points) on high-burden surveys compared to low-burden surveys.


**These inferences can be used to design a new survey (cost vs. response rate)**

## Example 2: Multistage decision making for medical screening

95-year-old man with an apparently malignant tumor in the lung




<Multi-stages decision problems>



Bayesian inference is particularly useful in updating the state of knowledge with the information gained at each step.

## Example 2: Multistage decision making for medical screening

### Single decision point (without test)




Cost of Radiotherapy and Surgery = 1

Quality-adjusted life expectancy under each treatment:

- With no treatment :  $0.1 \times 34.8 + 0.9 \times 5.6 = 8.5 m$
- With radiotherapy :  $0.1 \times 34.8 + 0.9 \times 16.7 - 1 = 17.5 m$  ← Best option
- With surgery :  $0.35 \times 0 + 0.65 \times (0.1 \times 34.8 + 0.9 \times 20.3 - 1) = 13.5 m$

## Example 2: Multistage decision making for medical screening

### Sequential decision making




- Based on the test result, the probability of cancer can be updated using Bayes rule:

$$\Pr(C|T) = \frac{\Pr(C) p(T|C)}{\Pr(C) p(T|C) + \Pr(NC) p(T|NC)}$$

- Updated information can be used for sequential decision making

## Example 2: Multistage decision making for medical screening

### Update information based on the test results



$$\Pr(T = +) = \Pr(NC) p(T = +|NC) + \Pr(C) p(T = +|C) \\ = 0.1 \cdot 0.02 + 0.9 \cdot 0.7 = 0.632$$

$$\Pr(T = -) = \Pr(NC) p(T = -|NC) + \Pr(C) p(T = -|C) \\ = 0.1 \cdot 0.98 + 0.9 \cdot 0.3 = 0.368$$

$$\Pr(C|T) = \frac{\Pr(C) p(T|C)}{\Pr(T)}$$

$$\Pr(C|T = +) = \frac{0.9 \cdot 0.7}{0.1 \cdot 0.02 + 0.9 \cdot 0.7} = 0.997$$

$$\Pr(C|T = -) = \frac{0.9 \cdot 0.3}{0.1 \cdot 0.98 + 0.9 \cdot 0.3} = 0.734$$

$$\Pr(NC|T) = \frac{\Pr(NC) p(T|NC)}{\Pr(T)}$$

$$\Pr(NC|T = +) = \frac{0.1 \cdot 0.02}{0.1 \cdot 0.02 + 0.9 \cdot 0.7} = 0.003$$


$$\Pr(NC|T = -) = \frac{0.1 \cdot 0.98}{0.1 \cdot 0.98 + 0.9 \cdot 0.3} = 0.266$$

## Example 2: Multistage decision making for medical screening

Choose the best action at the second stage using the updated information


$$Pr(\text{cancer} | T = +) = \frac{0.9 \cdot 0.7}{0.9 \cdot 0.7 + 0.1 \cdot 0.02} = 0.997$$

Quality-adjusted life expectancy under each treatment:

- With no treatment :  $0.997 \cdot 5.6 + 0.003 \cdot 34.8 = 5.7$  months
- With radiotherapy :  $0.997 \cdot 16.7 + 0.003 \cdot 34.8 - 1 = 15.8$  months  Best action
- With surgery :  $0.35 \cdot 0.65(0.997 \cdot 20.3 + 0.003 \cdot 34.8 - 1) = 12.6$  months

$$Pr(\text{cancer} | T = -) = \frac{0.9 \cdot 0.3}{0.9 \cdot 0.3 + 0.1 \cdot 0.98} = 0.734$$


Quality-adjusted life expectancy under each treatment:

- With no treatment :  $0.734 \cdot 5.6 + 0.266 \cdot 34.8 = 13.4$  months
- With radiotherapy :  $0.734 \cdot 16.7 + 0.266 \cdot 34.8 - 1 = 20.5$  months  Best action
- With surgery :  $0.35 \cdot 0.65(0.734 \cdot 20.3 + 0.266 \cdot 34.8 - 1) = 15.1$  months

It is clear conducting a test is not a good idea since no change in the selected option!

## Example 2: Multistage decision making for medical screening

### Sequential decision making




When selecting a decision at the first state, we should assume that all the decisions at the subsequent stages will be made optimum!! (key idea of sequential decision making: Control, MDP)



## Example 2: Multistage decision making for medical screening

### Decision analysis for performing test






- No test : 17.5 months is the best
- Test :  $0.368 \cdot 19.5 + 0.632 \cdot 15.0 = 16.6$  months


No change in the selected option and no improvement in the expected cost

→ Not conducting the test is the best option

### Example 3: Hierarchical decision analysis for home radon



### Example 3: Hierarchical decision analysis for home radon



#### Two tasks:

- Given prior information about Radon concentration, decide whether to conduct a measurement test (first decision)
- Given the measurement of Radon concentration (we have a posterior of Radon con.), decide whether to remediate (second decision)

Performing the decision analysis requires estimating for the risks, which is done by using hierarchical Bayesian model

### Example 3: Hierarchical decision analysis for home radon

#### Define utility : trades off dollars and lives

- $D_d$ , the dollar value associated with a reduction of  $10^{-6}$  in probability of death for lung cancer (the money that you are willing to pay to reduce  $10^{-6}$  in probability of death for lung cancer)
- $D_r$ , the dollar value associated with a reduction of 1 pCi/L in home radon level for 30-year period
  - depends on the number of lives saved by a drop in the Radon level
  - affected by variety of factors including gender, smoking status...
  - $D_r = 4800D_d$  (typical U.S. household)
- $R_{action}$ , the home radon level above which you should remediate if your radon level is known
  - depends on the dollar value of radon reduction and the benefits of remediation

Action level  $R_{action}$  is determined as the value at which the benefit of remediation

$$\$D_r(R_{action} - R_{remed}) = \text{remediation cost of \$2,000}$$

$$R_{action} = \frac{\$2,000}{D_r} + R_{remedy}$$

$$\text{where } R_{remed} = \begin{cases} 2 & \text{if home radon level} > 2 \\ \text{home radon level} & \text{otherwise} \end{cases}$$

$R_{action} = 4$  pCi/L is used as exemplary value, which can vary depending on financial resources, general risk tolerance, attitude toward risk, smoking status, etc.

### Example 3: Hierarchical decision analysis for home radon

#### Bayesian inference for country radon level

Two data sets:

- Long-term measurements from approximately 5,000 houses, selected as a cluster sample from 125 randomly selected counties
- Short-term measurements from about 80,000 houses, sampled at random from all the counties in the U.S.


a relatively small amount of accurate data VS. a large amount of biased and imprecise data

#### Challenges:

Use the good data to calibrate the bad data, so that inference can be made about the entire counties, not merely the 125 counties in the sample of long-term measurements


### Example 3: Hierarchical decision analysis for home radon

#### Hierarchical modeling for individual house radon level



### Example 3: Hierarchical decision analysis for home radon

#### Hierarchical modeling for individual house radon level



$W_j$ : county-level predictors including climate data and a measure of the uranium level in the soil

$X_i$ : household-level predictors including indicators for whether the house has a basement and

**measurement type** ( $\sigma_i^2$  varies depending on measurement type (long-term or short term))

## Example 3: Hierarchical decision analysis for home radon

### Bayesian inference for individual home level

$R_i$  = radon concentration in house  $i$

$$\theta_i = \log(R_i) \sim N(M_i, S_i^2)$$

- The mean  $M_i = X_i \hat{B} + \hat{\alpha}_j$  is computed from the posterior simulations of the model estimation.  
 $\hat{B}$  and  $\hat{\alpha}$  are the posterior means from the analysis in the appropriate region of the country.  
(computed from county level measurement data)
- The variance  $S_i^2$  is computed from the posterior simulations of the model estimation taking into account the posterior uncertainty in the coefficients  $\alpha, \beta$  and also hierarchical variance components  $\eta^2$  and  $\tau^2$
- Serves as a **prior distribution** for the homeowner in that the distribution is constructed solely based on the basement information  $X_i$  and the county level parameter  $\hat{\alpha}_j$

$$\theta_i \sim N(X_i \hat{B} + \hat{\alpha}_j, S_i^2)$$

- Using  $p(\theta_i)$ , decision whether to perform measurement test or not can be made



## Example 3: Hierarchical decision analysis for home radon

### Bayesian inference for individual home level

Likelihood :

$$Y_i \sim N(\theta, \sigma_Y^2) \rightarrow p(y_i | \theta, \sigma_Y^2) = \frac{1}{\sqrt{2\pi\sigma_Y^2}} \exp\left(-\frac{(y_i - \theta)^2}{2\sigma_Y^2}\right)$$

Prior:

$$\theta \sim N(\mu_0, \tau_0^2) \rightarrow p(\theta) = \frac{1}{\sqrt{2\pi\tau_0^2}} \exp\left(-\frac{(\theta - \mu_0)^2}{2\tau_0^2}\right)$$

Posterior on  $\theta = \mu_Y$

$$P(\theta | y) = N\left(\theta \left| \frac{\frac{\mu_0}{\tau_0^2} + \frac{n\bar{y}}{\sigma_Y^2}}{\frac{1}{\tau_0^2} + \frac{n}{\sigma_Y^2}}, \left(\frac{1}{\tau_0^2} + \frac{n}{\sigma_Y^2}\right)^{-1} \right.\right)$$

$R_i$  = radon concentration in house  $i$

$\theta_i = \log(R_i)$

Assume we measure Radon concentration  $y_i \sim N(\theta_i, \sigma_i^2)$

Likelihood

$$y_i \sim N(\theta_i, \sigma_i^2)$$

Prior (from previous slide) :

$$p(\theta_i) = N(M_i, S_i^2)$$

The posterior distribution :

$$\theta_i | M_i, y_i \sim N(\Lambda_i, V_i) \quad \Lambda_i = \frac{\frac{M_i}{S_i^2} + \frac{y_i}{\sigma_i^2}}{\frac{1}{S_i^2} + \frac{1}{\sigma_i^2}} \quad V_i = \frac{1}{\frac{1}{S_i^2} + \frac{1}{\sigma_i^2}}$$

### Example 3: Hierarchical decision analysis for home radon

#### Decision analysis for individual homeowners

$$\begin{aligned} R_i &= \text{radon concentration in house } i \\ \theta_i &= \log(R_i) \end{aligned}$$


Prior :  $\theta_i = N(M_i, S_i^2)$  Predictive distribution based on hierarchical model

Decision whether to perform measurement test or not

Posterior :  $\theta_i | M_i, y_i \sim N(\Lambda_i, V_i)$

Decision whether to remediate or not

### Example 3: Hierarchical decision analysis for home radon



$$\text{Expected loss} = \text{cost} + D_r E(R)$$

### Example 3: Hierarchical decision analysis for home radon

#### Decision analysis for individual homeowners

$$\begin{aligned} R_i &= \text{radon concentration in house } i \\ \theta_i &= \log(R_i) \end{aligned}$$

$$R = e^\theta, \theta \sim N(M, S^2)$$

$$E[R] = E[e^\theta] = e^{M + \frac{1}{2}S^2}$$

$$E[R|\theta > a] = E[e^\theta | \theta > a] = e^{M + \frac{1}{2}S^2} \left( 1 - \Phi \left( \frac{M + S^2 - a}{S} \right) \right)$$

Do nothing

$$L_1 = D_r E(R) = D_r e^{M + \frac{1}{2}S^2}$$

Remediate without test

$$\begin{aligned} L_2 &= \$2000 + D_r E(\min(R, R_{\text{remed}})) \\ &= \$2000 + D_r [R_{\text{remed}} \Pr(R \geq R_{\text{remed}}) + E(R|R < R_{\text{remed}}) \Pr(R < R_{\text{remed}})] \\ &= \$2000 + D_r \left[ R_{\text{remed}} \Phi \left( \frac{M - \log(R_{\text{remed}})}{S} \right) + e^{M + \frac{1}{2}S^2} \left( 1 - \Phi \left( \frac{M + S^2 - \log(R_{\text{remed}})}{S} \right) \right) \right] \end{aligned}$$

## Example 3: Hierarchical decision analysis for home radon

### Decision analysis for individual homeowners

#### Remediate without test

$$\begin{aligned} L_2 &= \$2000 + D_r E(\min(R, R_{\text{remed}})) \\ &= \$2000 + D_r [R_{\text{remed}} \Pr(R \geq R_{\text{remed}}) + E(R|R < R_{\text{remed}}) \Pr(R < R_{\text{remed}})] \\ &= \$2000 + D_r \left[ R_{\text{remed}} \Phi\left(\frac{M - \log(R_{\text{remed}})}{S}\right) + e^{M + \frac{1}{2}S^2} \left(1 - \Phi\left(\frac{M + S^2 - \log(R_{\text{remed}})}{S}\right)\right) \right] \end{aligned}$$

Prior  
 $(M, S^2)$



$(\Lambda, V)$   
Posterior

#### Perform measurement test

- Do nothing

$$L_{3a} = \$50 + D_r \frac{1}{30} e^{M + \frac{1}{2}S^2} + D_r e^{\Lambda + \frac{1}{2}V}$$


Exposure of 1 year      29 years of exposure after test is done

- Remediate

$$\begin{aligned} L_{3b} &= \$50 + D_r \frac{1}{30} e^{M + \frac{1}{2}S^2} + \$2000 \\ &\quad + D_r \left[ R_{\text{remed}} \Phi\left(\frac{\Lambda - \log(R_{\text{remed}})}{\sqrt{V}}\right) + e^{\Lambda + \frac{1}{2}V} \left(1 - \Phi\left(\frac{\Lambda + V - \log(R_{\text{remed}})}{\sqrt{V}}\right)\right) \right] \end{aligned}$$

## Example 3: Hierarchical decision analysis for home radon

### Decision analysis for individual homeowners



Expected loss for performing test :  $L^3 = E(\min(L_{3a}, L_{3b}))$  :  
(optimum decision in the second state is imbedded)

1. Simulate 5000 draws of  $y \sim N(M, S^2 + \sigma^2)$
2. For each draw of  $y$ , compute  $\min(L_{3a}, L_{3b})$
3. Estimate  $L_3$  as the average of these 5000 values

(Considering uncertainty of  $y$ )

### Example 3: Hierarchical decision analysis for home radon

#### Decision analysis for individual homeowners

