

Ax-Grothendieck in lean

Joseph Hua

June 10, 2022

Imperial College London

Polynomial maps

Definition (Polynomial maps)

Polynomial maps on a field K are regular endomorphisms on K^n , i.e. n polynomials in $K[x_1, \dots, x_n]$.

Polynomial maps

Definition (Polynomial maps)

Polynomial maps on a field K are regular endomorphisms on K^n , i.e. n polynomials in $K[x_1, \dots, x_n]$.

Examples

- Surjective but not injective: $f : \mathbb{C} \rightarrow \mathbb{C} := x \mapsto x^2$

Polynomial maps

Definition (Polynomial maps)

Polynomial maps on a field K are regular endomorphisms on K^n , i.e. n polynomials in $K[x_1, \dots, x_n]$.

Examples

- Surjective but not injective: $f : \mathbb{C} \rightarrow \mathbb{C} := x \mapsto x^2$
- Neither surjective nor injective: $f : \mathbb{C}^2 \rightarrow \mathbb{C}^2 := (x, y) \mapsto (x, xy)$

Polynomial maps

Definition (Polynomial maps)

Polynomial maps on a field K are regular endomorphisms on K^n , i.e. n polynomials in $K[x_1, \dots, x_n]$.

Examples

- Surjective but not injective: $f : \mathbb{C} \rightarrow \mathbb{C} := x \mapsto x^2$
- Neither surjective nor injective: $f : \mathbb{C}^2 \rightarrow \mathbb{C}^2 := (x, y) \mapsto (x, xy)$
- Bijective: $f : \mathbb{C}^3 \rightarrow \mathbb{C}^3 := (x, y, z) \mapsto (x, y, z + xy)$

Polynomial maps

Definition (Polynomial maps)

Polynomial maps on a field K are regular endomorphisms on K^n , i.e. n polynomials in $K[x_1, \dots, x_n]$.

Examples

- Surjective but not injective: $f : \mathbb{C} \rightarrow \mathbb{C} := x \mapsto x^2$
- Neither surjective nor injective: $f : \mathbb{C}^2 \rightarrow \mathbb{C}^2 := (x, y) \mapsto (x, xy)$
- Bijective: $f : \mathbb{C}^3 \rightarrow \mathbb{C}^3 := (x, y, z) \mapsto (x, y, z + xy)$

Code

```
def poly_map (K : Type*) [comm_semiring K] (n : ℕ) : Type* :=  
  fin n → mv_polynomial (fin n) K  
  
def eval : poly_map K n → (fin n → K) → (fin n → K) :=  
  λ ps as k, mv_polynomial.eval as (ps k)
```

Theorem (Ax-Grothendieck)

Any injective polynomial map over an algebraically closed field is surjective. In particular injective polynomial maps over \mathbb{C} are surjective.

Theorem (Ax-Grothendieck)

Any injective polynomial map over an algebraically closed field is surjective. In particular injective polynomial maps over \mathbb{C} are surjective.

```
theorem Ax_Groth {n : ℕ} {ps : poly_map K n}
  (hinject : function.injective (poly_map.eval ps)) :
  function.surjective (poly_map.eval ps) := sorry
```


Theorem (Ax-Grothendieck)

Any injective polynomial map over an algebraically closed field is surjective. In particular injective polynomial maps over \mathbb{C} are surjective.

```
theorem Ax_Groth {n : ℕ} {ps : poly_map K n}
  (hinject : function.injective (poly_map.eval ps)) :
  function.surjective (poly_map.eval ps) := sorry
```

- Amazing fact: the Lefschetz principle implies that we only need to show this for a single algebraically closed field of characteristic n (ACF_n) and it will be true for all of ACF_n (n is zero or prime).

Theorem (Ax-Grothendieck)

Any injective polynomial map over an algebraically closed field is surjective. In particular injective polynomial maps over \mathbb{C} are surjective.

```
theorem Ax_Groth {n : ℕ} {ps : poly_map K n}
  (hinject : function.injective (poly_map.eval ps)) :
  function.surjective (poly_map.eval ps) := sorry
```

- Amazing fact: the Lefschetz principle implies that we only need to show this for a single algebraically closed field of characteristic n (ACF_n) and it will be true for all of ACF_n (n is zero or prime).
- Amazing fact: Lefschetz also says that if we show it for all ACF_p for large p then it is also true for ACF_0 .

Theorem (Ax-Grothendieck)

Any injective polynomial map over an algebraically closed field is surjective. In particular injective polynomial maps over \mathbb{C} are surjective.

```
theorem Ax_Groth {n : ℕ} {ps : poly_map K n}
  (hinject : function.injective (poly_map.eval ps)) :
  function.surjective (poly_map.eval ps) := sorry
```

- Amazing fact: the Lefschetz principle implies that we only need to show this for a single algebraically closed field of characteristic n (ACF_n) and it will be true for all of ACF_n (n is zero or prime).
- Amazing fact: Lefschetz also says that if we show it for all ACF_p for large p then it is also true for ACF_0 .
- Good news: We can easily show it for algebraic closures of \mathbb{F}_p for any p .

Locally finite fields

Definition (Locally finite fields)

Let K be a field of characteristic p a prime. Then the following are equivalent definitions for K being a *locally finite field*:

1. The minimal subfield generated by any finite subset of K is finite.
2. $\mathbb{F}_p \rightarrow K$ is algebraic.
3. K embeds into an algebraic closure of \mathbb{F}_p .

Theorem

Locally finite fields satisfy Ax-Grothendieck.

Proof.

$$\begin{array}{c} \mathbb{F}_p \hookrightarrow \mathbb{F}_p(\text{coeffs}) \hookrightarrow K \\ \downarrow \mathbb{F}_p(\text{coeffs}) \text{ respects injectivity} \end{array}$$



The Lefschetz principle

Theorem (Lefschetz principle)

Let ϕ be a sentence in the language of rings. Then the following are equivalent:

- 1. Some model of ACF_0 satisfies ϕ . (If you like $\mathbb{C} \models \phi$.)*
- 2. $\text{ACF}_0 \models \phi$*
- 3. There exists $n \in \mathbb{N}$ such that for any prime p greater than n , $\text{ACF}_p \models \phi$*
- 4. There exists $n \in \mathbb{N}$ such that for any prime p greater than n , some model of ACF_p satisfies ϕ .*

(1. \leftrightarrow 2.) and (3. \leftrightarrow 4.) are due to the theories ACF_p being complete for any p (0 or prime); (2. \leftrightarrow 3.) due to compactness theorem.

Proof overview

An overview of the proof:

