

Normality inducing transformations

Adding missing feature as a normality inducing transformation

The Linear Regression model is

$$\mathbf{y} = \Theta^T \mathbf{x} + \epsilon$$

As explained before, Regression produces a conditional probability

$$p(\hat{\mathbf{y}}|\mathbf{x})$$

where $\hat{\mathbf{y}}$ and ϵ are *Normally distributed variables*.

Assumptions of the Linear Regression model are violated if

- ϵ is not Normal
- the individual $\epsilon^{(i)}$ display a pattern
- the individual $\epsilon^{(i)}$ have different variances (heteroscedastic)

One reason for failure of these assumptions is a missing feature

- "curvy" data set and Linear model
 - we saw pattern of errors: larger in tails
 - variances increased in tail


Adding a feature (e.g., second order polynomial term for the curvy data set) can be seen as a normality inducing transformation.

Log transformation

We've seen this in our lecture on Logistic Regression

- the probabilities are *not* normally distributed
- the odds are *not* normally distributed
- the *log odds* **is** normally distributed

```
In [4]: tf = tmh.TransformHelper()  
tf.plot_odds()
```



$$\begin{aligned}
 \frac{\hat{p}}{1-\hat{p}} &= \frac{\frac{1}{1+e^{-s}}}{1-\frac{1}{1+e^{-s}}} \\
 &= \frac{\frac{1}{1+e^{-s}}}{\frac{1+e^{-s}-1}{1+e^{-s}}} \\
 &= \frac{1}{e^{-s}} \\
 &= e^s
 \end{aligned}$$

So LogisticRegression is really just a LinearRegression with a transformed target

$$\log\left(\frac{\hat{p}}{1-\hat{p}}\right) = \Theta^T \cdot x$$

Other transformations

Centering

Transforming a feature to have mean 0.

$$\mathbf{x}_j^{(i)} = \mathbf{x}_j^{(i)} - \bar{\mathbf{x}}_j$$

- low values now become negative
 - more clearly indicates deleterious effect than a low, positive number
 - example: Star Ratings for movies
- some algorithms (PCA) need centered data

Bucketing/Binning

- Target may be linear in a feature only in broad ranges of the feature
 - income vs age
 - very young (below working age) all income is identical (0)
 - very old (above retirement) - no job related income
 - Latitude/Longitude
 - small changes matter MUCH less than big changes
- Converts numerical feature
 - into categorical **Is bucket 1, Is bucket 2, ...**
 - ordinal: replace value with center value of bin

Bucket size choices:

- Equal spaced buckets
- Equal quantile buckets

Lesson Don't fit a square peg (non-linear response) into a round hole (linear model)

Outliers

Pull in extreme values to reduce their influence on the fit.

- Clipping, Winsorization

In [9]: `print("Done")`

Done

