

# ***STATISTICAL INFERENCES (2cr)***

## Chapter 9 One- and Two-Sample Estimation Problems

Zhong Guan

Math, IUSB

# Outline

- 1 9.10 Estimating a Proportion
  - Estimating Single Proportion
  - Sample Size for Estimating  $p$
  
- 2 9.11 Difference of Two Proportions

# Outline

- 1 9.10 Estimating a Proportion
  - Estimating Single Proportion
  - Sample Size for Estimating  $p$
  
- 2 9.11 Difference of Two Proportions

# Outline

- 1 9.10 Estimating a Proportion
  - Estimating Single Proportion
  - Sample Size for Estimating  $p$
  
- 2 9.11 Difference of Two Proportions

# Do you know how to read the polls' results?

- If the the opinion poll says candidate A leads B 3%: 45% vs 42%, what does it mean?
- The poll gives sample proportions not the population proportions.
- You need to find the margin of error.

# Do you know how to read the polls' results?

- If the the opinion poll says candidate A leads B 3%: 45% vs 42%, what does it mean?
- The poll gives sample proportions not the population proportions.
- You need to find the margin of error.

# Do you know how to read the polls' results?

- If the the opinion poll says candidate A leads B 3%: 45% vs 42%, what does it mean?
- The poll gives sample proportions not the population proportions.
- You need to find the margin of error.

# Do you know how to read the polls' results?

- If the the opinion poll says candidate A leads B 3%: 45% vs 42%, what does it mean?
- The poll gives sample proportions not the population proportions.
- You need to find the margin of error.



# How does the poll work?

- The poll collects the opinions of  $n$  voters.
- $Y$  of the  $n$  voters will vote for A.
- The sample proportion is  $\hat{p} = Y/n$  which is a point estimate of the population proportion  $p$ .
- $Y$  has a binomial distribution  $b(n, p)$ .
- How to construct a confidence interval for  $p$ ?

# How does the poll work?

- The poll collects the opinions of  $n$  voters.
- $Y$  of the  $n$  voters will vote for A.
- The sample proportion is  $\hat{p} = Y/n$  which is a point estimate of the population proportion  $p$ .
- $Y$  has a binomial distribution  $b(n, p)$ .
- How to construct a confidence interval for  $p$ ?

# How does the poll work?

- The poll collects the opinions of  $n$  voters.
- $Y$  of the  $n$  voters will vote for A.
- The sample proportion is  $\hat{p} = Y/n$  which is a point estimate of the population proportion  $p$ .
- $Y$  has a binomial distribution  $b(n, p)$ .
- How to construct a confidence interval for  $p$ ?

# How does the poll work?

- The poll collects the opinions of  $n$  voters.
- $Y$  of the  $n$  voters will vote for A.
- The sample proportion is  $\hat{p} = Y/n$  which is a point estimate of the population proportion  $p$ .
- $Y$  has a binomial distribution  $b(n, p)$ .
- How to construct a confidence interval for  $p$ ?

# How does the poll work?

- The poll collects the opinions of  $n$  voters.
- $Y$  of the  $n$  voters will vote for A.
- The sample proportion is  $\hat{p} = Y/n$  which is a point estimate of the population proportion  $p$ .
- $Y$  has a binomial distribution  $b(n, p)$ .
- How to construct a confidence interval for  $p$ ?

# Normal Approximation of Sampling Distribution of $\hat{p}$ :

- By the C.L.T., if  $n$  is large, then

$$P\left[-z_{\alpha/2} \leq \frac{\bar{X}-p}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2}\right] \approx 1 - \alpha.$$

- Since  $p \approx \hat{p}$ ,

$$P\left[-z_{\alpha/2} \leq \frac{\bar{X}-p}{\sqrt{\hat{p}(1-\hat{p})/n}} \leq z_{\alpha/2}\right] \approx 1 - \alpha.$$

# Normal Approximation of Sampling Distribution of $\hat{p}$ :

- By the C.L.T., if  $n$  is large, then

$$P\left[-z_{\alpha/2} \leq \frac{\bar{X}-p}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2}\right] \approx 1 - \alpha.$$

- Since  $p \approx \hat{p}$ ,

$$P\left[-z_{\alpha/2} \leq \frac{\bar{X}-p}{\sqrt{\hat{p}(1-\hat{p})/n}} \leq z_{\alpha/2}\right] \approx 1 - \alpha.$$

# Confidence Interval for Population Proportion $p$ :

- If sample size  $n$  is large, i.e.  $n\hat{p} > 5$  and  $n\hat{q} = n(1 - \hat{p}) > 5$ ,
- the  $100(1 - \alpha)\%$  confidence interval for the population proportion  $p$  is

$$\bar{X} \pm E, \quad E = z_{\alpha/2} \sqrt{\hat{p}(1 - \hat{p})/n}.$$

- $E$  is called the margin of error, and  $z_{\alpha/2}$  satisfies



# Confidence Interval for Population Proportion $p$ :

- If sample size  $n$  is large, i.e.  $n\hat{p} > 5$  and  $n\hat{q} = n(1 - \hat{p}) > 5$ ,
- the  $100(1 - \alpha)\%$  confidence interval for the population proportion  $p$  is

$$\bar{X} \pm E, \quad E = z_{\alpha/2} \sqrt{\hat{p}(1 - \hat{p})/n}.$$

- $E$  is called the margin of error, and  $z_{\alpha/2}$  satisfies

# Confidence Interval for Population Proportion $p$ :

- If sample size  $n$  is large, i.e.  $n\hat{p} > 5$  and  $n\hat{q} = n(1 - \hat{p}) > 5$ ,
- the  $100(1 - \alpha)\%$  confidence interval for the population proportion  $p$  is

$$\bar{X} \pm E, \quad E = z_{\alpha/2} \sqrt{\hat{p}(1 - \hat{p})/n}.$$

- $E$  is called the **margin of error**, and  $z_{\alpha/2}$  satisfies

# Confidence Interval for Population Proportion $p$ :

- If sample size  $n$  is large, i.e.  $n\hat{p} > 5$  and  $n\hat{q} = n(1 - \hat{p}) > 5$ ,
- the  $100(1 - \alpha)\%$  confidence interval for the population proportion  $p$  is

$$\bar{X} \pm E, \quad E = z_{\alpha/2} \sqrt{\hat{p}(1 - \hat{p})/n}.$$


- $E$  is called the **margin of error**, and  $z_{\alpha/2}$  satisfies

# Confidence Interval for Population Proportion $p$ :

- If sample size  $n$  is large, i.e.  $n\hat{p} > 5$  and  $n\hat{q} = n(1 - \hat{p}) > 5$ ,
- the  $100(1 - \alpha)\%$  confidence interval for the population proportion  $p$  is

$$\bar{X} \pm E, \quad E = z_{\alpha/2} \sqrt{\hat{p}(1 - \hat{p})/n}.$$

- $E$  is called the **margin of error**, and  $z_{\alpha/2}$  satisfies



# A Little Better Confidence Interval for $p$ :

- Since  $-z_{\alpha/2} \leq \frac{\bar{X}-p}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2}$  is equivalent to

$$\frac{|\bar{X}-p|}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2} \quad \text{or} \quad \frac{(\bar{X}-p)^2}{p(1-p)/n} \leq z_{\alpha/2}^2$$

- That is  $H(p) = (\bar{X}-p)^2 - z_{\alpha/2}^2 p(1-p)/n \leq 0$
- Let  $a = 1 + \frac{z_{\alpha/2}^2}{n}$ ,  $b = -2\bar{X} - \frac{z_{\alpha/2}^2}{n}$ ,  $c = \bar{X}^2$ .  
 $H(p) = ap^2 + bp + c$ .
- The solutions of  $H(p) = 0$  are

$$\hat{p}_L = \frac{-b - \sqrt{b^2 - 4ac}}{2a}, \quad \hat{p}_U = \frac{-b + \sqrt{b^2 - 4ac}}{2a}.$$

So an approximate  $100(1 - \alpha)\%$  CI for  $p$  is  $[\hat{p}_L, \hat{p}_U]$ .

# A Little Better Confidence Interval for $p$ :

- Since  $-z_{\alpha/2} \leq \frac{\bar{X}-p}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2}$  is equivalent to

$$\frac{|\bar{X}-p|}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2} \quad \text{or} \quad \frac{(\bar{X}-p)^2}{p(1-p)/n} \leq z_{\alpha/2}^2$$

- That is  $H(p) = (\bar{X} - p)^2 - z_{\alpha/2}^2 p(1 - p)/n \leq 0$

- Let  $a = 1 + \frac{z_{\alpha/2}^2}{n}$ ,  $b = -2\bar{X} - \frac{z_{\alpha/2}^2}{n}$ ,  $c = \bar{X}^2$ .  
 $H(p) = ap^2 + bp + c$ .

- The solutions of  $H(p) = 0$  are

$$\hat{p}_L = \frac{-b - \sqrt{b^2 - 4ac}}{2a}, \quad \hat{p}_U = \frac{-b + \sqrt{b^2 - 4ac}}{2a}.$$

So an approximate  $100(1 - \alpha)\%$  CI for  $p$  is  $[\hat{p}_L, \hat{p}_U]$ .

# A Little Better Confidence Interval for $p$ :

- Since  $-z_{\alpha/2} \leq \frac{\bar{X}-p}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2}$  is equivalent to

$$\frac{|\bar{X}-p|}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2} \quad \text{or} \quad \frac{(\bar{X}-p)^2}{p(1-p)/n} \leq z_{\alpha/2}^2$$

- That is  $H(p) = (\bar{X} - p)^2 - z_{\alpha/2}^2 p(1 - p)/n \leq 0$

- Let  $a = 1 + \frac{z_{\alpha/2}^2}{n}$ ,  $b = -2\bar{X} - \frac{z_{\alpha/2}^2}{n}$ ,  $c = \bar{X}^2$ .  
 $H(p) = ap^2 + bp + c$ .

- The solutions of  $H(p) = 0$  are

$$\hat{p}_L = \frac{-b - \sqrt{b^2 - 4ac}}{2a}, \quad \hat{p}_U = \frac{-b + \sqrt{b^2 - 4ac}}{2a}.$$

So an approximate  $100(1 - \alpha)\%$  CI for  $p$  is  $[\hat{p}_L, \hat{p}_U]$ .

# A Little Better Confidence Interval for $p$ :

- Since  $-z_{\alpha/2} \leq \frac{\bar{X}-p}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2}$  is equivalent to

$$\frac{|\bar{X}-p|}{\sqrt{p(1-p)/n}} \leq z_{\alpha/2} \quad \text{or} \quad \frac{(\bar{X}-p)^2}{p(1-p)/n} \leq z_{\alpha/2}^2$$

- That is  $H(p) = (\bar{X} - p)^2 - z_{\alpha/2}^2 p(1 - p)/n \leq 0$
- Let  $a = 1 + \frac{z_{\alpha/2}^2}{n}$ ,  $b = -2\bar{X} - \frac{z_{\alpha/2}^2}{n}$ ,  $c = \bar{X}^2$ .  
 $H(p) = ap^2 + bp + c$ .
- The solutions of  $H(p) = 0$  are

$$\hat{p}_L = \frac{-b - \sqrt{b^2 - 4ac}}{2a}, \quad \hat{p}_U = \frac{-b + \sqrt{b^2 - 4ac}}{2a}.$$

So an approximate  $100(1 - \alpha)\%$  CI for  $p$  is  $[\hat{p}_L, \hat{p}_U]$ .



## Example 1: Poll Results in The News

### Example 1: USA Today/Gallup Poll:

<http://www.usatoday.com/>

- (a) For a poll result  $\hat{p}$  of interest, find the sample size  $n$ , the margin of error, and calculate the 95% confidence interval for  $p$ .
- (b) Find the margin of error for 99% confidence level and confidence interval.

## Example 1: Poll Results in The News

### Example 1: USA Today/Gallup Poll:

<http://www.usatoday.com/>

- (a) For a poll result  $\hat{p}$  of interest, find the sample size  $n$ , the margin of error, and calculate the 95% confidence interval for  $p$ .
- (b) Find the margin of error for 99% confidence level and confidence interval.

## Example 2:

A machine shop manufactures toggle levers. A lever is flawed if a standard nut cannot be screwed onto the threads. Let  $p$  equal the proportion of flawed toggle levers that the shop manufactures. If there were 24 flawed levers out of a sample of 642 that were selected randomly from the production line.

- (a) Give a point estimate of  $p$ .
- (b) Find a 95% confidence interval for  $p$ .

## Solution of Example 2

### Solution

(a) The point estimate is  $\hat{p} = 24/642 = 0.037$ .

(b)  $\alpha = 0.05$ ,  $z = 1.96$ ,  $s_{\hat{p}} = \sqrt{\hat{p}(1 - \hat{p})/n} = 0.0075$ .

$E = 0.0147$  and 95% CI for  $p$  is  $[0.0223, 0.0517]$ .

If the machine shop wants to be 99% confident with a margin of error as small as 0.001, how large a sample should be selected?

What is the most conservative sample size?

## Solution of Example 2

### Solution

(a) The point estimate is  $\hat{p} = 24/642 = 0.037$ .

(b)  $\alpha = 0.05$ ,  $z = 1.96$ ,  $s_{\hat{p}} = \sqrt{\hat{p}(1 - \hat{p})/n} = 0.0075$ .

$E = 0.0147$  and 95% CI for  $p$  is  $[0.0223, 0.0517]$ .

If the machine shop wants to be 99% confident with a margin of error as small as 0.001, how large a sample should be selected?

What is the most conservative sample size?

# Outline

- 1 9.10 Estimating a Proportion
  - Estimating Single Proportion
  - Sample Size for Estimating  $p$
  
- 2 9.11 Difference of Two Proportions

# Determine Sample Size by Margin of Error

- If  $\hat{p}$  is available,

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

$$E^2 = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{n}$$

$$n = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{E^2} = \left( \frac{z_{\frac{\alpha}{2}} \sqrt{\hat{p}(1 - \hat{p})}}{E} \right)^2$$

- If  $\hat{p}$  is not available, use the *most conservative* estimate

$$n = \left( \frac{z_{\frac{\alpha}{2}}}{2E} \right)^2$$

# Determine Sample Size by Margin of Error

- If  $\hat{p}$  is available,

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

$$E^2 = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{n}$$

$$n = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{E^2} = \left( \frac{z_{\frac{\alpha}{2}} \sqrt{\hat{p}(1 - \hat{p})}}{E} \right)^2$$

- If  $\hat{p}$  is not available, use the *most conservative* estimate

$$n = \left( \frac{z_{\frac{\alpha}{2}}}{2E} \right)^2$$



# Determine Sample Size by Margin of Error

- If  $\hat{p}$  is available,

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

$$E^2 = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{n}$$

$$n = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{E^2} = \left( \frac{z_{\frac{\alpha}{2}} \sqrt{\hat{p}(1 - \hat{p})}}{E} \right)^2$$

- If  $\hat{p}$  is not available, use the *most conservative* estimate

$$n = \left( \frac{z_{\frac{\alpha}{2}}}{2E} \right)^2$$

# Determine Sample Size by Margin of Error

- If  $\hat{p}$  is available,

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

$$E^2 = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{n}$$

$$n = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{E^2} = \left( \frac{z_{\frac{\alpha}{2}} \sqrt{\hat{p}(1 - \hat{p})}}{E} \right)^2$$

- If  $\hat{p}$  is not available, use the *most conservative* estimate

$$n = \left( \frac{z_{\frac{\alpha}{2}}}{2E} \right)^2$$

# Determine Sample Size by Margin of Error

- If  $\hat{p}$  is available,

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

$$E^2 = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{n}$$

$$n = z_{\frac{\alpha}{2}}^2 \frac{\hat{p}(1 - \hat{p})}{E^2} = \left( \frac{z_{\frac{\alpha}{2}} \sqrt{\hat{p}(1 - \hat{p})}}{E} \right)^2$$

- If  $\hat{p}$  is not available, use the *most conservative* estimate

$$n = \left( \frac{z_{\frac{\alpha}{2}}}{2E} \right)^2$$

## Example 3:

A machine shop manufactures toggle levers. A lever is flawed if a standard nut cannot be screwed onto the threads. Let  $p$  equal the proportion of flawed toggle levers that the shop manufactures. If there were 24 flawed levers out of a sample of 642 that were selected randomly from the production line.

In §6.7 Example 2, have found a point estimate of  $p$ :  $\hat{p} = 0.037$  and the margin of error  $E = 0.0147$  with 95% confidence.

(a) If the machine shop wants to be 99% confident with a margin of error as small as 0.001, how large a sample should be selected?

(b) What is the most conservative sample size?

## Example 3:

A machine shop manufactures toggle levers. A lever is flawed if a standard nut cannot be screwed onto the threads. Let  $p$  equal the proportion of flawed toggle levers that the shop manufactures. If there were 24 flawed levers out of a sample of 642 that were selected randomly from the production line. In §6.7 Example 2, have found a point estimate of  $p$ :  $\hat{p} = 0.037$  and the margin of error  $E = 0.0147$  with 95% confidence.

- (a) If the machine shop wants to be 99% confident with a margin of error as small as 0.001, how large a sample should be selected?
- (b) What is the most conservative sample size?

## Example 3:

A machine shop manufactures toggle levers. A lever is flawed if a standard nut cannot be screwed onto the threads. Let  $p$  equal the proportion of flawed toggle levers that the shop manufactures. If there were 24 flawed levers out of a sample of 642 that were selected randomly from the production line.

In §6.7 Example 2, have found a point estimate of  $p$ :  $\hat{p} = 0.037$  and the margin of error  $E = 0.0147$  with 95% confidence.

(a) If the machine shop wants to be 99% confident with a margin of error as small as 0.001, how large a sample should be selected?

(b) What is the most conservative sample size?

# Solution of Example 3

## Solution

(a)  $\alpha = 0.01$ ,  $z_{\frac{\alpha}{2}} = 2.575$

$$n = \left( \frac{z}{E} \right)^2 \hat{p}(1 - \hat{p}) = \left( \frac{2.575}{0.001} \right)^2 (0.037)(1 - 0.037) = 236,256$$

(b) The most conservative sample size is

$$n = \left( \frac{z_{\frac{\alpha}{2}}}{2E} \right)^2 = \left( \frac{2.575}{2(0.001)} \right)^2 = 1,657,657$$

## Example 4:

- Consider two groups of women:
    - Group 1:** women who spend less than \$500 annually on clothes.
    - Group 2:** women who spend over \$1000 annually on clothes.
  - Let  $p_1$  and  $p_2$  be the proportions of women in the two groups who believe that the clothes are too expensive, respectively.
  - If  $Y_1 = 1009$  out of  $n_1 = 1230$  women from Group 1 and  $Y_2 = 207$  out of  $n_2 = 340$  from Group 2 believe that the clothes are too expensive.
- (a) Find a point estimate of  $p_1 - p_2$ .
- (b) Find an approximate 95% CI for  $p_1 - p_2$ .



## Example 4:

- Consider two groups of women:
    - Group 1:** women who spend less than \$500 annually on clothes.
    - Group 2:** women who spend over \$1000 annually on clothes.
  - Let  $p_1$  and  $p_2$  be the proportions of women in the two groups who believe that the clothes are too expensive, respectively.
  - If  $Y_1 = 1009$  out of  $n_1 = 1230$  women from Group 1 and  $Y_2 = 207$  out of  $n_2 = 340$  from Group 2 believe that the clothes are too expensive.
- (a) Find a point estimate of  $p_1 - p_2$ .
- (b) Find an approximate 95% CI for  $p_1 - p_2$ .

## Example 4:

- Consider two groups of women:
  - Group 1:** women who spend less than \$500 annually on clothes.
  - Group 2:** women who spend over \$1000 annually on clothes.
- Let  $p_1$  and  $p_2$  be the proportions of women in the two groups who believe that the clothes are too expensive, respectively.
- If  $Y_1 = 1009$  out of  $n_1 = 1230$  women from Group 1 and  $Y_2 = 207$  out of  $n_2 = 340$  from Group 2 believe that the clothes are too expensive.

(a) Find a point estimate of  $p_1 - p_2$ .

(b) Find an approximate 95% CI for  $p_1 - p_2$ .

## Example 4:

- Consider two groups of women:

**Group 1:** women who spend less than \$500 annually on clothes.

**Group 2:** women who spend over \$1000 annually on clothes.

- Let  $p_1$  and  $p_2$  be the proportions of women in the two groups who believe that the clothes are too expensive, respectively.
- If  $Y_1 = 1009$  out of  $n_1 = 1230$  women from Group 1 and  $Y_2 = 207$  out of  $n_2 = 340$  from Group 2 believe that the clothes are too expensive.

- (a) Find a point estimate of  $p_1 - p_2$ .
- (b) Find an approximate 95% CI for  $p_1 - p_2$ .

# Solution of Example 4:

- (a) Calculate the sample proportions

$$\hat{p}_1 = \frac{Y_1}{n_1} = 0.82, \quad \hat{p}_2 = \frac{Y_2}{n_2} = 0.61$$

The point estimate of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = 0.21$ .

- (b) Since  $Y_1$  and  $Y_2$  are independent r.v.'s.  $Y_1 \sim b(n_1, p_1)$  and  $Y_2 \sim b(n_2, p_2)$ , by the C.L.T.,

$$\frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}_1(1 - \hat{p}_1)/n_1 + \hat{p}_2(1 - \hat{p}_2)/n_2}} \sim N(0, 1)$$

From this we can approximate 100(1 -  $\alpha$ )% CI for  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 \pm E$ , where

$$E = z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} = 0.056.$$

Thus  $\hat{p}_1 - \hat{p}_2 \pm E = [0.154, 0.266]$ .

# Solution of Example 4:

- (a) Calculate the sample proportions

$$\hat{p}_1 = \frac{Y_1}{n_1} = 0.82, \quad \hat{p}_2 = \frac{Y_2}{n_2} = 0.61$$

The **point estimate** of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = 0.21$ .

- (b) Since  $Y_1$  and  $Y_2$  are independent r.v.'s.  $Y_1 \sim b(n_1, p_1)$  and  $Y_2 \sim b(n_2, p_2)$ , by the C.L.T.,

$$\frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}_1(1 - \hat{p}_1)/n_1 + \hat{p}_2(1 - \hat{p}_2)/n_2}} \sim N(0, 1)$$

From this we can approximate 100(1 -  $\alpha$ )% CI for  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 \pm E$ , where

$$E = z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} = 0.056.$$

Thus  $\hat{p}_1 - \hat{p}_2 \pm E = [0.154, 0.266]$ .

## Solution of Example 4:

- (a) Calculate the sample proportions

$$\hat{p}_1 = \frac{Y_1}{n_1} = 0.82, \quad \hat{p}_2 = \frac{Y_2}{n_2} = 0.61$$

The **point estimate** of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = 0.21$ .

- (b) Since  $Y_1$  and  $Y_2$  are independent r.v.'s.  $Y_1 \sim b(n_1, p_1)$  and  $Y_2 \sim b(n_2, p_2)$ , by the C.L.T.,

$$\frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}_1(1 - \hat{p}_1)/n_1 + \hat{p}_2(1 - \hat{p}_2)/n_2}} \sim N(0, 1)$$

From this we can approximate 100(1 -  $\alpha$ )% CI for  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 \pm E$ , where

$$E = z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} = 0.056.$$

Thus  $\hat{p}_1 - \hat{p}_2 \pm E = [0.154, 0.266]$ .

## Solution of Example 4:

- (a) Calculate the sample proportions

$$\hat{p}_1 = \frac{Y_1}{n_1} = 0.82, \quad \hat{p}_2 = \frac{Y_2}{n_2} = 0.61$$

The **point estimate** of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = 0.21$ .

- (b) Since  $Y_1$  and  $Y_2$  are independent r.v.'s.  $Y_1 \sim b(n_1, p_1)$  and  $Y_2 \sim b(n_2, p_2)$ , by the C.L.T.,

$$\frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}_1(1 - \hat{p}_1)/n_1 + \hat{p}_2(1 - \hat{p}_2)/n_2}} \sim N(0, 1)$$

From this we an **approximate**  $100(1 - \alpha)\%$  CI for  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 \pm E$ , where

$$E = z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} = 0.056.$$

Thus  $\hat{p}_1 - \hat{p}_2 \pm E = [0.154, 0.266]$ .

# Solution of Example 4:

- (a) Calculate the sample proportions

$$\hat{p}_1 = \frac{Y_1}{n_1} = 0.82, \quad \hat{p}_2 = \frac{Y_2}{n_2} = 0.61$$

The **point estimate** of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = 0.21$ .

- (b) Since  $Y_1$  and  $Y_2$  are independent r.v.'s.  $Y_1 \sim b(n_1, p_1)$  and  $Y_2 \sim b(n_2, p_2)$ , by the C.L.T.,

$$\frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}_1(1 - \hat{p}_1)/n_1 + \hat{p}_2(1 - \hat{p}_2)/n_2}} \sim N(0, 1)$$

From this we an **approximate 100(1 -  $\alpha$ )% CI** for  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 \pm E$ , where

$$E = z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} = 0.056.$$

Thus  $\hat{p}_1 - \hat{p}_2 \pm E = [0.154, 0.266]$ .



## Solution of Example 4:

- (a) Calculate the sample proportions

$$\hat{p}_1 = \frac{Y_1}{n_1} = 0.82, \quad \hat{p}_2 = \frac{Y_2}{n_2} = 0.61$$

The **point estimate** of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = 0.21$ .

- (b) Since  $Y_1$  and  $Y_2$  are independent r.v.'s.  $Y_1 \sim b(n_1, p_1)$  and  $Y_2 \sim b(n_2, p_2)$ , by the C.L.T.,

$$\frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}_1(1 - \hat{p}_1)/n_1 + \hat{p}_2(1 - \hat{p}_2)/n_2}} \sim N(0, 1)$$

From this we can **approximate 100(1 -  $\alpha$ )% CI** for  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 \pm E$ , where

$$E = z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} = 0.056.$$

Thus  $\hat{p}_1 - \hat{p}_2 \pm E = [0.154, 0.266]$ .