

# ***STATISTICAL INFERENCES (2cr)***

## Chapter 10 One- and Two-Sample Tests of Hypotheses

Zhong Guan

Math, IUSB

# Outline

- 1 10.1 Statistical Hypotheses: General Concepts
- 2 10.2 Testing a Statistical Hypothesis
- 3 10.3 Using  $p$ -value

## Example 1.

- Let  $p$  be the proportion of printed circuits that fail. Current procedure has  $p_0 = 0.06$ .
- A new method is proposed. To see if the new method results in an improvement.
- We want to test **statistical hypotheses**

$$H_0 : p = p_0 = 0.06 \quad \text{vs} \quad H_1 : p < p_0 = 0.06.$$

- We decided to select  $n = 200$  circuits to test. Let  $Y$  be the number of circuits that fail.
- Decision rule:** accept the improvement hypothesis  $H_1$  if  $Y \leq k$  (reject  $H_0$ ), otherwise reject  $H_1$  (accept  $H_0$ ).
- $k = ?$

## Example 1.

- Let  $p$  be the proportion of printed circuits that fail. Current procedure has  $p_0 = 0.06$ .
- A new method is proposed. To see if the new method results in an improvement.
- We want to test **statistical hypotheses**

$$H_0 : p = p_0 = 0.06 \quad \text{vs} \quad H_1 : p < p_0 = 0.06.$$

- We decided to select  $n = 200$  circuits to test. Let  $Y$  be the number of circuits that fail.
- Decision rule:** accept the improvement hypothesis  $H_1$  if  $Y \leq k$  (reject  $H_0$ ), otherwise reject  $H_1$  (accept  $H_0$ ).
- $k = ?$

# Example 1.

- Let  $p$  be the proportion of printed circuits that fail. Current procedure has  $p_0 = 0.06$ .
- A new method is proposed. To see if the new method results in an improvement.
- We want to test **statistical hypotheses**

$$H_0 : p = p_0 = 0.06 \quad \text{vs} \quad H_1 : p < p_0 = 0.06.$$

- We decided to select  $n = 200$  circuits to test. Let  $Y$  be the number of circuits that fail.
- Decision rule:** accept the improvement hypothesis  $H_1$  if  $Y \leq k$  (reject  $H_0$ ), otherwise reject  $H_1$  (accept  $H_0$ ).
- $k = ?$

## Example 1.

- Let  $p$  be the proportion of printed circuits that fail. Current procedure has  $p_0 = 0.06$ .
- A new method is proposed. To see if the new method results in an improvement.
- We want to test **statistical hypotheses**

$$H_0 : p = p_0 = 0.06 \quad \text{vs} \quad H_1 : p < p_0 = 0.06.$$

- We decided to select  $n = 200$  circuits to test. Let  $Y$  be the number of circuits that fail.
- Decision rule:** accept the improvement hypothesis  $H_1$  if  $Y \leq k$  (reject  $H_0$ ), otherwise reject  $H_1$  (accept  $H_0$ ).
- $k = ?$

## Example 1.

- Let  $p$  be the proportion of printed circuits that fail. Current procedure has  $p_0 = 0.06$ .
- A new method is proposed. To see if the new method results in an improvement.
- We want to test **statistical hypotheses**

$$H_0 : p = p_0 = 0.06 \quad \text{vs} \quad H_1 : p < p_0 = 0.06.$$

- We decided to select  $n = 200$  circuits to test. Let  $Y$  be the number of circuits that fail.
- **Decision rule:** accept the improvement hypothesis  $H_1$  if  $Y \leq k$  (reject  $H_0$ ), otherwise reject  $H_1$  (accept  $H_0$ ).
- $k = ?$

## Example 1.

- Let  $p$  be the proportion of printed circuits that fail. Current procedure has  $p_0 = 0.06$ .
- A new method is proposed. To see if the new method results in an improvement.
- We want to test **statistical hypotheses**

$$H_0 : p = p_0 = 0.06 \quad \text{vs} \quad H_1 : p < p_0 = 0.06.$$

- We decided to select  $n = 200$  circuits to test. Let  $Y$  be the number of circuits that fail.
- **Decision rule**: accept the improvement hypothesis  $H_1$  if  $Y \leq k$  (reject  $H_0$ ), otherwise reject  $H_1$  (accept  $H_0$ ).
- $k = ?$



## Two Types of Errors

- For example, we choose  $k = 7$  so that if  $Y \leq 7$  or  $\hat{p} \leq 0.035$ , then we accept the improvement hypothesis.
- However, whatever  $k$  is, the decision made based on a sample could be wrong. There are two types of errors.

	$H_0$ true	$H_0$ false
Reject $H_0$	Type I error	Correct
Accept $H_0$	Correct	Type II error

- If we accept  $H_0$ , then we will keep the current procedure. This is called no change hypothesis. It is also called **null hypothesis**.
- $H_1$  is called **alternative hypothesis**.
- $y \leq k$  is called **rejection** or **critical region** of  $H_0$ .
- $y$  or  $\hat{p}$  is called the **test statistic**.

# Two Types of Errors

- For example, we choose  $k = 7$  so that if  $Y \leq 7$  or  $\hat{p} \leq 0.035$ , then we accept the improvement hypothesis.
- However, whatever  $k$  is, the decision made based on a sample could be wrong. There are two types of errors.

	$H_0$ true	$H_0$ false
Reject $H_0$	Type I error	Correct
Accept $H_0$	Correct	Type II error

- If we accept  $H_0$ , then we will keep the current procedure. This is called no change hypothesis. It is also called **null hypothesis**.
- $H_1$  is called **alternative hypothesis**.
- $y \leq k$  is called **rejection** or **critical region** of  $H_0$ .
- $y$  or  $\hat{p}$  is called the **test statistic**.

## Two Types of Errors

- For example, we choose  $k = 7$  so that if  $Y \leq 7$  or  $\hat{p} \leq 0.035$ , then we accept the improvement hypothesis.
- However, whatever  $k$  is, the decision made based on a sample could be wrong. There are two types of errors.

	$H_0$ true	$H_0$ false
Reject $H_0$	Type I error	Correct
Accept $H_0$	Correct	Type II error

- If we accept  $H_0$ , then we will keep the current procedure. This is called no change hypothesis. It is also called **null hypothesis**.
- $H_1$  is called **alternative hypothesis**.
- $y \leq k$  is called **rejection** or **critical region** of  $H_0$ .
- $y$  or  $\hat{p}$  is called the test statistic.

# Two Types of Errors

- For example, we choose  $k = 7$  so that if  $Y \leq 7$  or  $\hat{p} \leq 0.035$ , then we accept the improvement hypothesis.
- However, whatever  $k$  is, the decision made based on a sample could be wrong. There are two types of errors.

	$H_0$ true	$H_0$ false
Reject $H_0$	Type I error	Correct
Accept $H_0$	Correct	Type II error

- If we accept  $H_0$ , then we will keep the current procedure. This is called no change hypothesis. It is also called **null hypothesis**.
- $H_1$  is called **alternative hypothesis**.
- $y \leq k$  is called **rejection** or **critical region** of  $H_0$ .
- $y$  or  $\hat{p}$  is called the test statistic.

## Two Types of Errors

- For example, we choose  $k = 7$  so that if  $Y \leq 7$  or  $\hat{p} \leq 0.035$ , then we accept the improvement hypothesis.
- However, whatever  $k$  is, the decision made based on a sample could be wrong. There are two types of errors.

	$H_0$ true	$H_0$ false
Reject $H_0$	Type I error	Correct
Accept $H_0$	Correct	Type II error

- If we accept  $H_0$ , then we will keep the current procedure. This is called no change hypothesis. It is also called **null hypothesis**.
- $H_1$  is called **alternative hypothesis**.
- $y \leq k$  is called **rejection** or **critical region** of  $H_0$ .
- $y$  or  $\hat{p}$  is called the **test statistic**.

## Two Types of Errors

- For example, we choose  $k = 7$  so that if  $Y \leq 7$  or  $\hat{p} \leq 0.035$ , then we accept the improvement hypothesis.
- However, whatever  $k$  is, the decision made based on a sample could be wrong. There are two types of errors.

	$H_0$ true	$H_0$ false
Reject $H_0$	Type I error	Correct
Accept $H_0$	Correct	Type II error

- If we accept  $H_0$ , then we will keep the current procedure. This is called no change hypothesis. It is also called **null hypothesis**.
- $H_1$  is called **alternative hypothesis**.
- $y \leq k$  is called **rejection** or **critical region** of  $H_0$ .
- $y$  or  $\hat{p}$  is called the **test statistic**.

# Two Types of Errors

- For example, we choose  $k = 7$  so that if  $Y \leq 7$  or  $\hat{p} \leq 0.035$ , then we accept the improvement hypothesis.
- However, whatever  $k$  is, the decision made based on a sample could be wrong. There are two types of errors.

	$H_0$ true	$H_0$ false
Reject $H_0$	Type I error	Correct
Accept $H_0$	Correct	Type II error

- If we accept  $H_0$ , then we will keep the current procedure. This is called no change hypothesis. It is also called **null hypothesis**.
- $H_1$  is called **alternative hypothesis**.
- $y \leq k$  is called **rejection** or **critical region** of  $H_0$ .
- $y$  or  $\hat{p}$  is called the **test statistic**.

# Chances to Make Mistakes: Type I Error

- The Type I error of the above rule is that  $Y \leq 7$  but the truth is  $p = 0.06$ .
- The probability of this error is

$$\begin{aligned}P(Y \leq 7 | p = 0.06) &= \sum_{y=0}^7 \binom{200}{y} (0.06)^y (0.94)^{200-y} \\&= 0.0829 = 8.29\%.\end{aligned}$$



## Chances to Make Mistakes: Type I Error

- The Type I error of the above rule is that  $Y \leq 7$  but the truth is  $p = 0.06$ .
- The probability of this error is

$$\begin{aligned}P(Y \leq 7 | p = 0.06) &= \sum_{y=0}^7 \binom{200}{y} (0.06)^y (0.94)^{200-y} \\&= 0.0829 = 8.29\%.\end{aligned}$$

## Chances to Make Mistakes: Type I Error

- The Type I error of the above rule is that  $Y \leq 7$  but the truth is  $p = 0.06$ .
- The probability of this error is

$$P(Y \leq 7 | p = 0.06) = \sum_{y=0}^7 \binom{200}{y} (0.06)^y (0.94)^{200-y}$$
$$= 0.0829 = 8.29\%.$$

## Chances to Make Mistakes: Type I Error

- The Type I error of the above rule is that  $Y \leq 7$  but the truth is  $p = 0.06$ .
- The probability of this error is

$$\begin{aligned} P(Y \leq 7 | p = 0.06) &= \sum_{y=0}^7 \binom{200}{y} (0.06)^y (0.94)^{200-y} \\ &= 0.0829 = 8.29\%. \end{aligned}$$

# Chances to Make Mistakes: Type II Error

- Type II error:  $Y > 7$  but the truth is  $p < 0.06$ .
- The probability of Type II error is, for  $p < 0.06$ ,

$$\beta(p) = P(Y > 7|p) = \sum_{y=8}^{200} \binom{200}{y} p^y (1-p)^{200-y}$$

- For example,

$$\beta(0.03) = 1 - \text{binomialcdf}(7, n = 200, p = 0.03) = 0.267$$

$$\beta(0.02) = 1 - \text{binomialcdf}(7, n = 200, p = 0.02) = 0.038$$

$$\beta(0.01) = 1 - \text{binomialcdf}(7, n = 200, p = 0.01) = 4.64 \times 10^{-5}$$

## Chances to Make Mistakes: Type II Error

- Type II error:  $Y > 7$  but the truth is  $p < 0.06$ .
- The probability of Type II error is, for  $p < 0.06$ ,

$$\beta(p) = P(Y > 7|p) = \sum_{y=8}^{200} \binom{200}{y} p^y (1-p)^{200-y}$$

- For example,

$$\beta(0.03) = 1 - \text{binomialcdf}(7, n = 200, p = 0.03) = 0.267$$

$$\beta(0.02) = 1 - \text{binomialcdf}(7, n = 200, p = 0.02) = 0.038$$

$$\beta(0.01) = 1 - \text{binomialcdf}(7, n = 200, p = 0.01) = 4.64 \times 10^{-5}$$

# Chances to Make Mistakes: Type II Error

- Type II error:  $Y > 7$  but the truth is  $p < 0.06$ .
- The probability of Type II error is, for  $p < 0.06$ ,

$$\beta(p) = P(Y > 7|p) = \sum_{y=8}^{200} \binom{200}{y} p^y (1-p)^{200-y}$$

- For example,

$$\beta(0.03) = 1 - \text{binomialcdf}(7, n = 200, p = 0.03) = 0.267$$

$$\beta(0.02) = 1 - \text{binomialcdf}(7, n = 200, p = 0.02) = 0.038$$

$$\beta(0.01) = 1 - \text{binomialcdf}(7, n = 200, p = 0.01) = 4.64 \times 10^{-5}$$

## Chances to Make Mistakes: Type II Error

- Type II error:  $Y > 7$  but the truth is  $p < 0.06$ .
- The probability of Type II error is, for  $p < 0.06$ ,

$$\beta(p) = P(Y > 7|p) = \sum_{y=8}^{200} \binom{200}{y} p^y (1-p)^{200-y}$$

- For example,

$$\beta(0.03) = 1 - \text{binomialcdf}(7, n = 200, p = 0.03) = 0.267$$

$$\beta(0.02) = 1 - \text{binomialcdf}(7, n = 200, p = 0.02) = 0.038$$

$$\beta(0.01) = 1 - \text{binomialcdf}(7, n = 200, p = 0.01) = 4.64 \times 10^{-5}$$

# Chances to Make Mistakes: Type II Error

- Type II error:  $Y > 7$  but the truth is  $p < 0.06$ .
- The probability of Type II error is, for  $p < 0.06$ ,

$$\beta(p) = P(Y > 7|p) = \sum_{y=8}^{200} \binom{200}{y} p^y (1-p)^{200-y}$$

- For example,

$$\beta(0.03) = 1 - \text{binomialcdf}(7, n = 200, p = 0.03) = 0.267$$

$$\beta(0.02) = 1 - \text{binomialcdf}(7, n = 200, p = 0.02) = 0.038$$

$$\beta(0.01) = 1 - \text{binomialcdf}(7, n = 200, p = 0.01) = 4.64 \times 10^{-5}$$



## Significance Level and Power of the Test

- Generally, we want to control the Type I error first: for a given **significance level**  $\alpha$ , choose  $k$  so that

$$P(\text{Type I error}) \leq \alpha$$

- The **power** of the test is,

$$\text{Power}(p) = 1 - \beta(p).$$

- In the above example, for  $p < 0.06$ ,

$$\text{Power}(p) = P(Y \leq 7|p) = 1 - P(Y > 7|p).$$

$$\text{Power}(0.03) = 0.833$$

$$\text{Power}(0.02) = 0.962,$$

$$\text{Power}(0.01) \approx 1.$$

# Significance Level and Power of the Test

- Generally, we want to control the Type I error first: for a given **significance level**  $\alpha$ , choose  $k$  so that

$$P(\text{Type I error}) \leq \alpha$$

- The **power** of the test is,

$$\text{Power}(p) = 1 - \beta(p).$$

- In the above example, for  $p < 0.06$ ,

$$\text{Power}(p) = P(Y \leq 7|p) = 1 - P(Y > 7|p).$$

$$\text{Power}(0.03) = 0.833$$

$$\text{Power}(0.02) = 0.962,$$

$$\text{Power}(0.01) \approx 1.$$

# Significance Level and Power of the Test

- Generally, we want to control the Type I error first: for a given **significance level**  $\alpha$ , choose  $k$  so that

$$P(\text{Type I error}) \leq \alpha$$

- The **power** of the test is,

$$\text{Power}(p) = 1 - \beta(p).$$

- In the above example, for  $p < 0.06$ ,

$$\text{Power}(p) = P(Y \leq 7|p) = 1 - P(Y > 7|p).$$

$$\text{Power}(0.03) = 0.833$$

$$\text{Power}(0.02) = 0.962,$$

$$\text{Power}(0.01) \approx 1.$$

# Find Critical Value $k$

If  $H_0$  is true, approximately,

$$Z = \frac{Y - np_0}{\sqrt{np_0(1-p_0)}} \sim N(0, 1).$$

Thus

$$\alpha = P(Y \leq k | p = p_0) \approx \Phi\left(\frac{k - np_0}{\sqrt{np_0(1-p_0)}}\right)$$

So

$$\frac{k - np_0}{\sqrt{np_0(1-p_0)}} = -Z_\alpha.$$

and

$$k = np_0 - z_\alpha \sqrt{np_0(1-p_0)}.$$

# Rejection Region of $H_0$

- The **rejection region** of  $H_0$  is  $R = \{Y : Y \leq k\}$
- or  $R = \left\{Z = \frac{Y - np_0}{\sqrt{np_0(1-p_0)}} \leq -z_\alpha\right\}$ .
- Here  $k$  and  $-z_\alpha$  are called the critical values of the **test statistics**  $Y$  and  $Z$  respectively.
- In the above example,  $n = 200$ ,  $p_0 = 0.06$ . Choose  $\alpha = 0.05$ .  $z_\alpha = z_{0.05} = 1.645$ .

$$k = np_0 - z_\alpha \sqrt{np_0(1-p_0)} = 6.42$$

- The rejection region of  $H_0$  is

$$R = \{Y : Y \leq 6.4167\}$$

or

$$R = \left\{Z = \frac{Y - 12}{3.358571} \leq -1.645\right\}.$$

# Rejection Region of $H_0$

- The **rejection region** of  $H_0$  is  $R = \{Y : Y \leq k\}$
- or  $R = \left\{Z = \frac{Y - np_0}{\sqrt{np_0(1-p_0)}} \leq -z_\alpha\right\}$ .
- Here  $k$  and  $-z_\alpha$  are called the critical values of the **test statistics**  $Y$  and  $Z$  respectively.
- In the above example,  $n = 200$ ,  $p_0 = 0.06$ . Choose  $\alpha = 0.05$ .  $z_\alpha = z_{0.05} = 1.645$ .

$$k = np_0 - z_\alpha \sqrt{np_0(1-p_0)} = 6.42$$

- The rejection region of  $H_0$  is

$$R = \{Y : Y \leq 6.4167\}$$

or

$$R = \left\{Z = \frac{Y - 12}{3.358571} \leq -1.645\right\}.$$

# Rejection Region of $H_0$

- The **rejection region** of  $H_0$  is  $R = \{Y : Y \leq k\}$
- or  $R = \left\{Z = \frac{Y - np_0}{\sqrt{np_0(1-p_0)}} \leq -z_\alpha\right\}$ .
- Here  $k$  and  $-z_\alpha$  are called the critical values of the **test statistics**  $Y$  and  $Z$  respectively.
- In the above example,  $n = 200$ ,  $p_0 = 0.06$ . Choose  $\alpha = 0.05$ .  $z_\alpha = z_{0.05} = 1.645$ .

$$k = np_0 - z_\alpha \sqrt{np_0(1-p_0)} = 6.42$$

- The rejection region of  $H_0$  is

$$R = \{Y : Y \leq 6.4167\}$$

or

$$R = \left\{Z = \frac{Y - 12}{3.358571} \leq -1.645\right\}.$$

# Rejection Region of $H_0$

- The **rejection region** of  $H_0$  is  $R = \{Y : Y \leq k\}$
- or  $R = \left\{Z = \frac{Y - np_0}{\sqrt{np_0(1-p_0)}} \leq -z_\alpha\right\}$ .
- Here  $k$  and  $-z_\alpha$  are called the critical values of the **test statistics**  $Y$  and  $Z$  respectively.
- In the above example,  $n = 200$ ,  $p_0 = 0.06$ . Choose  $\alpha = 0.05$ .  $z_\alpha = z_{0.05} = 1.645$ .

$$k = np_0 - z_\alpha \sqrt{np_0(1-p_0)} = 6.42$$

- The rejection region of  $H_0$  is

$$R = \{Y : Y \leq 6.4167\}$$

or

$$R = \left\{Z = \frac{Y - 12}{3.358571} \leq -1.645\right\}.$$



# Rejection Region of $H_0$

- The **rejection region** of  $H_0$  is  $R = \{Y : Y \leq k\}$
- or  $R = \left\{Z = \frac{Y - np_0}{\sqrt{np_0(1-p_0)}} \leq -z_\alpha\right\}$ .
- Here  $k$  and  $-z_\alpha$  are called the critical values of the **test statistics**  $Y$  and  $Z$  respectively.
- In the above example,  $n = 200$ ,  $p_0 = 0.06$ . Choose  $\alpha = 0.05$ .  $z_\alpha = z_{0.05} = 1.645$ .

$$k = np_0 - z_\alpha \sqrt{np_0(1-p_0)} = 6.42$$

- The rejection region of  $H_0$  is

$$R = \{Y : Y \leq 6.4167\}$$

or

$$R = \left\{Z = \frac{Y - 12}{3.358571} \leq -1.645\right\}.$$

# A Continuous Random Variable Example

**Example 2:** Let  $X$  be the Brinell hardness measurement of ductile iron substantially annealed. Assume that the distribution of  $X$  is  $N(\mu, 10^2)$ . We shall test the null hypothesis  $H_0 : \mu = 170$  against the alternative hypothesis  $H_1 : \mu > 170$  based on a sample of size  $n$ .

- (a) Define the test statistic;
- (b) Define a critical region with a significance level  $\alpha = 0.05$ .
- (c) A random sample of  $n = 25$  observations of  $X$  yielded the following measurements.

170 167 174 179 179 156 163 156 187 156 183 179 174 179  
170 156 187 179 183 174 187 167 159 170 179

Calculate the value of the test statistic.

- (d) State your conclusion.

# A Continuous Random Variable Example

**Example 2:** Let  $X$  be the Brinell hardness measurement of ductile iron substantially annealed. Assume that the distribution of  $X$  is  $N(\mu, 10^2)$ . We shall test the null hypothesis  $H_0 : \mu = 170$  against the alternative hypothesis  $H_1 : \mu > 170$  based on a sample of size  $n$ .

- (a) Define the test statistic;
- (b) Define a critical region with a significance level  $\alpha = 0.05$ .
- (c) A random sample of  $n = 25$  observations of  $X$  yielded the following measurements.

170 167 174 179 179 156 163 156 187 156 183 179 174 179  
170 156 187 179 183 174 187 167 159 170 179

Calculate the value of the test statistic.

- (d) State your conclusion.

# Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined critical value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$



## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

# Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

# Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$

for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

## Solution of Example 2

- (a) Since  $\bar{X}$  is an estimator of  $\mu$ , we use  $\bar{X}$  as the test statistic.
- (b) The rejection region of  $H_0 : \mu = 170$  in favor of  $H_1 : \mu > 170$ :

$$\bar{X} > c$$


for some pre-determined **critical** value  $c$  according to  $\alpha$ .

$$0.05 = \alpha = \text{Prob}(\text{Type I}) = ?$$

$$P(\text{Type I}) = P(\text{Rejec } H_0 \mid H_0 \text{ is true}) = P(\bar{X} > c \mid \mu = 170)$$

$$0.05 = \alpha = P\left(Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > \frac{c - 170}{10/\sqrt{n}}\right)$$

# Solution of Example 2




So

$$\frac{c - 170}{10/\sqrt{n}} = z_\alpha \rightarrow c = 170 + \frac{10z_\alpha}{\sqrt{n}}$$




## Solution of Example 2



So

$$\frac{c - 170}{10/\sqrt{n}} = z_\alpha \rightarrow c = 170 + \frac{10z_\alpha}{\sqrt{n}}$$

## Solution of Example 2



So

$$\frac{c - 170}{10/\sqrt{n}} = z_\alpha \quad \rightarrow \quad c = 170 + \frac{10z_\alpha}{\sqrt{n}}$$

## Solution of Example 2

The rejection region is

$$\bar{X} > 170 + \frac{10z_{\alpha}}{\sqrt{n}}, \quad \text{or} \quad Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > z_{\alpha}$$

If  $\alpha = 0.05$  then  $z_{\alpha} = 1.645$ .

(c)  $n = 25$ ,  $\bar{x} = 172.52$ ,  $\alpha = 0.05$ ,  $z_{\alpha} = 1.645$ .

$$z = \frac{\bar{x} - 170}{10/\sqrt{n}} = \frac{172.52 - 170}{10/\sqrt{25}} = 1.26$$

(d) Conclusion: Since  $z = 1.26 < z_{\alpha} = 1.645$ , don't reject  $H_0$ .

## Solution of Example 2

The rejection region is

$$\bar{X} > 170 + \frac{10z_{\alpha}}{\sqrt{n}}, \quad \text{or} \quad Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > z_{\alpha}$$

If  $\alpha = 0.05$  then  $z_{\alpha} = 1.645$ .

(c)  $n = 25$ ,  $\bar{x} = 172.52$ ,  $\alpha = 0.05$ ,  $z_{\alpha} = 1.645$ .

$$z = \frac{\bar{x} - 170}{10/\sqrt{n}} = \frac{172.52 - 170}{10/\sqrt{25}} = 1.26$$

(d) Conclusion: Since  $z = 1.26 < z_{\alpha} = 1.645$ , don't reject  $H_0$ .

## Solution of Example 2

The rejection region is

$$\bar{X} > 170 + \frac{10z_{\alpha}}{\sqrt{n}}, \quad \text{or} \quad Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > z_{\alpha}$$

If  $\alpha = 0.05$  then  $z_{\alpha} = 1.645$ .

(c)  $n = 25$ ,  $\bar{x} = 172.52$ ,  $\alpha = 0.05$ ,  $z_{\alpha} = 1.645$ .

$$z = \frac{\bar{x} - 170}{10/\sqrt{n}} = \frac{172.52 - 170}{10/\sqrt{25}} = 1.26$$

(d) Conclusion: Since  $z = 1.26 < z_{\alpha} = 1.645$ , don't reject  $H_0$ .

## Solution of Example 2

The rejection region is

$$\bar{X} > 170 + \frac{10z_{\alpha}}{\sqrt{n}}, \quad \text{or} \quad Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > z_{\alpha}$$

If  $\alpha = 0.05$  then  $z_{\alpha} = 1.645$ .

(c)  $n = 25$ ,  $\bar{x} = 172.52$ ,  $\alpha = 0.05$ ,  $z_{\alpha} = 1.645$ .

$$z = \frac{\bar{x} - 170}{10/\sqrt{n}} = \frac{172.52 - 170}{10/\sqrt{25}} = 1.26$$

(d) Conclusion: Since  $z = 1.26 < z_{\alpha} = 1.645$ , don't reject  $H_0$ .

## Solution of Example 2

The rejection region is

$$\bar{X} > 170 + \frac{10z_{\alpha}}{\sqrt{n}}, \quad \text{or} \quad Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > z_{\alpha}$$

If  $\alpha = 0.05$  then  $z_{\alpha} = 1.645$ .

(c)  $n = 25$ ,  $\bar{x} = 172.52$ ,  $\alpha = 0.05$ ,  $z_{\alpha} = 1.645$ .

$$z = \frac{\bar{x} - 170}{10/\sqrt{n}} = \frac{172.52 - 170}{10/\sqrt{25}} = 1.26$$

(d) Conclusion: Since  $z = 1.26 < z_{\alpha} = 1.645$ , don't reject  $H_0$ .

## Solution of Example 2

The rejection region is

$$\bar{X} > 170 + \frac{10z_{\alpha}}{\sqrt{n}}, \quad \text{or} \quad Z = \frac{\bar{X} - 170}{10/\sqrt{n}} > z_{\alpha}$$

If  $\alpha = 0.05$  then  $z_{\alpha} = 1.645$ .

(c)  $n = 25$ ,  $\bar{x} = 172.52$ ,  $\alpha = 0.05$ ,  $z_{\alpha} = 1.645$ .

$$z = \frac{\bar{x} - 170}{10/\sqrt{n}} = \frac{172.52 - 170}{10/\sqrt{25}} = 1.26$$

(d) **Conclusion:** Since  $z = 1.26 < z_{\alpha} = 1.645$ , don't reject  $H_0$ .



# The $p$ -value Approach

- One important statistic which is usually reported in statistical analysis is the  **$p$ -value**:
- the probability of rejecting true null  $H_0$  using the observed value of test statistic as the critical value.
- For example, for the above hypotheses  
 $H_0 : p = p_0$  vs  $H_1 : p < p_0$ , if the observed value of  $Y$  is  $y$ , then the observed value of  $Z$  is

$$Z = \frac{y - np_0}{\sqrt{np_0(1-p_0)}}$$

- the  $p$ -value is

$$\begin{aligned} p_{\text{value}} &= P(Z \leq z | p = p_0) = P(Y \leq y | p = p_0) \\ &\approx \Phi\left(\frac{y - np_0}{\sqrt{np_0(1-p_0)}}\right). \end{aligned}$$

# The $p$ -value Approach

- One important statistic which is usually reported in statistical analysis is the  **$p$ -value**:
- the probability of rejecting true null  $H_0$  using the observed value of test statistic as the critical value.
- For example, for the above hypotheses  
 $H_0 : p = p_0$  vs  $H_1 : p < p_0$ , if the observed value of  $Y$  is  $y$ , then the observed value of  $Z$  is

$$Z = \frac{y - np_0}{\sqrt{np_0(1-p_0)}}$$

- the  $p$ -value is

$$\begin{aligned} p_{\text{value}} &= P(Z \leq z | p = p_0) = P(Y \leq y | p = p_0) \\ &\approx \Phi\left(\frac{y - np_0}{\sqrt{np_0(1-p_0)}}\right). \end{aligned}$$

# The $p$ -value Approach

- One important statistic which is usually reported in statistical analysis is the  **$p$ -value**:
- the probability of rejecting true null  $H_0$  using the observed value of test statistic as the critical value.
- For example, for the above hypotheses  
 $H_0 : p = p_0$  vs  $H_1 : p < p_0$ , if the observed value of  $Y$  is  $y$ , then the observed value of  $Z$  is

$$Z = \frac{y - np_0}{\sqrt{np_0(1-p_0)}}$$

- the  $p$ -value is

$$\begin{aligned} p_{\text{value}} &= P(Z \leq z | p = p_0) = P(Y \leq y | p = p_0) \\ &\approx \Phi\left(\frac{y - np_0}{\sqrt{np_0(1-p_0)}}\right). \end{aligned}$$

# The $p$ -value Approach

- One important statistic which is usually reported in statistical analysis is the  **$p$ -value**:
- the probability of rejecting true null  $H_0$  using the observed value of test statistic as the critical value.
- For example, for the above hypotheses  
 $H_0 : p = p_0$  vs  $H_1 : p < p_0$ , if the observed value of  $Y$  is  $y$ , then the observed value of  $Z$  is

$$Z = \frac{y - np_0}{\sqrt{np_0(1-p_0)}}$$

- the  $p$ -value is

$$\begin{aligned} p_{\text{value}} &= P(Z \leq z | p = p_0) = P(Y \leq y | p = p_0) \\ &\approx \Phi\left(\frac{y - np_0}{\sqrt{np_0(1-p_0)}}\right). \end{aligned}$$

## Decision Based on $p$ -value

- If  $p\text{-value} < \alpha$ , then we reject  $H_0$  at the significance level  $\alpha$ . Otherwise accept  $H_0$ .
- If in Example 1, the observed value of  $Y$  is  $y = 4$ , then the  $p$ -value is

$$p_{\text{value}} \approx \Phi\left(\frac{4-12}{3.358571}\right) = 0.0086.$$

- So we can reject  $H_0$  at the significance level  $\alpha = 0.05$ . We can also reject  $H_0$  at the significance level  $\alpha = 0.01$ .

## Decision Based on $p$ -value

- If  $p\text{-value} < \alpha$ , then we reject  $H_0$  at the significance level  $\alpha$ . Otherwise accept  $H_0$ .
- If in Example 1, the observed value of  $Y$  is  $y = 4$ , then the  $p$ -value is

$$p_{\text{value}} \approx \Phi\left(\frac{4-12}{3.358571}\right) = 0.0086.$$

- So we can reject  $H_0$  at the significance level  $\alpha = 0.05$ . We can also reject  $H_0$  at the significance level  $\alpha = 0.01$ .

## Decision Based on $p$ -value

- If  $p\text{-value} < \alpha$ , then we reject  $H_0$  at the significance level  $\alpha$ . Otherwise accept  $H_0$ .
- If in Example 1, the observed value of  $Y$  is  $y = 4$ , then the  $p$ -value is

$$p_{\text{value}} \approx \Phi\left(\frac{4-12}{3.358571}\right) = 0.0086.$$

- So we can reject  $H_0$  at the significance level  $\alpha = 0.05$ . We can also reject  $H_0$  at the significance level  $\alpha = 0.01$ .

## Decision Based on $p$ -value

- If  $p\text{-value} < \alpha$ , then we reject  $H_0$  at the significance level  $\alpha$ . Otherwise accept  $H_0$ .
- If in Example 1, the observed value of  $Y$  is  $y = 4$ , then the  $p$ -value is

$$p_{\text{value}} \approx \Phi\left(\frac{4-12}{3.358571}\right) = 0.0086.$$

- So we can reject  $H_0$  at the significance level  $\alpha = 0.05$ . We can also reject  $H_0$  at the significance level  $\alpha = 0.01$ .



# $P$ -value—the formal definition

## Definition

If  $W$  is a test statistic, the  $p$ -value, or *attained significance level*, is the smallest level of significance  $\alpha$  for which the observed data indicate that the null hypothesis should be rejected.

Let  $RR_\alpha$  be the rejection region of significance level  $\alpha$ :

$$P(W \in RR_\alpha | H_0) = \alpha.$$

If the data resulted in an observed value  $w$  of  $W$ , then the

$$p\text{-value} = \min\{\alpha : w \in RR_\alpha\}.$$

Rejection Region:  $p\text{-value} < \alpha$ .

# $P$ -value—the formal definition

## Definition

If  $W$  is a test statistic, the  $p$ -value, or *attained significance level*, is the smallest level of significance  $\alpha$  for which the observed data indicate that the null hypothesis should be rejected.

Let  $RR_\alpha$  be the rejection region of significance level  $\alpha$ :

$$P(W \in RR_\alpha | H_0) = \alpha.$$

If the data resulted in an observed value  $w$  of  $W$ , then the

$$p\text{-value} = \min\{\alpha : w \in RR_\alpha\}.$$

Rejection Region:  $p\text{-value} < \alpha$ .

# $P$ -value—the formal definition

## Definition

If  $W$  is a test statistic, the  $p$ -value, or *attained significance level*, is the smallest level of significance  $\alpha$  for which the observed data indicate that the null hypothesis should be rejected.

Let  $RR_\alpha$  be the rejection region of significance level  $\alpha$ :

$$P(W \in RR_\alpha | H_0) = \alpha.$$

If the data resulted in an observed value  $w$  of  $W$ , then the

$$p\text{-value} = \min\{\alpha : w \in RR_\alpha\}.$$

**Rejection Region:**  $p\text{-value} < \alpha$ .

# Example

A health club claims that its members lose an average of 10 pounds or more within the 1st month after joining the club. A consumer agency wanted to test this claim.

A sample 36 members was selected and an average weight lost of 9.2 pounds was obtained. Assume the population standard deviation is known to be 2.4 pounds. What is the  $p$ -value of this test?

Solution:

Step 1. Hypotheses:

$$H_0 : \mu \geq 10 \quad H_A : \mu < 10$$

Step 2. Since  $\sigma = 2.4$  known,  $n = 36 > 30$ , use normal distribution and  $z$  test statistic  $Z = (\bar{X} - \mu_0)/(\sigma/\sqrt{n})$  and rejection region  $RR_\alpha = \{Z < -z_\alpha\}$ .

## Example

A health club claims that its members lose an average of 10 pounds or more within the 1st month after joining the club. A consumer agency wanted to test this claim.

A sample 36 members was selected and an average weight lost of 9.2 pounds was obtained. Assume the population standard deviation is known to be 2.4 pounds. What is the  $p$ -value of this test?

**Solution:**

**Step 1.** Hypotheses:

$$H_0 : \mu \geq 10 \quad H_A : \mu < 10$$

**Step 2.** Since  $\sigma = 2.4$  known,  $n = 36 > 30$ , use normal distribution and  $z$  test statistic  $Z = (\bar{X} - \mu_0)/(\sigma/\sqrt{n})$  and rejection region  $RR_\alpha = \{Z < -z_\alpha\}$ .

**Step 3.**  $\bar{x} = 9.2$ , the observed value of  $Z$  is  
 $z = (\bar{x} - \mu_0)/(\sigma/\sqrt{n}) = -2.00$ .

The  $p$ -value is

$$p\text{-value} = \min\{\alpha : -2.00 < -z_\alpha\} = P(Z < -2.00) = 0.0228$$

If  $\alpha = 0.01$ , then since  $p\text{-value} > 0.01$ , we do not reject  $H_0$  at the significance level 0.01.

But  $p\text{-value} < 0.05$ , we reject  $H_0$  at the significance level 0.05.

Step 3.  $\bar{x} = 9.2$ , the observed value of  $Z$  is  
 $z = (\bar{x} - \mu_0)/(\sigma/\sqrt{n}) = -2.00$ .

The  $p$ -value is

$$p\text{-value} = \min\{\alpha : -2.00 < -z_\alpha\} = P(Z < -2.00) = 0.0228$$

If  $\alpha = 0.01$ , then since  $p\text{-value} > 0.01$ , we do not reject  $H_0$  at the significance level 0.01.

But  $p\text{-value} < 0.05$ , we reject  $H_0$  at the significance level 0.05.

Step 3.  $\bar{x} = 9.2$ , the observed value of  $Z$  is  
 $z = (\bar{x} - \mu_0)/(\sigma/\sqrt{n}) = -2.00$ .

The  $p$ -value is

$$p\text{-value} = \min\{\alpha : -2.00 < -z_\alpha\} = P(Z < -2.00) = 0.0228$$

If  $\alpha = 0.01$ , then since  $p\text{-value} > 0.01$ , we do not reject  $H_0$  at the significance level 0.01.

But  $p\text{-value} < 0.05$ , we reject  $H_0$  at the significance level 0.05.



Step 3.  $\bar{x} = 9.2$ , the observed value of  $Z$  is  
 $z = (\bar{x} - \mu_0)/(\sigma/\sqrt{n}) = -2.00$ .

The  $p$ -value is

$$p\text{-value} = \min\{\alpha : -2.00 < -z_\alpha\} = P(Z < -2.00) = 0.0228$$

If  $\alpha = 0.01$ , then since  $p\text{-value} > 0.01$ , we do not reject  $H_0$  at the significance level 0.01.

But  $p\text{-value} < 0.05$ , we reject  $H_0$  at the significance level 0.05.

# Finding $p$ -value

Let the observed value of the test statistic be  $w$ . If the rejection region is  $\{W < k\}$ , then the  $p$ -value is

$$p\text{-value} = P(W < w | H_0).$$

Given a significance level  $\alpha$ , if  $p$ -value is smaller than  $\alpha$ , we reject the null hypothesis  $H_0$ . This is equivalent to the critical value approach.

Click [here](#) for tables of the commonly used tests.

## Finding $p$ -value

Let the observed value of the test statistic be  $w$ . If the rejection region is  $\{W < k\}$ , then the  $p$ -value is

$$p - \text{value} = P(W < w | H_0).$$

Given a significance level  $\alpha$ , if  $p$  - value is **smaller than**  $\alpha$ , we **reject** the null hypothesis  $H_0$ . This is equivalent to the critical value approach.

Click [here](#) for tables of the commonly used tests.

## Finding $p$ -value

Let the observed value of the test statistic be  $w$ . If the rejection region is  $\{W < k\}$ , then the  $p$ -value is

$$p - \text{value} = P(W < w | H_0).$$

Given a significance level  $\alpha$ , if  $p$  - value is **smaller than**  $\alpha$ , we **reject** the null hypothesis  $H_0$ . This is equivalent to the critical value approach.

Click **here** for tables of the commonly used tests.

# An Example of One-Sided Test

**Example 1.** Let  $x$  be the Brinell hardness measurement of ductile iron subcritically annealed. Assume that  $x$  has a normal distribution with mean  $\mu$ . Test  $H_0 : \mu = 170$  against the alternative hypothesis that the mean hardness is **greater than** 170 based on the following 25 observations (Use significance level  $\alpha = 0.05$ .)

170, 167, 174, 179, 179, 156, 163, 156, 187  
156, 183, 179, 174, 179, 170, 156, 187,  
179, 183, 174, 187, 167, 159, 170, 179

Hypotheses:

$$H_0 : \mu = 170 \quad H_1 : \mu > 170$$

# An Example of One-Sided Test

**Example 1.** Let  $x$  be the Brinell hardness measurement of ductile iron subcritically annealed. Assume that  $x$  has a normal distribution with mean  $\mu$ . Test  $H_0 : \mu = 170$  against the alternative hypothesis that the mean hardness is **greater than** 170 based on the following 25 observations (Use significance level  $\alpha = 0.05$ .)

170, 167, 174, 179, 179, 156, 163, 156, 187  
156, 183, 179, 174, 179, 170, 156, 187,  
179, 183, 174, 187, 167, 159, 170, 179

Hypotheses:

$$H_0 : \mu = 170 \quad H_1 : \mu > 170$$

# An Example of Two-Sided Test

**Example 2.** A company that manufactures brackets for an auto maker selected 15 brackets from the production line and performs a torque test. The goal is for mean torque to **equal** 125. Let the torque have a normal distribution. The 15 observations are

128, 149, 136, 114, 126, 142, 124,  
122, 118, 122, 129, 118, 122, 129, 136

Hypotheses:  $H_0 : \mu = 125$     $H_1 : \mu \neq 125$

# An Example of Two-Sided Test

**Example 2.** A company that manufactures brackets for an auto maker selected 15 brackets from the production line and performs a torque test. The goal is for mean torque to **equal** 125. Let the torque have a normal distribution. The 15 observations are


128, 149, 136, 114, 126, 142, 124,  
122, 118, 122, 129, 118, 122, 129, 136

Hypotheses:  $H_0 : \mu = 125$     $H_1 : \mu \neq 125$



## Another Example of Two-Sided Test


**Example 3.** A Casino want to test whether the probability  $p$  that a red number comes up on a Nevada roulette wheel equals  $18/38$  or not.



Hypotheses:  $H_0 : p = 18/38$     $H_1 : p \neq 18/38$ .

## Another Example of Two-Sided Test

**Example 3.** A Casino want to test whether the probability  $p$  that a red number comes up on a Nevada roulette wheel equals  $18/38$  or not.



Hypotheses:  $H_0 : p = 18/38$     $H_1 : p \neq 18/38$ .