

STATISTICAL INFERENCES (2cr)

Chapter 9 One- and Two-Sample Estimation Problems

Zhong Guan

Math, IUSB

Outline

1 9.12 Estimating a Variance

2 9.13 Estimating Ratio of Variances

Outline

- 1 9.12 Estimating a Variance
- 2 9.13 Estimating Ratio of Variances

Assumptions and Point Estimator

- Assume that X_1, \dots, X_n is a random sample from normal population with mean μ and variance σ^2 .
- Both μ and σ^2 are unknown.
- The point estimator is the sample variance $S^2 = (n-1)^{-1} \sum_{i=1}^n (X_i - \bar{X})^2$. This is an unbiased and consistent estimator.

Assumptions and Point Estimator

- Assume that X_1, \dots, X_n is a random sample from normal population with mean μ and variance σ^2 .
- Both μ and σ^2 are unknown.
- The point estimator is the sample variance $S^2 = (n-1)^{-1} \sum_{i=1}^n (X_i - \bar{X})^2$. This is an unbiased and consistent estimator.

Assumptions and Point Estimator

- Assume that X_1, \dots, X_n is a random sample from normal population with mean μ and variance σ^2 .
- Both μ and σ^2 are unknown.
- The point estimator is the sample variance $S^2 = (n-1)^{-1} \sum_{i=1}^n (X_i - \bar{X})^2$. This is an unbiased and consistent estimator.

Assumptions and Point Estimator

- Assume that X_1, \dots, X_n is a random sample from normal population with mean μ and variance σ^2 .
- Both μ and σ^2 are unknown.
- The point estimator is the sample variance $S^2 = (n-1)^{-1} \sum_{i=1}^n (X_i - \bar{X})^2$. This is an unbiased and consistent estimator.

Confidence Interval

- $X^2 = (n - 1)S^2/\sigma^2$ has a chi-squared distribution with $n - 1$ degrees of freedom.
- There exist a and b such that

$$\begin{aligned} 1 - \alpha &= P(a \leq X^2 \leq b) = P(a \leq (n - 1)S^2/\sigma^2 \leq b) \\ &= P((n - 1)S^2/b \leq \sigma^2 \leq (n - 1)S^2/a) \end{aligned}$$

- For convenience we choose $a = \chi_{1-\alpha/2}^2$ and $b = \chi_{\alpha/2}^2$.
Thus we have a $100(1 - \alpha)\%$ confidence interval for σ^2 :

$$\left[\frac{(n - 1)S^2}{\chi_{\alpha/2}^2}, \frac{(n - 1)S^2}{\chi_{1-\alpha/2}^2} \right]$$

Confidence Interval

- $X^2 = (n - 1)S^2/\sigma^2$ has a chi-squared distribution with $n - 1$ degrees of freedom.
- There exist a and b such that

$$\begin{aligned}1 - \alpha &= P(a \leq X^2 \leq b) = P(a \leq (n - 1)S^2/\sigma^2 \leq b) \\ &= P((n - 1)S^2/b \leq \sigma^2 \leq (n - 1)S^2/a)\end{aligned}$$

- For convenience we choose $a = \chi_{1-\alpha/2}^2$ and $b = \chi_{\alpha/2}^2$.
Thus we have a $100(1 - \alpha)\%$ confidence interval for σ^2 :

$$\left[\frac{(n - 1)S^2}{\chi_{\alpha/2}^2}, \frac{(n - 1)S^2}{\chi_{1-\alpha/2}^2} \right]$$

Confidence Interval

- $X^2 = (n - 1)S^2/\sigma^2$ has a chi-squared distribution with $n - 1$ degrees of freedom.
- There exist a and b such that

$$\begin{aligned}1 - \alpha &= P(a \leq X^2 \leq b) = P(a \leq (n - 1)S^2/\sigma^2 \leq b) \\&= P((n - 1)S^2/b \leq \sigma^2 \leq (n - 1)S^2/a)\end{aligned}$$

- For convenience we choose $a = \chi_{1-\alpha/2}^2$ and $b = \chi_{\alpha/2}^2$.
Thus we have a $100(1 - \alpha)\%$ confidence interval for σ^2 :

$$\left[\frac{(n - 1)S^2}{\chi_{\alpha/2}^2}, \frac{(n - 1)S^2}{\chi_{1-\alpha/2}^2} \right]$$

Find Critical Value χ^2_{α}

- Using Table A.5.
- Using TI 8x:
 - (1) graph cdf curve $y = \chi^2 \text{cdf}(0, X, df)$ and horizontal line $y = 1 - \alpha$;
 - (2) the x-coordinate of the intersection is χ^2_{α} .


Find Critical Value χ^2_{α}

- Using Table A.5.
- Using TI 8x:
 - (1) graph cdf curve $y = \chi^2cdf(0, X, df)$ and horizontal line $y = 1 - \alpha$;
 - (2) the x-coordinate of the intersection is χ^2_{α} .

Use TI-83/84 to Find Critical χ^2 Value


For example, $n = 20$, $df = n - 1 = 19$, to find $\chi^2_{.025}$:

- Enter equation for χ^2 cdf : $Y=$ -> $Y1 = \chi^2\text{cdf}(0, X, 19)$
- Enter equation: $Y2 = 1 - .025 = 0.975$.



```
Plot1 Plot2 Plot3
Y1=χ²cdf(0,X,19
Y2=.975
Y3=
Y4=
Y5=
Y6=
```


- Press **GRAPH** and adjust **WINDOW** if needed.
- Press **2ND** and **CALC** to find intersection of the two graphs. We can obtain $\chi^2_{.025} = 32.85$.



Use TI-83/84 to Find Critical χ^2 Value


For example, $n = 20$, $df = n - 1 = 19$, to find $\chi^2_{.025}$:

- Enter equation for χ^2 cdf : $Y_1 = \chi^2\text{cdf}(0, X, 19)$
- Enter equation: $Y_2 = 1 - .025 = 0.975$.



```
Plot1 Plot2 Plot3
Y1=X^2cdf(0,X,19
Y2=.975
Y3=
Y4=
Y5=
Y6=
```


- Press **GRAPH** and adjust **WINDOW** if needed.
- Press **2ND** and **CALC** to find intersection of the two graphs. We can obtain $\chi^2_{.025} = 32.85$.



Use TI-83/84 to Find Critical χ^2 Value

For example, $n = 20$, $df = n - 1 = 19$, to find $\chi^2_{.025}$:


- Enter equation for χ^2 cdf : $Y=$ $\rightarrow Y1 = \chi^2\text{cdf}(0, X, 19)$
- Enter equation: $Y2 = 1 - .025 = 0.975$.



```

Plot1 Plot2 Plot3
Y1=χ²cdf(0,X,19)
Y2=.975
Y3=
Y4=
Y5=
Y6=
    
```


- Press **GRAPH** and adjust **WINDOW** if needed.
- Press **2ND** and **CALC** to find intersection of the two graphs. We can obtain $\chi^2_{.025} = 32.85$.



Use TI-83/84 to Find Critical χ^2 Value


For example, $n = 20$, $df = n - 1 = 19$, to find $\chi^2_{.025}$:

- Enter equation for χ^2 cdf : $Y_1 = \chi^2 \text{cdf}(0, X, 19)$
- Enter equation: $Y_2 = 1 - .025 = 0.975$.



Plot1 Plot2 Plot3
Y1 $\chi^2 \text{cdf}(0, X, 19)$
Y2 $.975$
Y3 =
Y4 =
Y5 =
Y6 =


- Press **GRAPH** and adjust **WINDOW** if needed.
- Press **2ND** and **CALC** to find intersection of the two graphs. We can obtain $\chi^2_{.025} = 32.85$.



Use TI-83/84 to Find Critical χ^2 Value


For example, $n = 20$, $df = n - 1 = 19$, to find $\chi^2_{.025}$:

- Enter equation for χ^2 cdf : $Y_1 = \chi^2 \text{cdf}(0, X, 19)$
- Enter equation: $Y_2 = 1 - .025 = 0.975$.



Plot1 Plot2 Plot3
Y1 $\chi^2 \text{cdf}(0, X, 19)$
Y2 $.975$
Y3 =
Y4 =
Y5 =
Y6 =

- Press **GRAPH** and adjust **WINDOW** if needed.
- Press **2ND** and **CALC** to find intersection of the two graphs. We can obtain $\chi^2_{.025} = 32.85$.



Challenging Problems

- Find a and b so that

$$1 - \alpha = P((n-1)S^2/b \leq \sigma^2 \leq (n-1)S^2/a)$$

and $\frac{1}{a} - \frac{1}{b}$ is minimized.

- If μ is known, how do you estimate σ^2 using point estimator and confidence interval?

Assumptions and Point Estimator

- Assume that X_{11}, \dots, X_{1n_1} is a random sample from normal population with unknown mean μ_1 and variance σ_1^2 , and
- X_{21}, \dots, X_{2n_2} is a random sample from normal population with unknown mean μ_2 and variance σ_2^2 , and that
- the two samples are **independent**.
- The sample variances and sample means are

$$S_i^2 = (n_i - 1)^{-1} \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_i)^2,$$

$$\bar{X}_i = n_i^{-1} \sum_{j=1}^{n_i} X_{ij}, \quad i = 1, 2.$$

- The point estimator of σ_1^2/σ_2^2 is S_1^2/S_2^2 . This is a consistent but **biased** estimator.

Assumptions and Point Estimator

- Assume that X_{11}, \dots, X_{1n_1} is a random sample from normal population with unknown mean μ_1 and variance σ_1^2 , and
- X_{21}, \dots, X_{2n_2} is a random sample from normal population with unknown mean μ_2 and variance σ_2^2 , and that
- the two samples are **independent**.
- The sample variances and sample means are

$$S_i^2 = (n_i - 1)^{-1} \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_i)^2,$$

$$\bar{X}_i = n_i^{-1} \sum_{j=1}^{n_i} X_{ij}, \quad i = 1, 2.$$

- The point estimator of σ_1^2/σ_2^2 is S_1^2/S_2^2 . This is a consistent but **biased** estimator.

Assumptions and Point Estimator

- Assume that X_{11}, \dots, X_{1n_1} is a random sample from normal population with unknown mean μ_1 and variance σ_1^2 , and
- X_{21}, \dots, X_{2n_2} is a random sample from normal population with unknown mean μ_2 and variance σ_2^2 , and that
- the two samples are **independent**.
- The sample variances and sample means are

$$S_i^2 = (n_i - 1)^{-1} \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_i)^2,$$

$$\bar{X}_i = n_i^{-1} \sum_{j=1}^{n_i} X_{ij}, \quad i = 1, 2.$$

- The point estimator of σ_1^2/σ_2^2 is S_1^2/S_2^2 . This is a consistent but **biased** estimator.

Assumptions and Point Estimator

- Assume that X_{11}, \dots, X_{1n_1} is a random sample from normal population with unknown mean μ_1 and variance σ_1^2 , and
- X_{21}, \dots, X_{2n_2} is a random sample from normal population with unknown mean μ_2 and variance σ_2^2 , and that
- the two samples are **independent**.
- The sample variances and sample means are

$$S_i^2 = (n_i - 1)^{-1} \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_i)^2,$$

$$\bar{X}_i = n_i^{-1} \sum_{j=1}^{n_i} X_{ij}, \quad i = 1, 2.$$

- The point estimator of σ_1^2/σ_2^2 is S_1^2/S_2^2 . This is a consistent but **biased** estimator.

Assumptions and Point Estimator

- Assume that X_{11}, \dots, X_{1n_1} is a random sample from normal population with unknown mean μ_1 and variance σ_1^2 , and
- X_{21}, \dots, X_{2n_2} is a random sample from normal population with unknown mean μ_2 and variance σ_2^2 , and that
- the two samples are **independent**.
- The sample variances and sample means are

$$S_i^2 = (n_i - 1)^{-1} \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_i)^2,$$

$$\bar{X}_i = n_i^{-1} \sum_{j=1}^{n_i} X_{ij}, \quad i = 1, 2.$$

- The point estimator of σ_1^2/σ_2^2 is S_1^2/S_2^2 , This is a consistent but **biased** estimator.

Assumptions and Point Estimator

- Assume that X_{11}, \dots, X_{1n_1} is a random sample from normal population with unknown mean μ_1 and variance σ_1^2 , and
- X_{21}, \dots, X_{2n_2} is a random sample from normal population with unknown mean μ_2 and variance σ_2^2 , and that
- the two samples are **independent**.
- The sample variances and sample means are

$$S_i^2 = (n_i - 1)^{-1} \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_i)^2,$$

$$\bar{X}_i = n_i^{-1} \sum_{j=1}^{n_i} X_{ij}, \quad i = 1, 2.$$

- The point estimator of σ_1^2/σ_2^2 is S_1^2/S_2^2 , This is a consistent but **biased** estimator.

Confidence Interval

- $X_1^2 = (n_1 - 1)S_1^2/\sigma_1^2$ and $X_2^2 = (n_2 - 1)S_2^2/\sigma_2^2$ are independent and have chi-squared distributions with $v_1 = n_1 - 1$ and $v_2 = n_2 - 1$ degrees of freedom, respectively.
- So $F = \frac{X_1^2/v_1}{X_2^2/v_2} = \frac{S_1^2/\sigma_1^2}{S_2^2/\sigma_2^2}$ has an F distribution with numerator and denominator degrees of freedom $v_1 = n_1 - 1$ and $v_2 = n_2 - 1$, respectively.

Confidence Interval

- $X_1^2 = (n_1 - 1)S_1^2/\sigma_1^2$ and $X_2^2 = (n_2 - 1)S_2^2/\sigma_2^2$ are independent and have chi-squared distributions with $\nu_1 = n_1 - 1$ and $\nu_2 = n_2 - 1$ degrees of freedom, respectively.
- So $F = \frac{X_1^2/\nu_1}{X_2^2/\nu_2} = \frac{S_1^2/\sigma_1^2}{S_2^2/\sigma_2^2}$ has an F distribution with numerator and denominator degrees of freedom $\nu_1 = n_1 - 1$ and $\nu_2 = n_2 - 1$, respectively.

Confidence Interval

- $X_1^2 = (n_1 - 1)S_1^2/\sigma_1^2$ and $X_2^2 = (n_2 - 1)S_2^2/\sigma_2^2$ are independent and have chi-squared distributions with $\nu_1 = n_1 - 1$ and $\nu_2 = n_2 - 1$ degrees of freedom, respectively.
- So $F = \frac{X_1^2/\nu_1}{X_2^2/\nu_2} = \frac{S_1^2/\sigma_1^2}{S_2^2/\sigma_2^2}$ has an F distribution with numerator and denominator degrees of freedom $\nu_1 = n_1 - 1$ and $\nu_2 = n_2 - 1$, respectively.

Confidence Interval(Cont.)

- There exist a and b such that

$$\begin{aligned}1 - \alpha &= P(a \leq F \leq b) = P\left(a \leq \frac{S_1^2 \sigma_2^2}{S_2^2 \sigma_1^2} \leq b\right) \\&= P\left(\frac{S_1^2}{b S_2^2} \leq \frac{\sigma_1^2}{\sigma_2^2} \leq \frac{S_1^2}{a S_2^2}\right)\end{aligned}$$

- For convenience we choose $a = F_{1-\alpha/2}$ and $b = F_{\alpha/2}$.

Thus we have a $100(1 - \alpha)\%$ confidence interval for $\frac{\sigma_1^2}{\sigma_2^2}$:

$$\left[\frac{S_1^2}{F_{\alpha/2} S_2^2}, \frac{S_1^2}{F_{1-\alpha/2} S_2^2} \right]$$

Confidence Interval(Cont.)

- There exist a and b such that

$$\begin{aligned} 1 - \alpha &= P(a \leq F \leq b) = P\left(a \leq \frac{S_1^2 \sigma_2^2}{S_2^2 \sigma_1^2} \leq b\right) \\ &= P\left(\frac{S_1^2}{b S_2^2} \leq \frac{\sigma_1^2}{\sigma_2^2} \leq \frac{S_1^2}{a S_2^2}\right) \end{aligned}$$

- For convenience we choose $a = F_{1-\alpha/2}$ and $b = F_{\alpha/2}$.
Thus we have a $100(1 - \alpha)\%$ confidence interval for $\frac{\sigma_1^2}{\sigma_2^2}$:

$$\left[\frac{S_1^2}{F_{\alpha/2} S_2^2}, \frac{S_1^2}{F_{1-\alpha/2} S_2^2} \right]$$

Find Critical Value F_{α}


- Using Table A.6.
- Using TI 8x:
 - (1) graph $Fcdf$ $Y = Fcdf(0, X, v_1, v_2)$ and horizontal line $Y = 1 - \alpha$;
 - (2) the x -coordinate of the intersection is F_{α} .

Find Critical Value F_{α}

- Using Table A.6.
- Using TI 8x:
 - (1) graph $Fcdf$ $Y = Fcdf(0, X, v_1, v_2)$ and horizontal line $Y = 1 - \alpha$;
 - (2) the x -coordinate of the intersection is F_{α} .

The F Critical Value using Technology


- In Excel: $F_{.025} = F.INV.RT(0.025, 5, 7) = 5.285236852$, or $=F.INV(1-0.025, 5, 7) = 5.285236852$.
- In TI-8x: Graph $Y_1 = Fcdf(0, X, 5, 7)$ and $Y_2 = 1 - .025 = 0.975$, the x-coordinate of the intersection is $F_{.025} = 5.2852369$.



- In R: $F_{.975} = \text{"qf(.975, 5, 7)"} = 5.285237$.

The F Critical Value using Technology


- In Excel: $F_{.025} = F.INV.RT(0.025, 5, 7) = 5.285236852$, or $=F.INV(1-0.025, 5, 7) = 5.285236852$.
- In TI-8x: Graph $Y_1 = Fcdf(0, X, 5, 7)$ and $Y_2 = 1 - .025 = 0.975$, the x-coordinate of the intersection is $F_{.025} = 5.2852369$.



- In R: $F_{.975} = \text{"qf(.975, 5, 7)"} = 5.285237$.

The F Critical Value using Technology

- In Excel: $F_{.025} = F.INV.RT(0.025, 5, 7) = 5.285236852$, or $=F.INV(1-0.025, 5, 7) = 5.285236852$.
- In TI-8x: Graph $Y_1 = Fcdf(0, X, 5, 7)$ and $Y_2 = 1 - .025 = 0.975$, the x-coordinate of the intersection is $F_{.025} = 5.2852369$.



- In R: $F_{.975} = \text{"qf(.975, 5, 7)"} = 5.285237$.

Challenging Problems

- Find a and b so that

$$1 - \alpha = P\left(\frac{S_1^2}{bS_2^2} \leq \frac{\sigma_1^2}{\sigma_2^2} \leq \frac{S_1^2}{aS_2^2}\right)$$

and $\frac{1}{a} - \frac{1}{b}$ is minimized.

- If μ_1 and μ_2 are known, how do you estimate σ_1^2/σ_2^2 using point estimator and confidence interval?