

# Proof of existence of square roots

Joaquín Gómez

April 4, 2025

Let  $T$  be a subset of the real numbers given by

$$T = \{t \in \mathbb{R} : t^2 < 2\}$$

We want to prove that the supremum of this set, which should be irrational, exists. We define  $\alpha = \sup T$ . We want to get that  $\alpha^2 = 2$  by ruling out the possibilities of  $\alpha^2 < 2$  and  $\alpha^2 > 2$

First, imagine a line of real numbers, we can try to push the supremum by a factor of  $1/n$

$$\begin{aligned}\left(\alpha + \frac{1}{n}\right)^2 &= \alpha^2 + \frac{2\alpha}{n} + \frac{1}{n^2} \\ &< \alpha^2 + \frac{2\alpha}{n} + \frac{1}{n} \\ &= \alpha^2 + \frac{2\alpha + 1}{n}\end{aligned}$$

We now assume that there's some value of  $n$  such that

$$\alpha^2 + \frac{2\alpha + 1}{n} < 2$$

After a rearrangement we get

$$\frac{2\alpha + 1}{n} > 2 - \alpha^2$$

Now, we can get back to the first inequality. Notice that

$$\left(\alpha + \frac{1}{n}\right)^2 < \alpha^2 + (2 - \alpha^2) = 2$$

This contradicts the fact that  $\alpha = \sup T$ . Thus, it means that we still can “move towards the right” after  $\alpha$  and get a number in  $T$ .

Now, we look to a contradiction for  $\alpha^2 > 2$ . Assuming this is true, we could subtract some amount to  $\alpha$  and get a number in  $T$ .

$$\begin{aligned}\left(\alpha - \frac{1}{n}\right)^2 &= \alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2} \\ &> \alpha^2 - \frac{2\alpha}{n}\end{aligned}$$

Now, analogous to the first part of the Proof

$$\alpha^2 - \frac{2\alpha}{n} > 2$$

Rearranging the inequality we get  $2 - \alpha^2 < \frac{-2\alpha}{n}$ , this means that

$$\left(\alpha - \frac{1}{n}\right)^2 > \alpha^2 + (2 - \alpha^2) = 2$$

This means that  $\alpha^2 > 2$  is false.

And because both  $\alpha^2 < 2$  and  $\alpha^2 > 2$  are false, this means that  $\alpha^2 = 2$ , which proves the existence of the square roots in  $\mathbb{R}$ .