

Exercício 7

Variables Entered/Removed

Model	Variables Entered	Variables Removed	Method
1	Horas, QI ^a	.	Enter

a. All requested variables entered.

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,884 ^a	,782	,719	10,096

a. Predictors: (Constant), Horas, QI

b. Dependent Variable: Resultado

ANOVA^b

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	2554,156	2	1277,078	12,530	,005 ^a
	Residual	713,444	7	101,921		
	Total	3267,600	9			

a. Predictors: (Constant), Horas, QI

b. Dependent Variable: Resultado

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients		
		B	Std. Error	Beta	t	Sig.
1	(Constant)	-124,568	47,874		-2,602	,035
	QI	1,659	,446	,733	3,719	,007
	Horas	1,439	1,067	,266	1,349	,219

a. Dependent Variable: Resultado

Coefficients^a

Model		95,0% Confidence Interval for B	
		Lower Bound	Upper Bound
1	(Constant)	-237,773	-11,364
	QI	,604	2,714
	Horas	-1,083	3,962

a. Dependent Variable: Resultado

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Studentized Residual	10	100,0%	0	,0%	10	100,0%

Descriptives

		Statistic	Std. Error
Studentized Residual	Mean	,0095740	,31978363
95% Confidence Interval for Mean	Lower Bound	-,7138268	
	Upper Bound	,7329748	
	5% Trimmed Mean	,0325599	
	Median	,4270071	
	Variance	1,023	
	Std. Deviation	1,01124464	
	Minimum	-1,63807	
	Maximum	1,24347	
	Range	2,88154	
	Interquartile Range	1,70487	
	Skewness	-,405	,687
	Kurtosis	-1,443	1,334

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Studentized Residual	,260	10	,054	,893	10	,184

a. Lilliefors Significance Correction

Studentized Residual

Normal Q-Q Plot of Studentized Residual



