

Asymptotics of Partition Functions of Coulomb Gases

João Victor Alcântara Pimenta

Dissertação de Mestrado do Programa de Pós-Graduação em
Matemática (PPG-Mat)

SERVIÇO DE PÓS-GRADUAÇÃO DO ICMC-USP

Data de Depósito:

Assinatura: _____

João Victor Alcântara Pimenta

Asymptotics of Partition Functions of Coulomb Gases

Master dissertation submitted to the Institute of Mathematics and Computer Sciences – ICMC-USP, in partial fulfillment of the requirements for the degree of the Master Program in Mathematics. *EXAMINATION BOARD PRESENTATION COPY*

Concentration Area: Mathematics

Advisor: Prof. Dr. Guilherme Silva

USP – São Carlos
July de 2026

Ficha catalográfica elaborada pela Biblioteca Prof. Achille Bassi
e Seção Técnica de Informática, ICMC/USP,
com os dados fornecidos pelo(a) autor(a)

A634m	Antonelli, Humberto Lidio Modelo de teses e dissertações em LaTeX do ICMC / Humberto Lidio Antonelli; orientadora Renata Pontin de Mattos Fortes. -- São Carlos, 2017. 79 p. Tese (Doutorado - Programa de Pós-Graduação em Ciências de Computação e Matemática Computacional) -- Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo, 2017. 1. Modelo. 2. Monografia de qualificação. 3. Dissertação. 4. Tese. 5. Latex. I. Fortes, Renata Pontin de Mattos, orient. II. Título.
-------	---

João Victor Alcântara Pimenta

Asymptotics of Partition Functions of Coulomb Gases

Master dissertation submitted to the Institute of Mathematics and Computer Sciences – ICMC-USP, in partial fulfillment of the requirements for the degree of the Master Program in Mathematics. *EXAMINATION BOARD PRESENTATION COPY*

Concentration Area: Mathematics

Advisor: Prof. Dr. Guilherme Silva

USP – São Carlos
July de 2026

*"En remontant chez moi pour y passer la soirée à travailler de mon mieux,
je me disais que le monde n'est pas construit pour l'équilibre.*

Le monde est désordre.

L'équilibre n'est pas la règle, c'est l'exception.

Et je faisais le serment de travailler pour l'ordre et l'équilibre.

Que de serments! Tu vas sourire."

G.Duhamel, Maitres, 1937

ACKNOWLEDGEMENTS

For all of those who were before and are now:

Your courage, patience, daring, and caring has brought me here.

To hoping that I become deserving of such act, for I feel loved and dear; of immense luck.

For all of those who will be after:

To promise that I can for you as it was done for me.

*“João, é bom ver que você aprendeu e fez tantas coisas bonitas!
Continue assim, muito feliz! Brinque e aprenda muito.”
Te amo muito, Papai.*

RESUMO

JOÃO V. A. PIMENTA. **Assintóticas de Funções Partição em Gases de Coulomb**. 2026. 49 p. Dissertação (Mestrado em Ciências – Matemática) – Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo, São Carlos – SP, 2026.

As matrizes aleatórias podem ser vistas de duas formas: como uma função mensurável de um espaço de probabilidade num conjunto de matrizes ou, mais concretamente, como uma matriz cujas entradas são variáveis aleatórias com alguma distribuição conjunta. Um conjunto de tais matrizes com uma dada distribuição arbitrária é designado por ensemble. Para ensembles de interesse, pode-se estudar estatísticas como autovetores e distribuições de autovalores. As simetrias do conjunto escolhido irão refletir sobre quais modelos tais estatísticas podem representar. Por exemplo, as distâncias dos valores característicos de uma matriz simétrica com coeficientes aleatórios modelam as distâncias entre níveis sucessivos de energias de núcleos atômicos pesados. Mais interessante ainda, existe uma certa universalidade nas estatísticas encontradas entre classes de ensembles, sugerindo a relevância da teoria no estudo da universalidade nos fenómenos modelados em geral. Particularmente importante é o estudo de tais matrizes quando estas crescem infinitamente, considerando assintoticamente o seu comportamento estatístico. O nosso foco é uma técnica para derivar assintóticas para a chamada função partição de alguns ensembles interessantes de matrizes aleatórias.

Palavras-chave: Teoria de Matrizes Aleatórias, Ensembles de Matrizes Planares, Assintótica.

ABSTRACT

JOÃO V. A. PIMENTA. **Asymptotics of Partition Functions of Coulomb Gases** . 2026. 49 p. Dissertação (Mestrado em Ciências – Matemática) – Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo, São Carlos – SP, 2026.

Random Matrices can be thought of in two ways: As a measurable function from a probability space into a set of matrices or, more concretely, as a matrix whose entries are random variables with some joint distribution. A set of such matrices with a given arbitrary distribution is called an ensemble. One can, for ensembles of interest, study statistic such as the eigenvectors and eigenvalue distributions. The symmetries of chosen set will reflect on what models such statistics can model. For example, distances of the characteristic values of a symmetric matrix with random coefficients model distances between successive levels of energies of heavy nuclei. More interestingly, there is some universality in the statistics found among classes of ensembles, suggesting the relevance of the theory in studying the universality in general modeled phenomena. Particularly important is the study of such matrices when they grow infinitely large, taking asymptotically its statistical behavior. We focus on a technique for deriving asymptotics for the so-called partition function of some interesting ensembles of random matrices.

Keywords: Random Matrix Theory, Planar Matrix Ensembles, Asymptotics.

LIST OF FIGURES

Figure 1 – In the Plot it is represented the plot for $e^{-\lambda(\cosh(x)-1)}$ and an increasing $\lambda = 1, 2, 10, 70$ - darker represent larger values of the parameter. The dotted patten represents $e^{-\lambda x^2}$ for large lambda.	27
Figure 2 – The lines represent the solution to $\text{Im}[f(z) - f(t_c)] = 0$ for the three possible cases of the Hermite Polynomial. These are the curves we use to perform the asymptotic analysis on the integral. Note the case $\alpha = 2$; the coalescence of critical point gives origin to the Airy function.	39

CONTENTS

1	INTRODUCTION	19
2	RANDOM MATRIX THEORY	21
3	ASYMPTOTICS	23
3.1	Quadrature Technique	23
3.2	Asymptotic on Integrals	25
3.2.1	<i>Watson's Lemma</i>	27
3.2.2	<i>Laplace Method</i>	31
3.2.3	<i>Steepest Descent</i>	36
4	EQUILIBRIUM	41
5	PLANAR ENSEMBLES	43
	BIBLIOGRAPHY	45

INTRODUCTION

RANDOM MATRIX THEORY

ASYMPTOTICS

There is a sense in the previous Chapters that we want to study some distributions when our matrices are large enough - be that for feasibility of computations or approximation of infinite dimension operators. We also have mentioned that our main interest here is the Invariant Ensembles, or specially in Orthogonal Polynomial ones. Because of this, studying large matrices is equivalent, in some sense, to studying the regime of large n for orthogonal polynomials. For some of those polynomials, a integral representation is available, allowing us to use asymptotic methods on integrals to solve for the large n behavior. This is the main discussion of this chapter.

Techniques and structuring here are based on the book by [Olver *et al.* \(2010\)](#) and [Alpert \(1999\)](#) for quadrature and on the lecture notes by [Silva \(2021\)](#) for asymptotics on integrals.

3.1 Quadrature Technique

Before doing asymptotics on integrals we discuss a technique that will become important when approximating large summations: quadratures, i.e., approximating integrals by summations and vice-versa. One can see that asymptotics on integrals can then also perform a role on discrete summations. We first start briefly introducing a useful tool in the development of the formula we wish to use: the Bernoulli numbers B_n , polynomials $B_n(x)$ and periodic polynomials $\tilde{B}_n(x)$. The Bernoulli numbers B_n are a sequence of rational numbers that frequently appears in the Taylor series expansion of trigonometric-type functions. These number can also be seen as special values obtained by the Bernoulli polynomials $B_n(x)$. This is because these polynomials can be written explicitly as

$$B_k(x) = \sum_{n=0}^k \binom{k}{n} B_n x^{k-n},$$

then, we must have that $B_k(0) = B_k$. More interestingly for us is the fact that these polynomials satisfy the differential equation

$$B'_n(x) = nB_{n-1}(x), \quad n = 1, 2, \dots \quad (3.1)$$

Finally, the periodic Bernoulli polynomial is simply the Bernoulli polynomials evaluated at the fractional part of the argument, i.e., $\tilde{B}_n(x) = B_n(x - \lfloor x \rfloor)$. Now, let B_n and $B_n(x)$ denote the n th Bernoulli number and polynomial, respectively, and $\tilde{B}_n(x)$ the n th Bernoulli periodic function $B_n(x - \lfloor x \rfloor)$. Assume that a, m and n are integers such that $n > a$, $m > 0$, and $f^{(2m)}(x)$ is absolutely integrable over $[a, n]$. Then,

Theorem 1. (*Euler-Maclaurin Formula, (OLVER et al., 2010)*) Following the context given above,

$$\begin{aligned} \sum_{j=a}^n f(j) &= \int_a^n f(x)dx + \frac{1}{2}(f(a) + f(n)) + \sum_{s=1}^{m-1} \frac{B_{2s}}{(2s)!} \left(f^{(2s-1)}(n) - f^{(2s-1)}(a) \right) \\ &\quad + \int_a^n \frac{B_{2m} - \tilde{B}_{2m}(x)}{(2m)!} f^{(2m)}(x)dx. \end{aligned}$$

This identity is called the *Euler-Maclaurin* formula. It can be extremely useful when estimating the asymptotic of the summation on the function f when its integral can be more easily evaluate. In essence, this is a quadrature technique; a numerical integration: the approximation of an integral $\int f(x)dx$ of some function f by a discrete summation $\sum w_i f(x_i)$ over a set of points x_i with some weight function w_i . For example, in the classical Riemann sums, one approximates the integral $\int f(x)dx$ over some interval $[a, b]$ with the limit of the summation $\sum f(x_i^*)\Delta x_i$, where $x_i^* \in [x_{i-1}, x_i]$ and $\Delta x_i = x_i - x_{i-1}$. Different methods can be generated by choosing x_i^* differently.

The trapezoidal rule, for example, is a Riemann sum where x_i^* is chosen to be the mean point $x_i^* = (x_{i-1} + x_i)/2$. In this way, we would estimate the area using trapezoids, as the name suggests. Then, one could write

$$\begin{aligned} \int_a^n f(x)dx &\approx \sum_{j=0}^n f(x_j^*) = \sum_{j=0}^n \frac{f(x_j + x_{j-1})}{2}, \\ &= \sum_{j=1}^{n-1} f(x_j^*) + \frac{1}{2}(f(n) + f(0)) = \sum_{j=0}^{n-1} f(x_j^*) + \frac{1}{2}(f(n) - f(n)). \end{aligned}$$

Of course, this has some error term. Suppose that for some (x_i, x_{i-1}) we have the same k -derivatives at both extreme points for every $k \geq 1$, i.e., the function is constant. Then, there is no error in the approximation. If however, the two extreme points differ only on the k th derivative, there is some curvature defined by this difference where the error comes from. Therefore, there is indication of some error function given in respect to higher order derivatives.

Proof. Following the computations in Alpert (1999), take a function $f \in C^p(\mathbb{R})$ for $p \geq 1$. To derive the Euler-Maclaurin for the interval $[a, b]$, let $h = (b - a)/n$ and $c = a + jh$, then we express

$$\int_a^b f(x)dx = \sum_{j=0}^{n-1} \int_{a+jh}^{a+(j+1)h} f(x)dx$$

where we use (3.1) and integration by parts to rewrite each integral as

$$\begin{aligned} \int_c^{c+h} f(x)dx &= h \int_0^1 B_0(1-x) f(c+xh)dx \\ &= -h \frac{B_1(1-x)}{1!} f(c+xh) \Big|_0^1 + h^2 \int_0^1 \frac{B_1(1-x)}{1!} f'(c+xh)dx, \\ &\vdots \\ &= - \sum_{r=0}^{p-1} \frac{h^{r+1} B_{r+1}(1-x)}{(r+1)!} f^{(r)}(c+xh) \Big|_0^1 + h^{p+1} \int_0^1 \frac{h^{r+1} B_p(1-x)}{(p)!} f^{(p)}(c+xh)dx, \\ &= h \frac{f(c) + f(c+h)}{2} - \sum_{r=0}^{p-1} \frac{h^{r+1} B_{r+1}(1-x)}{(r+1)!} [f^{(r)}(c+h) - f^{(r)}(c)] \\ &\quad + h^{p+1} \int_0^1 \frac{B_p(1-x)}{(p)!} f^{(p)}(c+xh)dx. \end{aligned}$$

where one can start to notice, for instance, the trapezoidal rule creeping out on the first term. Notice that it was necessary to use that $-B_1(0) = B_1(1) = 1/2$ and $B_n(0) = B_n(1) = B_n$ for $n \neq 1$. Returning to the summation, we rearrange terms as

$$\begin{aligned} h \left[\frac{f(a)}{2} + f(a+h) + \cdots + f(b-h) + \frac{f(b)}{2} \right] &= \frac{1}{2} (f(b) - f(a)) + \sum_{j=0}^{n-1} f(a+jh), \\ &= \int_a^b f(x)dx + \sum_{r=0}^{p-1} \frac{h^{r+1} B_{r+1}(1-x)}{(r+1)!} [f^{(r)}(b) - f^{(r)}(a)] - h^{p+1} \int_0^1 \frac{B_p(1-x)}{(p)!} f^{(p)}(c+xh)dx. \end{aligned}$$

□

3.2 Asymptotic on Integrals

The techniques we will expand on here refers to a specific type of complex function $\phi : \mathbb{C} \rightarrow \mathbb{R}$ in the regime where $\lambda \rightarrow \infty$. Those functions are of the form

$$\phi(\lambda) := \int_{\Gamma} g(s) e^{-\lambda f(s)} ds, \quad (3.2)$$

where Γ is a given contour in the plane and f, g are suitable functions. Suitable here means that we impose g and f to be continuous (or analytic); g is integrable; f have global minima with zero derivative and strictly positive second derivative at this point of minima. There is also a more technical regularity condition on f to ensure the approximations hold to still be made clear.

Example 1. (Airy Equation) Function such as in (3.2) are not uncommon. Consider the function that satisfies the equation $\psi''(x) = x\psi(x)$. Taking the Fourier transform we have

$$-\omega^2 \Phi(x) = i\Phi'(x)$$

for which we solve with $\Phi(x) = e^{-i\omega^3/3}$ and then take the inverse Fourier Transform to get

$$\phi(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i/3\omega^3 + ix\omega} d\omega, \quad x \in \mathbb{R}.$$

The imaginary part of the argument is an odd function and therefore zero when integrated. The real part is expressed

$$\phi(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \cos\left(\frac{\omega^3}{3} + x\omega\right) d\omega.$$

This is the Airy function. For this integral, the techniques applied in this chapter are of great use to understand its behavior as $\omega \rightarrow \infty$.

Moreover, one can somehow motivate the conditions we imposed on ϕ . Consider $\phi : \mathbb{R} \rightarrow \mathbb{R}$, f with a single global minima and $\Gamma \equiv \mathbb{R}$ - for simplicity; the argument's idea should hold for other choices given the appropriate considerations. If ϕ is as in (3.2), and satisfying the conditions given for such function, then take x^* the single minimal critical point for f . We first note that if that is true, then, for any $|\epsilon| > 0$, the following limit should hold:

$$\lim_{\lambda \rightarrow \infty} \frac{e^{-\lambda(f(x^*))}}{e^{-\lambda(f(x^*-\epsilon))}} = \lim_{\lambda \rightarrow \infty} e^{-\lambda(f(x^*)-f(x^*-\epsilon))} = \lim_{\lambda \rightarrow \infty} e^{\lambda|c|} \rightarrow +\infty.$$

That is, the main contribution for the integral comes from a single point. The main idea then is to separate the integral in two regions: A first interval $(x^* - c/\sqrt{\lambda}, x^* + c/\sqrt{\lambda})$, $c > 0$, including this critical point and another interval that in theory should contribute only in error for the integral asymptotic. We start by approximating f around its global maximum by Taylor's theorem

$$f(x) = f(x^*) - \frac{1}{2}|f''(x^*)|(x-x^*)^2 + \epsilon(x-x^*)$$

and then perform a change of variables $\frac{x}{\sqrt{\lambda}} = s - x^*$, so that

$$\int_{x^* - \frac{c}{\sqrt{\lambda}}}^{x^* + \frac{c}{\sqrt{\lambda}}} g(s) e^{-\lambda f(s)} d(s) = \frac{e^{-\lambda f(x^*)}}{\sqrt{\lambda}} \int_{-c}^c g\left(x^* + \frac{x}{\sqrt{\lambda}}\right) e^{\frac{1}{2}|f''(x^*)|x^2 - \epsilon\left(\frac{x^3}{\sqrt{\lambda}}\right)} dx.$$

Now, using the technical assumption of regularity on the function f , one can argue that the error term ϵ goes to zero sufficiently fast, leaving a Gaussian Integral. Moreover, one needs to argue that the integral outside this small neighborhood of x^* is exponentially smaller than $e^{-\lambda f(x^*)}$. This should follow by an argument where the asymptotic for the integral

$$\int_{x^* + \frac{c}{\sqrt{\lambda}}}^{\infty} |g(s)| e^{-\lambda f(s)} d(s) \leq e^{-\lambda f(x^* + c/\sqrt{\lambda})} \|g\|_{L^1}$$




Figure 1 – In the Plot it is represented the plot for $e^{-\lambda(\cosh(x)-1)}$ and an increasing $\lambda = 1, 2, 10, 70$ - darker represent larger values of the parameter. The dotted patten represents $e^{-\lambda x^2}$ for large lambda.

is exponentially smaller than the previous region.

With this, we only note a few details of interest. Firstly, more precise error estimates can be obtained if one uses higher order terms of the Taylor Series expansion of f . For maximizer x^* which have null second derivative (a double critical point), the Laplace Approximation can be used expanding to the third order terms of the Taylor Series $f(x) = f(x^*) = \frac{1}{6}f'''(x^*)(x-x^*)^3 + \epsilon(x-x^*)$ and making the change of variables $t/\sqrt[3]{n} = x-x^*$ instead. This argument and result could also be used on curves on the plane. This is motivated from the fact that every (smooth) curve is locally a straight line. This will lead us to the method of *Steepest Descent*. These next subsections are in the spirit of proving a bit more formally the expansion given by the *Laplace Method* and introducing the mentioned *Steepest Descent*.

3.2.1 Watson's Lemma

We start this topic on the real line by looking at the functions $F : \mathbb{R} \rightarrow \mathbb{R}$ of the form

$$F(\lambda) = \int_0^T f(t) e^{-\lambda t} dt, \quad (3.3)$$

which one may think of as the Laplace transform of a function f supported on $[0, T]$. Note that F fits well with the models we described previously, and so one should expect the main contribution to come from the minima of the exponent function. Let us formalize the conditions for such expectation and the consequent result.

Theorem 2. (Watson's Lemma) Take $f \in L^1(0, T)$ and suppose that for some $\delta > 0$ it is of the form

$$f(t) = t^\alpha f_0(t), \quad 0 < t \leq \delta,$$

where $\alpha \in (-1, \infty)$ and f_0 is continuously differentiable on $[0, \delta]$ with $f_0(0) \neq 0$. Then the

function F in (3.3) satisfies

$$F(\lambda) = \frac{f_0(0)\Gamma(\alpha+1)}{\lambda^{\alpha+1}}(1 + O(\lambda^{-1}))$$

when $\lambda \rightarrow \infty$. Also, Γ is the usual Gamma function.

We must first mention that a weak estimation of the integral could be given by

$$\int_0^T f(t) e^{-\lambda t} dt \leq \|f\|_{L^1(0,T)}$$

where we only use the fact that $f \in L^1(0, T)$. However, we have more information on the behavior of the function in the section $0 < t < \delta$. Because of that fact, it is expected that in this set we could refine the approximation.

Proof. We start by decomposing the function in two parts

$$F(\lambda) = \int_0^\delta f(t) e^{-\lambda t} dt + \int_\delta^T f(t) e^{-\lambda t} dt =: (1) + (2).$$

For (2), we use the same weak estimate as commented before,

$$\int_\delta^T f(t) e^{-\lambda t} dt \leq e^{-\lambda \delta} \int_\delta^T |f(t)| dt = O(e^{-\lambda \delta} \|f\|_{L^1(\delta, T)}).$$

We keep it this way as we have no further information on the f in this section of the real line. This leave us with

$$F(\lambda) = \int_0^\delta f(t) e^{-\lambda t} dt + O(e^{-\lambda \delta} \|f\|_{L^1(\delta, T)}).$$

Now, we remind of the condition of continuous differentiability and the hypothesis on $f(t)$ to get $f(t) = t^\alpha (f_0(0) + r(t))$ for some differentiable error function $r(t)$ satisfying $|r(t)| \leq \|f'_0\|_{L^\infty(0, \delta)} t$ with $\|f'_0\|_{L^\infty(0, \delta)} = \inf \{c \geq 0 \mid \mu|_{(0, \delta)}(\{|f'_0| > c\}) = 0\}$ as usual. Then, write (1) as

$$(1) = \int_0^\delta t^\alpha (f_0(0) + r(t)) e^{-\lambda t} dt = f_0(0) \int_0^\delta t^\alpha e^{-\lambda t} dt + \int_0^\delta t^\alpha r(t) e^{-\lambda t} dt.$$

For these two integrals we simplify the problem by rewriting

$$\begin{aligned} \int_0^\delta t^\alpha e^{-\lambda t} dt &= \int_0^\infty t^\alpha e^{-\lambda t} dt - \int_\delta^\infty t^\alpha e^{-\lambda t} dt = \frac{\Gamma(\alpha+1)}{\lambda^{\alpha+1}} - \int_\delta^\infty t^\alpha e^{-\lambda t} dt, \\ \left| \int_0^\delta t^\alpha r(t) e^{-\lambda t} dt \right| &\leq \|f'_0\|_{L^\infty(0, \delta)} \int_0^\delta t^{\alpha+1} e^{-\lambda t} dt, \\ &= \|f'_0\|_{L^\infty(0, \delta)} \left(\int_0^\infty t^{\alpha+1} e^{-\lambda t} dt - \int_\delta^\infty t^{\alpha+1} e^{-\lambda t} dt \right), \\ &= \|f'_0\|_{L^\infty(0, \delta)} \left(\frac{\Gamma(\alpha+2)}{\lambda^{\alpha+2}} - \int_\delta^\infty t^{\alpha+1} e^{-\lambda t} dt \right). \end{aligned}$$

We are now only missing one ingredient to get the result, the order of

$$G(\lambda, \beta) := \int_{\delta}^{\infty} t^{\beta} e^{-\lambda t} dt, \quad \lambda > 1, \quad \beta > -1.$$

Using Cauchy-Schwartz we estimate

$$\begin{aligned} |G(\lambda, \beta)|^2 &\leq \left(\int_{\delta}^{\infty} e^{-\lambda t} dt \right) \left(\int_{\delta}^{\infty} t^{2\beta} e^{-\lambda t} dt \right), \\ &= \left(\frac{e^{-\lambda \delta}}{\lambda} \right) \left(\int_{\delta}^{\infty} t^{2\beta} e^{-\lambda t} dt \right), \\ &\stackrel{(u=\lambda t)}{=} \left(\frac{e^{-\lambda \delta}}{\lambda} \right) \left(\int_{\delta}^{\infty} e^{-u} u^{2\beta} \frac{1}{\lambda^{2\beta+1}} du \right), \\ &= \frac{e^{-\lambda \delta}}{\lambda^{2\beta+2}} M_{\beta} \end{aligned}$$

where M_{β} is some constant independent of λ . That leave us with

$$G(\lambda, \beta) = O\left(\frac{e^{-\lambda \delta/2}}{\lambda^{\beta+1}}\right) = O\left(e^{-\epsilon \lambda}\right)$$

where $\epsilon > 0$ depends on β and δ . Finally we rewrite

$$\int_0^{\delta} t^{\alpha} e^{-\lambda t} dt = \frac{\Gamma(\alpha+1)}{\lambda^{\alpha+1}} - G(\lambda, \alpha) = \frac{\Gamma(\alpha+1)}{\lambda^{\alpha+1}} + O\left(e^{-\epsilon \lambda}\right),$$

$$\begin{aligned} \left| \int_0^{\delta} t^{\alpha} r(t) e^{-\lambda t} dt \right| &\leq \|f'\|_{L^{\infty}(0, \delta)} \left(\frac{\Gamma(\alpha+2)}{\lambda^{\alpha+2}} - G(\lambda, \alpha+1) \right) \\ &= \|f'\|_{L^{\infty}(0, \delta)} \left(\frac{\Gamma(\alpha+2)}{\lambda^{\alpha+2}} + O\left(e^{-\epsilon \lambda}\right) \right) = O\left(\lambda^{-\alpha-2}\right), \end{aligned}$$

and

$$F(\lambda) = O\left(\lambda^{-\alpha-2}\right) + f_0(0) \left(\frac{\Gamma(\alpha+1)}{\lambda^{\alpha+1}} + O\left(e^{-\epsilon \lambda}\right) \right) + \|f\|_{L^1(\delta, T)} e^{-\delta \lambda}.$$

By the dominance of the exponential error, we get the result

$$F(\lambda) = \frac{f_0(0)\Gamma(\alpha+1)}{\lambda^{\alpha+1}} (1 + O(\lambda^{-1})).$$

□

In the estimation above, the Gamma function and the factor of λ^{-1} is the result of integrating the linear exponent of $e^{-\lambda t}$. If we, however, change this exponent to be of the general type $-\lambda t^d$ we still have a similar result on the behavior of the function.

Proposition 1. For $T, S > 0$, define the function

$$G(\lambda) := \int_{-S}^T g(t) e^{-\lambda t^2} dt$$

and assume $g(x)$ absolutely integrable in $[-S, T]$, of class C^1 in a fixed neighborhood of the origin. Then the function satisfies

$$G(\lambda) = \frac{g(0)\Gamma(1/2)}{\lambda^{1/2}} + O\left(\frac{1}{\lambda^{3/2}}\right)$$

when $\lambda \rightarrow \infty$ and where Γ is the Gamma function.

Proof. The proof is a reduction of the problem to Watson's Lemma. Assume, $T > S$ and write

$$G(\lambda) = \int_{-S}^S g(t) e^{-\lambda t^2} dt + \int_S^T g(t) e^{-\lambda t^2} dt.$$

The second integral we know to be of order

$$\int_S^T |g(t)| e^{-\lambda t^2} dt \leq \|g\|_\infty e^{-\lambda S^2},$$

exponential on λ , so we drop it. For the first integral we rewrite

$$\begin{aligned} \int_{-S}^S g(t) e^{-\lambda t^2} dt &= \int_0^S (g(t) + g(-t)) e^{-\lambda t^2} dt, \\ &\stackrel{(u=t^2)}{=} \frac{1}{2} \int_0^{S^2} \left(\frac{g(\sqrt{u}) + g(-\sqrt{u})}{\sqrt{u}} \right) e^{-\lambda u} du \end{aligned}$$

where now we use Watson's Lemma to finish the proof. $\square$

Example 2. Take the function called Gamma given by the expression

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt. \quad (3.4)$$

We are interested in what happens to the function when we take $x \rightarrow +\infty$. To see what happens, let us fit it into [Equation 2](#). Rewrite (3.4) as

$$\int_0^\infty e^{-t+x \log(t) - \log(t)} dt.$$

This lead us to identify ϕ as $-\log(t)$, which would give us $\phi'(t) < 0$, that is, we would have its minimum at $t = +\infty$. However $e^{\log(t)} = t^1$ is not integrable near the origin. Something need to change about the identification. Let us introduce a variable to rescale the integral: $xs = t$, s.t.,

$$\Gamma(x) = x^x \int_0^\infty e^{-\lambda(s-\log(s))} e^{-s} ds,$$

where we have used $\lambda = x - 1$. Thus we identify $\phi(s) = s - \log(s)$ and $g(s) = e^{-s}$. In fact, now ϕ has a global minimum at $s = 1$ and $\phi''(1) = 1 \neq 0$. Hence

$$\begin{aligned} \Gamma(x) &= \frac{\sqrt{2\pi} x^x e^{-x}}{(x-1)^{(-1/2)}} (1 + O(x^{-1})), \\ &= \sqrt{2\pi} x^{x+1/2} e^{-x} (1 + O(x^{-1})), \end{aligned}$$

which is called the Stirling formula.

3.2.2 Laplace Method

A seemly extension of the type of function we were dealing with are functions of the type

$$\Phi(\lambda) := \int_a^b g(t) e^{-\lambda\phi(t)} dt \quad (3.5)$$

for $a, b \in \mathbb{R} \cup \{-\infty, \infty\}$ and $a < b$ and a well behaved function g . For now, ϕ is a real-valued function. To begin the study let us make some assumptions. The first one is that a is finite, that means that we set $a = 0$ without loss of generality. The second assumption is that ϕ is a increasing function on $[0, b]$, so that $\phi(0) < \phi(t)$ for any t in the interval. With that we note that the coefficient $e^{-\lambda\phi(t)} / e^{-\lambda\phi(0)}$ decreases exponentially, that is, the integral is accumulated as the neighborhood of 0, i.e.

$$\Phi(\lambda) \approx \int_0^\delta g(t) e^{-\lambda\phi(t)} dt.$$

As we know $\phi(t)$ to be increasing, its first non zero derivative must be of even order, so that when we expanded the function by Taylor,

$$\phi(t) \approx \phi(0) + \phi_{(2k+1)}(0)t^{2k+1},$$

for some $k \geq 0$ and $|t| < \delta$ where $\phi_{(2k+1)} := \phi_{(2k+1)} / (2k+1)!$. As ϕ_{2k+1} , the $(2k+1)$ th-derivative, must be positive we also have that, as $\lambda \rightarrow \infty$,

$$\begin{aligned} \Phi(\lambda) &\approx e^{-\lambda\phi(0)} \int_0^\delta g(t) e^{-\lambda\phi_{(2k+1)}(0)t^{2k+1}} dt, \\ &\stackrel{(u=t(\lambda\phi_{(2k+1)}(0))^{1/(2k+1)})}{\approx} e^{-\lambda\phi(0)} \frac{g(0)}{(\lambda\phi_{(2k+1)}(0))^{\frac{1}{2k+1}}} \int_0^\infty e^{-u^{2k+1}} du, \\ &\stackrel{(s=u^{2k+1})}{\approx} e^{-\lambda\phi(0)} \frac{g(0)}{(2k+1)(\lambda\phi_{(2k+1)}(0))^{\frac{1}{2k+1}}} \int_0^\infty e^{-s} ds. \end{aligned}$$

where the integral can be evaluated as a Gamma function. More than anything we need to focus on an important assumption that was made: that ϕ has a global minima at one of the endpoints of integration. If that were not the case, for example if $a < 0 < b$ and $\phi(t)$ had a global minima at $t = 0$ we would now have to consider the expansion

$$\phi(t) \approx \phi(0) + \phi_{2k}(0)t^{2k}$$

which would lead us to the expression

$$\Phi(\lambda) \approx e^{-\lambda\phi(0)} \frac{g(0)}{(\lambda\phi_{2k}(0))^{\frac{1}{2k}}} \int_{-\infty}^\infty e^{-u^{2k}} du.$$

This differentiates of the previously result more importantly by the limits of integration. This is closely related to physical boundaries of the problem. Before enunciating the proper Laplace Method we state the Implicit Function Theorem, which will become important in a series of results that follows.

Theorem 3. (*Implicit Function Theorem*) Assume that S is an open subset of $\mathbb{R}^2$ and that $F : S \rightarrow \mathbb{R}$ is a function of class C^1 . Assume also that (a, b) is a point in S such that

$$F(a, b) = 0, \quad \text{and} \quad D_y F(a, b) \neq 0.$$

Then

1. There exist $r_0, r_1 > 0$ such that for every $x \in \mathbb{R}$ such that $|x - a| < r_0$, there exists a unique $y \in \mathbb{R}$ such that $|y - b| < r_1$

$$F(x, y) = 0.$$

That is, defines a function $y = f(x)$ for $x \in \mathbb{R}$ near a and y close to b .

2. Moreover, the function $f : B(r_0, a) \rightarrow B(r_1, b) \subset \mathbb{R}$ from item 1 is of class C^1 , and its derivatives may be determined by differentiating the identity

$$F(x, f(x)) = 0$$

and solving to find the partial derivatives of f .

Now we turn to our main interest.

Theorem 4. (*Laplace's Method, endpoint contributions*) Let Φ be as in (3.5), with a finite and $g \in L^1(a, b)$. Assume that ϕ has a unique global minimum at $t = a$. In addition, suppose that for some $\delta > 0$ and some $\alpha > -1$, the function g takes the form

$$g(t) = (t - a)^\alpha g_0(t)$$

for $|t - a| < \delta$ and g_0 continuously differentiable in $[a, a + \delta]$ and $g_0(a) \neq 0$, whereas ϕ is twice continuously differentiable on $[a, a + \delta]$ with $\phi'(a) > 0$. Then the function Φ satisfies an estimate of the form

$$\Phi(\lambda) = \frac{1}{\lambda^{\alpha+1}} \frac{g_0(a) \Gamma(\alpha+1)}{\phi'(a)^{\alpha+1}} e^{-\lambda \phi(a)} \left(1 + O(\lambda^{-1})\right)$$

as $\lambda \rightarrow \infty$.

Proof. Start by assuming, without loss of generality, that $a = 0$ and dividing the integral in two intervals, one in the region where we have information about the behavior of g and another in the rest of the interval,

$$\Phi(\lambda) = \int_0^b g(t) e^{-\lambda \phi(t)} dt = \int_0^\delta g(t) e^{-\lambda \phi(t)} dt + \int_\delta^b g(t) e^{-\lambda \phi(t)} dt = (1) + (2).$$

Knowing that ϕ has a global minima at $t = 0$ we assume, changing δ as needed, that $\phi(t) > \phi(\delta)$

for any $t > \delta$. That give us a simple bound for (2):

$$|(2)| \leq \int_{\delta}^b |g(t)| e^{-\lambda\phi(t)} dt \leq e^{-\lambda\phi(\delta)} \int_{\delta}^b |g(t)| dt \leq e^{-\lambda\phi(\delta)} \|g\|_{L^1[0,b]}.$$

We now get back to (1) and start by studying the difference given by $\phi(t) - \phi(0) = s$ in a similar way that we would think of a change of variable from t to s . Define the multivariate function

$$H(t, s) := \phi(t) - \phi(0) - s$$

such that $H(0, 0) = 0$ and $\partial_t H(0, 0) \neq 0$. This means we are in the condition to use [Theorem 3](#) to say that there exists a unique bijective and continuously differentiable function $t(s) := t$ such that

$$\begin{cases} \phi(t(s)) - \phi(0) - s = 0, \\ t'(0) = \frac{\partial_s H(0, 0)}{\partial_t H(0, 0)} = \frac{1}{\phi'(0)}. \end{cases}$$

If we set $T = t^{-1}(\delta)$ we would then have for (1)

$$\int_0^{\delta} g(t) e^{-\lambda\phi(t)} dt = \int_0^T g(t(s)) e^{-\lambda\phi(t(s))} t'(s) ds.$$

Now, to apply the assumption on g we write $g(t(s)) = t^{\alpha}(s)g_0(t(s))$. As t is C^2 and vanishes at the origin, using its Taylor expansion, we have

$$t(s) = st'(0)(1 + r(s)) = \frac{s}{\phi'(0)}(1 + r(s)).$$

Joining both results,

$$\begin{aligned} \int_0^T g(t(s)) e^{-\lambda\phi(t(s))} t'(s) ds &= e^{\lambda\phi(0)} \int_0^T g(t(s)) e^{-\lambda(\phi(t(s)) - \phi(0))} t'(s) ds \\ &= e^{\lambda\phi(0)} \int_0^T t^{\alpha}(s) g_0(t(s)) e^{-\lambda s} t'(s) ds \\ &= \frac{e^{\lambda\phi(0)}}{\phi'(0)^{\alpha}} \int_0^T s^{\alpha} (1 + r(s))^{\alpha} g_0(t(s)) e^{-\lambda s} t'(s) ds \end{aligned}$$

where, if we set $f(s) := s^{\alpha} (1 + r(s))^{\alpha} g_0(t(s))$ we apply Watson's Lemma to finish the result. $\square$

Theorem 5. (*Laplace's Method, interior contributions*) Let Φ be as in (3.5), with a finite and $g \in L^1(a, b)$. Assume that ϕ has a unique global minimum at $t = c \in (a, b)$. In addition, suppose that for some $\delta > 0$ and some $\alpha > -1$, the function g takes the form

$$g(t) = (t - a)^{\alpha} g_0(t)$$

for $|t - a| < \delta$ and g_0 continuously differentiable in $[c - \delta, c + \delta]$ and $g_0(a) \neq 0$, whereas ϕ is three times continuously differentiable on $[c - \delta, c + \delta]$ with $\phi'(a) > 0$. Then the function

Φ satisfies an estimate of the form

$$\Phi(\lambda) = e^{-\lambda\phi(c)} \left(\frac{\sqrt{2\pi}g(c)}{\sqrt{|\phi''(c)|}\lambda^{1/2}} + O(\lambda^{-3/2}) \right)$$

as $\lambda \rightarrow \infty$.

Proof. Although the proof may be similar there is an important distinction in the solution. Suppose we wanted to try the same approach. We would first assume, w.l.g., that $c = 0$. Then, we could write

$$\begin{aligned} \Phi(\lambda) &= \int_a^b g(t) e^{-\lambda\phi(t)} dt \\ &= \int_a^{-\delta} g(t) e^{-\lambda\phi(t)} dt + \int_{-\delta}^{\delta} g(t) e^{-\lambda\phi(t)} dt + \int_{\delta}^b g(t) e^{-\lambda\phi(t)} dt \\ &=: (1) + (2) + (3). \end{aligned}$$

For (1) and (3) the result is as it was before

$$\begin{cases} |(1)| \leq \int_{\delta}^a |g(-t)| e^{-\lambda\phi(-t)} dt \leq e^{\lambda\phi(-\delta)} \|g\|_{L^1[a,b]}, \\ |(3)| \leq \int_{\delta}^b |g(t)| e^{-\lambda\phi(t)} dt \leq e^{\lambda\phi(\delta)} \|g\|_{L^1[a,b]}. \end{cases}$$

For (2) we would try to define a function

$$H(t, s) := \phi(t) - \phi(0) - s^2.$$

However we could not apply [Theorem 3](#) as the partial derivative $\partial_t H = \phi'$ vanishes at $t = 0$. A nice fact about this indetermination is that it does not come from the fact that there are no solutions to $H(t, s) = 0$ but from the fact that there are two. To solve this we will use a technique called *blow up*. First, we introduce a variable $v := v(t)$ such that $vs = t$. Now, we redefine our function of interest to be

$$L(s, v) := \frac{\phi(sv) - \phi(0)}{s^2} - 1.$$

for $s \in (-\delta, \delta) - \{0\}$. We would think this function is also singular but note that, using the Taylor Expansion of ϕ we write

$$\phi(t) = \frac{\phi''(0)}{2} t^2 + t^3 R(t)$$

for some error function R , which leave us with

$$L(s, v) = \frac{\phi''(0)}{2} v^2 + sv^3 R(sv) - 1,$$

and we can indeed apply the Implicit Function Theorem. In fact, we know $L \in C^2$ and as we want to solve for $L(s, v) = 0$, we need that the v_0 we take to satisfy $L(0, v_0) = 0$, ie,

$$v_0 = \sqrt{\frac{2}{\phi''(0)}}.$$

Now, we also know that

$$\partial_v L(0, v_0) = \phi''(0) v_0 \neq 0.$$

So we get a bijective and continuously differentiable function $v = v(s)$ such that

$$\begin{cases} L(s, v(s)) = 0 \\ v(0) = v_0 \end{cases}$$

But, $L(s, v(s))$ means exactly that

$$\phi(t) - \phi(0) = s^2$$

with $t = t(s) = sv(s)$. Getting back to our equation, we have that

$$e^{\lambda\phi(0)} \int_{-\delta}^{\delta} g(t) e^{-\lambda(\phi(t)-\phi(0))} dt = e^{\lambda\phi(0)} \int_{\alpha}^{\beta} g(sv(s)) (v(s) + sv'(s)) e^{-\lambda s^2} ds$$

where we calculate $\alpha = -\sqrt{\phi(-\delta) - \phi(0)}$ and $\beta = \sqrt{\phi(\delta) - \phi(0)}$. One can now again finish the proof with [Proposition 1](#). $\square$

Example 3. (NICA; ORTMANN, 2025) For $x \in \mathbb{R}$ fixed, i.e. $x = O(1)$, we have the following asymptotic as $n \rightarrow \infty$:

$$\text{He}_n(x) = \sqrt{2} \left(\frac{n}{e}\right)^{n/2} e^{x^2/2} \cos\left(\frac{n\pi}{2} - x\sqrt{n}\right) (1 + o(1)).$$

To see that, consider the integral representation of the Hermite Polynomial and re-write

$$\begin{aligned} \text{He}_n(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (x+it)^n e^{-t^2/2} dt = \frac{2}{\sqrt{2\pi}} \Re \left\{ \int_0^{\infty} (x+it)^n e^{-t^2/2} dt \right\} \\ &= \frac{2}{\sqrt{2\pi}} \Re \left\{ \int_{\Re(z)=x, \text{Im}(z)>0} e^{n \ln(z) + \frac{1}{2}(z-x)^2} \frac{dz}{i} \right\} \\ &= \frac{2}{\sqrt{2\pi}} \Re \left\{ \int_{\Re(z)=0, \text{Im}(z)>0} e^{n \ln(z) + \frac{1}{2}(z-x)^2} \frac{dz}{i} \right\} \end{aligned}$$

where you start with a line integral in $\mathbb{C}$ and move the contour over to the positive imaginary axis - justified by the Cauchy's Theorem and the exponential decay of the integrand near infinity. Now, we focus on $n \ln(z) + \frac{1}{2}(z-x)^2$ which has maximum at $z^* = i\sqrt{n} + o(\sqrt{n})$. The idea is to change variables by a factor of $\sqrt{z}$ so that the location of the maximum contribution stays $O(1)$ as n gets large. Specifically, we do

$$z = i\sqrt{n}s, \quad \frac{dz}{i} = \sqrt{n}ds, \quad \ln(z) = \ln(\sqrt{n}) + \ln(s) + i\frac{\pi}{2}$$

which give us

$$\frac{1}{2}(z-x)^2 = -\frac{1}{2}ns^2 + \frac{1}{2}x^2 - i\sqrt{n}sx$$

so that we remain with

$$\begin{aligned} \text{He}_n(x) &= \frac{2}{\sqrt{2\pi}} \Re \left\{ \int_0^{\infty} e^{n \ln(\sqrt{n})} e^{n \ln(s)} e^{ni\frac{\pi}{2}} e^{\frac{1}{2}x^2} e^{-i\sqrt{n}sx} \sqrt{n}ds \right\} \\ &= \frac{2}{\sqrt{2\pi}} \sqrt{n} e^{n \ln(\sqrt{n})} e^{\frac{1}{2}x^2} \int_0^{\infty} e^{n(\ln(s) - \frac{1}{2}s^2)} \cos\left(n\frac{\pi}{2} - \sqrt{n}sx\right) ds, \end{aligned}$$

where we can apply Laplace's approximation discussed before. Noting that $\phi(s) = \ln(s) - \frac{1}{2}s^2$ has $\phi'(s) = \frac{1}{s} - s$ and $\phi''(s) = -\frac{1}{s^2} - 1$, with global maximum at $s^* = 1$ with $\phi(1) = -\frac{1}{2}$ and $\phi''(1) = -2$, yielding

$$\begin{aligned} \text{He}_n(x) &= \frac{2}{\sqrt{2\pi}} \sqrt{n} e^{n \ln(\sqrt{n})} e^{\frac{1}{2}x^2} \left(e^{n\phi(1)} \sqrt{\frac{2\pi}{n|\phi''(1)|}} \cos\left(n\frac{\pi}{2} - \sqrt{n}x\right) (1 + o(1)) \right), \\ &= \sqrt{2} e^{n \ln(\sqrt{n})} e^{\frac{1}{2}(x^2 - n)} \cos\left(n\frac{\pi}{2} - \sqrt{n}x\right) (1 + o(1)), \\ &= \sqrt{2} \left(\frac{n}{e}\right)^{n/2} e^{x^2/2} \cos\left(\frac{n\pi}{2} - x\sqrt{n}\right) (1 + o(1)). \end{aligned}$$

3.2.3 Steepest Descent

Let us now finally explore integrals over the complex plane with complex valued functions. We retake the integral

$$\phi(\lambda) := \int_{\Gamma} g(s) e^{-\lambda f(s)} ds, \quad (3.6)$$

where Γ is a contour in the complex plane and f, g are given complex-valued, analytic functions. Suppose that Γ is parametrized by $\gamma : [a, b] \rightarrow \mathbb{C}$ and let us decompose the functions we have into their real and imaginary parts such as

$$f(z) = u(z) + iv(z), \quad g(z) = p(z) + iq(z), \quad \gamma(t) = \alpha(t) + i\beta(t)$$

with all functions in the decomposition being real valued. Of course we now have

$$e^{-\lambda f(z)} = e^{-\lambda u(z)} e^{-i\lambda v(z)},$$

and when λ grows large, we have the exponentially growing/decaying term $e^{-\lambda u(z)}$ and a oscillating term coming from $e^{i\lambda v(z)}$ which is hard to control. The idea to get around this fact is to use paths in which this term is constant. If we have such a case, we would write

$$\begin{aligned} \Phi(\lambda) &= e^{-i\lambda v_0} \int_a^b (p(\gamma(t))\alpha'(t) - q(\gamma(t))\beta'(t)) e^{-\lambda u(\gamma(t))} dt \\ &\quad + i e^{-i\lambda v_0} \int_a^b (p(\gamma(t))\beta'(t) - q(\gamma(t))\alpha'(t)) e^{-\lambda u(\gamma(t))} dt \quad (3.7) \end{aligned}$$

where each integral can now be solved by Laplace's method.

Theorem 6. (Steepest Descent) Consider Φ as in (3.6) with a smooth contour Γ . Assume that the exponent $f = u + iv$ is an analytic function on a neighborhood of Γ that satisfies

$$v(z) \equiv v_0, \quad z \in \Gamma$$

In addition, suppose that u has a unique minimum at an interior point of Γ , say on $z_0 \in \Gamma$,

with $f''(z_0) \neq 0$, and also that g is analytic on a neighborhood of Γ and satisfies

$$\int_{\Gamma} |g(z)| |dz| < \infty.$$

Then Φ has a leading asymptotic as $\lambda \rightarrow +\infty$ given by

$$\Phi(\lambda) = e^{-\lambda f(z_0)} \left(\frac{\sqrt{2\pi} g(z_0) e^{i\theta(z_0)}}{\lambda^{1/2} \sqrt{|f''(z_0)|}} + O(\lambda^{-3/2}) \right), \quad \lambda \rightarrow +\infty,$$

where $\theta(z_0)$ is the angle of the oriented tangent of Γ at z_0 .

Proof. Fix a parametrization $\gamma : [a, b] \rightarrow \mathbb{C}$ of Γ with $\gamma(c) = z_0, a < c < b$, and decompose Φ as before in (3.7). By the assumptions on the functions f, g we apply Laplace's method on both equations, and yields, as $\lambda \rightarrow +\infty$,

$$\int_a^b (p(\gamma(t))\alpha'(t) - q(\gamma(t))\beta'(t)) e^{-\lambda u(\gamma(t))} dt = \sqrt{2\pi} e^{-\lambda u(z_0)} \left(\frac{p(z_0)\alpha'(c) - q(z_0)\beta'(c)}{\sqrt{|(u \circ \gamma)''(c)|} \lambda^{1/2}} + O(\lambda^{3/2}) \right)$$

and

$$\int_a^b (p(\gamma(t))\beta'(t) - q(\gamma(t))\alpha'(t)) e^{-\lambda u(\gamma(t))} dt = \sqrt{2\pi} e^{-\lambda u(z_0)} \left(\frac{p(z_0)\beta'(c) - q(z_0)\alpha'(c)}{\sqrt{|(u \circ \gamma)''(c)|} \lambda^{1/2}} + O(\lambda^{3/2}) \right)$$

Using that $v_0 + u(z_0) = f(z_0)$ and also that,

$$(p(z_0)\alpha'(c) - q(z_0)\beta'(c)) + i(p(z_0)\beta'(c) - q(z_0)\alpha'(c)) = g(z_0)\gamma'(c),$$

and, combining the estimates for both parts we get

$$\Phi(\lambda) = \sqrt{2\pi} e^{-\lambda f(z_0)} \left(\frac{g(z_0)\gamma'(c)}{\sqrt{|(u \circ \gamma)''(c)|} \lambda^{1/2}} + O(\lambda^{3/2}) \right)$$

and it suffices now to get an expression for the second derivative in the denominator. Knowing the imaginary part is constant we write

$$\frac{d}{dt} u(\gamma(t)) = \frac{d}{dt} f(\gamma(t)) = \gamma'(t) f'(\gamma(t))$$

and

$$\frac{d^2}{dt^2} u(\gamma(t)) = \frac{d^2}{dt^2} f(\gamma(t)) = (\gamma'(t))^2 f''(\gamma(t)) + \gamma''(t) f'(\gamma(t)).$$

Hence,

$$\frac{\gamma'(c)}{\sqrt{|(u \circ \gamma)''(c)|}} = \frac{1}{\sqrt{|f''(z_0)|}} \frac{\gamma'(c)}{|\gamma'(c)|}$$

which concludes the proof. $\square$

We now need to discuss why this assumption of constant value of $\text{Im}(f(z))$ is not an assumption as strong as one might assume at first. Suppose we want to integrate in an arbitrary contour Γ of the plane. Of course, our assumption that the functions were analytic permit us to deform the contour keeping the value of the integral. We are going to use this to find a contour $\tilde{\Gamma}$ such that sections of the contour are such that $\text{Im}(f(z))$ is constant. We already know that being analytic means satisfying the *Cauchy-Riemann* equations,

$$u_x(z) = v_y(z), \quad \text{and} \quad u_y(z) = -v_x(z).$$

An immediate result from these equations is that the gradients of u and v must be normal. Indeed, if $\langle \cdot, \cdot \rangle$ is the inner product on $\mathbb{R}^2$,

$$\langle \nabla u(z), \nabla v(z) \rangle = u_x(z)v_x(z) + u_y(z)v_y(z) = u_x(z)v_x(z) - v_x(z)u_x(z) = 0.$$

As we are following the curves where $\text{Im}(f(z))$ is constant, we are also following the path of Steepest Descent of the real part of the integral. As we also are looking to use Laplace's method on the integrals, we may want to have our paths going through the critical points of the function. This fully motivates the method described.

Example 4. There are three asymptotics regimes for $\text{He}_n(x)$ when $x \rightarrow \infty$ depending on the value of $\lim_{N \rightarrow \infty} x/2\sqrt{N}$ for $N = n + 1$, each of which has qualitatively different behavior

- Small x values $-2\sqrt{N} < x < 2\sqrt{N}$: in this regime the Hermite Polynomials will be oscillating, and have a similar qualitative behavior as in [Example 3](#), with modification to phase and amplitude.
- Large x values $|x| > 2\sqrt{N}$: In this regime the polynomial does not have more oscillations and is monotone increasing. As x grows larger, this approximation behaves as $x^N e^{-N^2/2x^2}$.
- Edge x values $x = \pm 2\sqrt{N} + O(N^{-\frac{1}{6}})$: in this regime the polynomials will be approximated by the Airy Function. This interpolates between the oscillatory small x regime and the monotone large x regimes.

We write the integral representation given by:

$$\begin{aligned} \text{He}_n(x) &:= \frac{n!}{2\pi i} \int_{\Gamma} z^{-n-1} e^{xz - \frac{z^2}{2}} dz \\ &= \frac{n!}{2\pi i} \int_{\Gamma} e^{xz - \frac{z^2}{2} - (n+1)\ln(z)} dz \end{aligned}$$

where Γ is a contour that goes around the origin coming and going from minus infinity such as $\Gamma = \{\lambda - i\delta : \lambda < 0\} \cup \{\delta e^{i\theta} : -\pi/2 < \theta < \pi/2\} \cup \{\lambda + i\delta : \lambda < 0\}$. This is equivalent to using $\Gamma = \{e^{i\theta} : 0 \leq \theta < 2\pi\}$. To determine He_n , set $N = n + 1$ and consider the scaling

$x \mapsto N^a \alpha$ and $z = N^b \beta$

$$\begin{aligned} \text{He}_n(N^a \alpha) &= \frac{(N-1)!}{2\pi i} \int_{\Gamma} e^{(N^a \alpha)(N^b \beta) - \frac{1}{2}(N^b \beta)^2 - N \ln(N^b \beta)} d\beta \\ &\stackrel{(a=b=1/2)}{=} \frac{(N-1)! N^{\frac{1}{2}(1-N)}}{2\pi i} \int_{\Gamma} e^{-N(-\alpha \beta + \frac{1}{2}\beta^2 + \ln(\beta))} d\beta, \end{aligned}$$

so that we have

$$\text{He}_n(\sqrt{N}\alpha) = C_n^0 \int_{\Gamma} e^{-N\phi_{\alpha}(\beta)} d\beta$$

with

$$\phi_{\alpha}(\beta) = \frac{1}{2}\beta^2 - \alpha\beta + \ln(\beta).$$

As we want to calculate the asymptotic by steepest descent, we need to determine the critical points of such a function and then the steepest descent paths passing through this points. We can easily determine the critical points as $\beta_c^{\pm} = \frac{\alpha \pm \sqrt{\alpha^2 - 4}}{2}$ - solutions to the quadratic formula given by solving the critical point of the function (the zeros of its derivatives). Now, the behavior of the critical points will depend on the interval of the real parameter α ,

$$\beta_c^{\pm} = \frac{\alpha \pm \sqrt{\alpha^2 - 4}}{2} = \begin{cases} e^{\pm\gamma}, & \alpha = 2 \cosh(\gamma) > 2, \quad \gamma > 0; \\ e^{\pm i\gamma}, & \alpha = 2 \cos(\gamma) \in [0, 2), \quad \gamma \in (0, \pi/2]; \\ 1, & \alpha = 2, \end{cases}$$

each of the cases refers to one of the regimes we described before.




Figure 2 – The lines represent the solution to $\text{Im}[f(z) - f(t_c)] = 0$ for the three possible cases of the Hermite Polynomial. These are the curves we use to perform the asymptotic analysis on the integral. Note the case $\alpha = 2$; the coalescence of critical point gives origin to the Airy function.

EQUILIBRIUM

PLANAR ENSEMBLES

BIBLIOGRAPHY

ABUL-MAGD, A. Modelling gap-size distribution of parked cars using random-matrix theory. **Physica A: Statistical Mechanics and its Applications**, Elsevier BV, v. 368, n. 2, p. 536–540, Aug. 2006. ISSN 0378-4371. Available: <http://dx.doi.org/10.1016/j.physa.2005.10.059>. No citation.

ALPERT, B. K. Hybrid gauss-trapezoidal quadrature rules. **SIAM J. Sci. Comput.**, Society for Industrial and Applied Mathematics, USA, v. 20, n. 5, p. 1551–1584, Jan. 1999. ISSN 1064-8275. Available: <https://doi.org/10.1137/S1064827597325141>. Citations on pages 23 and 25.

BAIK, J.; DEIFT, P.; JOHANSSON, K. **On the Distribution of the Length of the Longest Increasing Subsequence of Random Permutations**. 1999. Available: <https://arxiv.org/abs/math/9810105>. No citation.

BALOGH, F.; GRAVA, T.; MERZI, D. Orthogonal polynomials for a class of measures with discrete rotational symmetries in the complex plane. 2016. No citation.

BYUN, S.-S.; KANG, N.-G.; SEO, S.-M. Partition functions of determinantal and pfaffian coulomb gases with radially symmetric potentials. **Communications in Mathematical Physics**, Springer Science and Business Media LLC, v. 401, n. 2, p. 1627–1663, Mar. 2023. ISSN 1432-0916. Available: <http://dx.doi.org/10.1007/s00220-023-04673-1>. No citation.

CHAFĀĪ, D.; FERRÉ, G. Simulating coulomb and log-gases with hybrid monte carlo algorithms. **Journal of Statistical Physics**, Springer Science and Business Media LLC, v. 174, n. 3, p. 692–714, Nov. 2018. ISSN 1572-9613. Available: <http://dx.doi.org/10.1007/s10955-018-2195-6>. No citation.

DEIFT, P. **Orthogonal Polynomials and Random Matrices: A Riemann-Hilbert Approach**. Courant Institute of Mathematical Sciences, New York University, 1999. (Courant lecture notes in mathematics). ISBN 9780821883440. Available: <https://books.google.com.br/books?id=SBR8yv0LkFgC>. No citation.

_____. **Universality for mathematical and physical systems**. 2006. Available: <https://arxiv.org/abs/math-ph/0603038>. No citation.

DYSON, F. J. Statistical theory of the energy levels of complex systems. I. **J. Math. Phys.**, v. 3, p. 140–156, 1962. No citation.

EDELMAN, A. **Eigenvalues and condition numbers of random matrices**. Phd Thesis (PhD Thesis) — Massachusetts Institute of Technology, Dept. of Mathematics, 1989. Unpublished. Available: <http://hdl.handle.net/1721.1/14322>. No citation.

GAUSS, C. **Theoria motus corporum coelestium in sectionibus conicis solem ambientium**. Sumtibus F. Perthes et I.H. Besser, 1809. (Carl Friedrich Gauss Werke). Available: <https://books.google.com.br/books?id=ORUOAAAAQAAJ>. No citation.

GINIBRE, J. Statistical ensembles of complex, quaternion, and real matrices. **Journal of Mathematical Physics**, v. 6, n. 3, p. 440–449, 03 1965. ISSN 0022-2488. Available: <https://doi.org/10.1063/1.1704292>. No citation.

GIROTTI, M. **Riemann-Hilbert approach to Gap Probabilities of Determinantal Point Processes**. Phd Thesis (PhD Thesis) — Concordia University, August 2014. Unpublished. Available: <https://spectrum.library.concordia.ca/id/eprint/978910/>. No citation.

GOLDSTON, J.; MONTGOMERY, G. Pair correlation of zeros and primes in short intervals. In: ADOLPHSON, A. C.; CONREY J. B. AND GHOSH, A.; YAGER, R. I. (Ed.). **Analytic Number Theory and Diophantine Problems: Proceedings of a Conference at Oklahoma State University, 1984**. Boston, MA: Birkhauser Boston, 1987. p. 183–203. ISBN 978-1-4612-4816-3. Available: https://doi.org/10.1007/978-1-4612-4816-3_10. No citation.

HAMMERSLEY, J. M. A few seedlings of research. In: . [s.n.], 1972. Available: <https://api.semanticscholar.org/CorpusID:36895634>. No citation.

HEDENMALM, H.; WENNMAN, A. **Planar orthogonal polynomials and boundary universality in the random normal matrix model**. 2020. Available: <https://arxiv.org/abs/1710.06493>. No citation.

JOHANSSON, K. Course 1 - random matrices and determinantal processes. In: BOVIER, A.; DUNLOP, F.; van Enter, A.; den Hollander, F.; DALIBARD, J. (Ed.). Elsevier, 2006, (Les Houches, v. 83). p. 1–56. Available: <https://www.sciencedirect.com/science/article/pii/S0924809906800387>. No citation.

KIEBURG, M.; VERBAARSCHOT, J. J. M.; ZAFEIROPOULOS, S. Dirac spectra of two-dimensional qcd-like theories. **Physical Review D**, American Physical Society (APS), v. 90, n. 8, Oct. 2014. ISSN 1550-2368. Available: <http://dx.doi.org/10.1103/PhysRevD.90.085013>. No citation.

KRBÁLEK, M.; SEBA, P. The statistical properties of the city transport in cuernavaca (mexico) and random matrix ensembles. **Journal of Physics A: Mathematical and General**, IOP Publishing, v. 33, n. 26, p. L229–L234, 2000. ISSN 1361-6447. Available: <http://dx.doi.org/10.1088/0305-4470/33/26/102>. No citation.

KUIJLAARS, A. **Riemann-Hilbert Problems and Multiple Orthogonal Polynomials**. 2014. <http://wis.kuleuven.be/analyse/arno>. [Online; accessed 08-April-2026]. No citation.

LEBLÈ, T.; SERFATY, S. Large deviation principle for empirical fields of log and riesz gases. **Invent. Math.**, Springer Science and Business Media LLC, v. 210, n. 3, p. 645–757, dec 2017. No citation.

LIVAN, G.; NOVAES, M.; VIVO, P. **Introduction to Random Matrices**. Springer International Publishing, 2018. ISSN 2197-1765. ISBN 9783319708850. Available: <http://dx.doi.org/10.1007/978-3-319-70885-0>. No citation.

LOGAN, B.; SHEPP, L. A variational problem for random young tableaux. **Advances in Mathematics**, v. 26, n. 2, p. 206–222, 1977. ISSN 0001-8708. Available: <https://www.sciencedirect.com/science/article/pii/0001870877900305>. No citation.

MAJUMDAR, S. N. **Random Matrices, the Ulam Problem, Directed Polymers, Growth Models, and Sequence Matching**. 2007. Available: <https://arxiv.org/abs/cond-mat/0701193>. No citation.

MAJUMDAR, S. N.; SCHEHR, G. Top eigenvalue of a random matrix: large deviations and third order phase transition. **Journal of Statistical Mechanics: Theory and Experiment**, IOP Publishing, v. 2014, n. 1, p. P01012, Jan. 2014. ISSN 1742-5468. Available: <http://dx.doi.org/10.1088/1742-5468/2014/01/P01012>. No citation.

METROPOLIS, N. The beginning of the monte carlo method. In: _____. [S.l.]: Los Alamos Science Special Issue, 1987. v. 15, p. 125–130. No citation.

NICA, M.; ORTMANN, J. **A Gaussian integral formula for the Hermite polynomials: Combinatorics, Asymptotics and Applications**. 2025. Available: <https://arxiv.org/abs/2508.13910>. Citation on page 35.

ODLYZKO, A. M.; RAINS, E. M. On longest increasing subsequences in random permutations. In: **Analysis, Geometry, Number Theory: The Mathematics of Leon Ehrenpreis**. American Mathematical Society, 2000. p. 439–451. Available: <https://doi.org/10.1090/conm/251/03886>. No citation.

OLVER, F.; LOZIER, D.; BOISVERT, R.; CLARK, C. Nist handbook of mathematical functions. 01 2010. Citations on pages 23 and 24.

PIMENTA, J. V. A.; SILVA, G. L. F. **Matrizes aleatórias e simulação de Gases de Coulomb**. 2024. No citation.

POZNIAK, M.; ZYCZKOWSKI, K.; KUS, M. Composed ensembles of random unitary matrices. **Journal of Physics A: Mathematical and General**, v. 31, n. 3, p. 1059, jan 1998. Available: <https://doi.org/10.1088/0305-4470/31/3/016>. No citation.

RAHMAN, A. A. **Recursive characterisations of random matrix ensembles and associated combinatorial objects**. 2023. Available: <https://arxiv.org/abs/2301.11428>. No citation.

SAFF, E.; TOTIK, V. **Logarithmic Potentials with External Fields**. Springer-Verlag, 2024. (Grundlehren der mathematischen Wissenschaften). ISBN 9783031651335. Available: <https://books.google.com.bo/books?id=z2wnEQAAQBAJ>. No citation.

SERFOZO, R. F. Chapter 1 point processes. In: **Stochastic Models**. Elsevier, 1990, (Handbooks in Operations Research and Management Science, v. 2). p. 1–93. Available: <https://www.sciencedirect.com/science/article/pii/S0927050705801653>. No citation.

SILVA, G. L. F. **Lecture Notes on Riemann-Hilbert Problems**. 2021. <https://sites.google.com/site/guilhermesilvamath/publications>. [Online; accessed 20-April-2026]. Citation on page 23.

TREMBLAY, N. **A short introduction to Determinantal Point Processes**. [S.l.]: Institut Henri Poincaré, 2025. No citation.

ULAM, S. M. On some statistical properties of dynamical systems. In: NEYMAN, J. (Ed.). **Fourth Berkeley Symposium on Mathematical Statistics and Probability**. [S.l.: s.n.], 1961. p. 315–320. No citation.

VERSHIK, S. V. K. A. M. Asymptotics of the plancherel measure of the symmetric group and the limiting form of young tableaux. **Dokl. Akad. Nauk SSSR**, v. 233, n. 6, p. 1024–1027, 1977. No citation.

WIGNER, E. P. On the distribution of the roots of certain symmetric matrices. **Annals of Mathematics**, v. 67, n. 2, p. 325, mar 1958. No citation.

_____. The unreasonable effectiveness of mathematics in the natural sciences. In: _____. **Philosophical Reflections and Syntheses**. Berlin, Heidelberg: Springer Berlin Heidelberg, 1995. p. 534–549. ISBN 978-3-642-78374-6. Available: <https://doi.org/10.1007/978-3-642-78374-6_41>. No citation.

