

U:%%- *GNU Emacs 27.2 (Build 20D91) of 2021-03-27 Copyright (C) 2021

For information

Welcome to GNU Emacs. To follow a link, type C-h C-l. To quit a partial match, type C-g.

Important Help menu items: Emacs Tutorial Read the Emacs Manual (Non)Warranty Copying Conditions More Manuals / Other

Useful tasks: Visit New File Open Home Directory Customize Start-up Options

10.0.0.74:3000

Not Secure

Apps Gmail YouTube Maps Translate News

Reading List

Papers

Latest Papers

Genetic dissection of assortative mating behavior

The evolution of new species is made easier when traits under divergent ecological selection are also mating cues. Such ecological mating cues are now considered more common than previously thought, but we still know little about the genetic changes underlying their evolution, or more generally about the genetic basis for assortative mating behaviors. The warning patterns of *Heliconius melpomene* and *H. cydno* are under disruptive selection due to increased predation of non-mimetic hybrids, and are used during mate recognition. We carried out a genome-wide quantitative trait locus (QTL) analysis of preference behaviors between these species and showed that divergent male preference has a simple genetic basis. Three QTLs each explain a large proportion of the differences in preference behavior observed between the parental species. Two of these QTLs are on chromosomes with major color pattern genes, including one that is tightly associated with the gene *optix*. Different loci influence different aspects of attraction, suggesting that behavioral isolation in *Heliconius* involves the evolution of independently segregating modules, similar to those for the corresponding wing pattern cues. Hybridization and