

Abstract Algebra Homework 11

Joe Loser

April 30, 2016

This problem set includes problems 3c, 4b, 24, and an extra problem from section 17.4.

3) Use the division algorithm to find $q(x)$ and $r(x)$ such that $a(x) = q(x)b(x) + r(x)$ with $\deg r(x) < \deg b(x)$.

3c) $a(x) = 4x^5 - x^3 + x^2 + 4$ and $b(x) = x^3 - 2$ where $a(x), b(x) \in \mathbb{Z}_5[x]$.

Solution: First note that in $\mathbb{Z}_5[x]$, $a(x) \equiv 4x^5 + 4x^3 + x^2 + 4$ and $b(x) \equiv x^3 + 3$.

Now performing long division, we have

$$\begin{array}{r} \overline{4x^2+4} \\ x^3+3 \overline{) 4x^5+4x^3+x^2+4} \\ \underline{-4x^5} -12x^2 \\ 4x^3-11x^2+4 \\ \underline{-4x^3} -12 \\ -11x^2-8 \end{array}$$

Thus,

$$\begin{aligned} a(x) &= (4x^2 + 4) \cdot (x^3 + 3) + (-11x^2 - 8) \\ &\equiv (4x^2 + 4) \cdot (x^3 + 3) + (4x^2 + 2). \end{aligned}$$

□

4) Find the greatest common divisor of each of the following pairs $p(x)$ and $q(x)$ of polynomials. If $d(x) = \gcd(p(x), q(x))$, find two polynomials $a(x)$ and $b(x)$ such that $d(x) = a(x)p(x) + b(x)q(x)$.

4b) $p(x) = x^3 + x^2 - x + 1$ and $q(x) = x^3 + x - 1$ where $p(x), q(x) \in \mathbb{Z}_2[x]$.

First note that $\mathbb{Z}_2[x]$, $p(x) \equiv x^3 + x^2 + x + 1$ and $q(x) \equiv x^3 + x + 1$.

Performing the first stage of long division, we have that

$$\begin{array}{r} \overline{1} \\ x^3+x^2+x+1 \overline{) x^3+x+1} \\ \underline{-x^3-x^2-x-1} \\ -x^2 \end{array}$$

Note that in $\mathbb{Z}_2[x]$, $-x^2 \equiv x^2$ and $2x^2 \equiv 0$. So

$$x^3 + x + 1 = 1 \cdot (x^3 + x^2 + x + 1) + (x^2). \quad (1)$$

In the second stage of long division, we get

$$\begin{array}{r} \overline{x+1} \\ x^2 \overline{) x^3+x^2+x+1} \\ \underline{-x^3} x^2 \\ -x^2 \end{array}$$

So we have that

$$x^3 + x^2 + x + 1 = (x + 1) \cdot (x^2) + (x + 1). \quad (2)$$

In the third stage, we get

$$\begin{array}{r} x+1 \overline{) \begin{array}{r} x^2 \\ -x^2-x \\ \hline -x \\ x+1 \\ \hline 1 \end{array}} \end{array}$$

Then

$$x^2 = (x + 1) \cdot (x + 1) + (1). \quad (3)$$

In the last stage of long division, we have

$$\begin{array}{r} x+1 \overline{) \begin{array}{r} x+1 \\ -x \\ \hline 1 \\ -1 \\ \hline 0 \end{array}} \end{array}$$

So

$$x + 1 = (1) \cdot (x + 1) + 0. \quad (4)$$

Thus $\gcd(p(x), q(x)) = 1$. Performing the back substitution, we have the following

$$\begin{aligned} 1 &= x^2 - (x + 1)(x + 1) \\ &= x^2 - (x + 1)[(x^3 + x^2 + x + 1) - x^2(x + 1)] \\ &= x^2 + (x + 1)(x^3 + x^2 + x + 1) + x^2(x + 1)^2 \\ &= x^2(1 + (x + 1)^2) + (x + 1)(x^3 + x^2 + x + 1) \\ &= [x^3 + x + 1 + (x^3 + x^2 + x + 1)](x^2) + (x + 1)(x^3 + x^2 + x + 1) \\ &= (x^3 + x^2 + x + 1)(x^2 + x + 1) + (x^2)(x^3 + x + 1). \end{aligned}$$

Thus

$$\begin{aligned} 1 &= (x^2 + x + 1)(x^3 + x^2 + x + 1) + (x^2)(x^3 + x + 1) \\ &= a(x)p(x) + b(x)q(x). \end{aligned}$$

To check this holds in $\mathbb{Z}_2[x]$, we multiply everything out. We get that

$$\begin{aligned} 1 &= (x^2 + x + 1)(x^3 + x^2 + x + 1) + (x^2)(x^3 + x + 1) \\ &= x^5 + x^4 + x^3 + x^2 + x^4 + x^3 + x^2 + x + x^3 + x^2 + x + 1 + x^5 + x^3 + x^2 \\ &= 2x^5 + 2x^4 + 4x^3 + 4x^2 + 2x + 1 \\ &\equiv 1. \end{aligned}$$

□

24) Show that $x^p - x$ has p distinct zeros in \mathbb{Z}_p for any prime p . Conclude that

$$x^p - x = x(x-1)(x-2)\cdots(x-(p-1)).$$

Proof: By Fermat's Little Theorem, for all $a \in \mathbb{Z}_p$ we have that $a^p = a$. So $a^p - a = 0$. Thus every $a \in \mathbb{Z}_p$ is a zero of the polynomial $x^p - x$. Note that the polynomial has degree p and p zeros in \mathbb{Z}_p . The numbers $0, 1, \dots, p-1$ are the roots of the equation $x^p - x$, i.e. the p distinct roots. Hence it must split into p distinct linear factors in $\mathbb{Z}_p[x]$ as follows:

$$x^p - x = x(x-1)(x-2)\cdots(x-(p-1)).$$

□

E1) Construct a field with 8 elements.

Solution: Since $8 = 2^3$ we start with a field \mathbb{Z}_2 of characteristic 2 and look for an irreducible polynomial of degree 3 in $\mathbb{Z}_2[x]$. Such a polynomial is $p(x) = x^3 + x + 1$.

We will show that

$$K := \frac{\mathbb{Z}_2[x]}{\langle x^3 + x + 1 \rangle}$$

is a field of 8 elements.

To see why $p(x)$ is irreducible in $\mathbb{Z}_2[x]$, since it of degree 3 or lower, we can look at all of the roots in \mathbb{Z}_2 since $g(x)$ is irreducible if and only if p does not have a root, i.e. $p(a) \neq 0$ for all $a \in \mathbb{Z}_2$. We have that $g(0) = 1$ and $g(1) = 3 \equiv 1$. So neither 0 or 1 are a root of $p(x)$. Hence we see that $p(x)$ is irreducible over $\mathbb{Z}_2[x]$.

By the Division Algorithm, we have that

$$p(x) + \langle x^3 + x + 1 \rangle = a_0 + a_1x + a_2x^2 + \langle x^3 + x + 1 \rangle$$

where $a_0, a_1, a_2 \in \mathbb{Z}_2$. So

$$K = \{a_0 + a_1x + a_2x^2 + \langle x^3 + x + 1 \rangle \mid a_0, a_1, a_2 \in \mathbb{Z}_2\}.$$

Note that since $a_0, a_1, a_2 \in \mathbb{Z}_2$ we see that the order of K which we denote $|K|$ is

$$\begin{aligned} |K| &\leq |\mathbb{Z}_2| \times |\mathbb{Z}_2| \times |\mathbb{Z}_2| \\ &= 2 \times 2 \times 2 \\ &= 8. \end{aligned}$$

To see why K has exactly 8 elements, suppose $a_0 + a_1x + a_2x^2 + \langle x^3 + x + 1 \rangle = b_0 + b_1x + b_2x^2 + \langle x^3 + x + 1 \rangle$. Then

$$(a_0 - b_0) + (a_1 - b_1)x + (a_2 - b_2)x^2 \in \langle x^3 + x + 1 \rangle.$$

That is,

$$\underbrace{x^3 + x + 1}_{\deg 3} \mid \underbrace{(a_0 - b_0) + (a_1 - b_1)x + (a_2 - b_2)x^2}_{\deg < 3}.$$

So $(a_0 - b_0) + (a_1 - b_1)x + (a_2 - b_2)x^2 = 0$. Hence $a_0 = b_0, a_1 = b_1$ and $a_2 = b_2$. Therefore $|K| = 8$.

To explicitly see what the elements of K are, let $\alpha := x + \langle x^3 + x + 1 \rangle \in K$. Now we want to compute the powers of α for $i = 1 \cdots 7$. This will give us the elements of K .

$$\begin{aligned} \alpha^2 &= (x + \langle x^3 + x + 1 \rangle)^2 \\ &= x^2 + 2x\langle x^3 + x + 1 \rangle + (\langle x^3 + x + 1 \rangle)^2 \\ &\equiv x^2 + \langle x^3 + x + 1 \rangle. \end{aligned}$$

In K , $x^3 + x + 1 = 0 \iff \alpha^3 + \alpha + 1 = 0 \iff \alpha^3 = \alpha + 1$. That is, $\alpha^3 = x + 1 + \langle x^3 + x + 1 \rangle$. Next we see that

$$\begin{aligned} \alpha^4 &= \alpha^3 \cdot \alpha \\ &= (\alpha + 1) \cdot \alpha \\ &= (x + 1) \cdot x \\ &\equiv x^2 + x + \langle x^3 + x + 1 \rangle. \end{aligned}$$

Also

$$\begin{aligned}\alpha^5 &= \alpha^3 \cdot \alpha^2 \\ &= (\alpha + 1) \cdot \alpha^2 \\ &= \alpha^3 + \alpha^2 \\ &= (\alpha + 1) + \alpha^2 \\ &\equiv x + 1 + x^2 \\ &\equiv x^2 + x + 1 + \langle x^3 + x + 1 \rangle.\end{aligned}$$

Continuing on, we have that

$$\begin{aligned}\alpha^6 &= \alpha^3 \cdot \alpha^3 \\ &= (x + 1)(x + 1) \\ &= x^2 + 2x + 1 \\ &\equiv x^2 + 1 + \langle x^3 + x + 1 \rangle.\end{aligned}$$

Lastly we have that

$$\begin{aligned}\alpha^7 &= \alpha^4 \cdot \alpha^3 \\ &= (\alpha^2 + \alpha) \cdot (\alpha + 1) \\ &= \alpha^3 + \alpha^2 + \alpha^2 + \alpha \\ &= \alpha^3 + \alpha \\ &= (\alpha + 1) + \alpha \\ &\equiv 1.\end{aligned}$$

Explicitly listing the elements of K , we have that

$$\begin{aligned}K &= \{0, x, x^2, x + 1, x^2 + x, x^2 + x + 1, x^2 + 1, 1\} \\ &= \{0, 1, x, x + 1, x^2, x^2 + 1, x^2 + x, x^2 + x + 1\}.\end{aligned}$$

□