

Abstract Algebra Homework 7

Joe Loser

March 18, 2016

This problem set includes problems 10.3 numbers 4d), 11.3 numbers 7, 16, 17 and 11.4 number 5.

4) Let T be the group of nonsingular upper triangular 2×2 matrices with entries in \mathbb{R} . Let U consist of matrices of the form

$$\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$$

where $x \in \mathbb{R}$.

4d) Show that T/U is abelian.

Proof: Note that we have already showed that U is normal in T in part 4c). To show that T/U is abelian, we need to show that $(AU)(BU) = (BU)(AU)$ for all $A, B \in T$.

Let $A = \begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$ and let $B = \begin{pmatrix} a' & b' \\ 0 & c' \end{pmatrix}$. Then we have that

$$AB = \begin{pmatrix} aa' & ab' + bc' \\ 0 & cc' \end{pmatrix}$$

and

$$BA = \begin{pmatrix} a'a & a'b + b'c \\ 0 & c'c \end{pmatrix}.$$

This shows that $AB \neq BA$ in general. However, we want to show that $(AU)(BU) = (BU)(AU)$. Note that $(AU)(BU) = ABU$ and $(BU)(AU) = BAU$ since U is normal. Let $C = \begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix} \in U$ where $z \in \mathbb{R}$. Then we have that

$$ABU = \begin{pmatrix} aa' & z(ab' + bc') \\ 0 & cc' \end{pmatrix}$$

and

$$BAU = \begin{pmatrix} a'a & z(a'b + b'c) \\ 0 & c'c \end{pmatrix}.$$

Notice that $aa' = a'a$ and $cc' = c'c$ since $a, a', c, c' \in \mathbb{R}$. So we see that ABU and BAU only differ in the upper right entry. This does not matter though since U is matrices of the form

$$\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$$

where $x \in \mathbb{R}$. Notice that both $z(ab' + bc')$ and $z(a'b + b'c) \in \mathbb{R}$. Thus, AB and BA define the same coset in U , meaning that $ABU = BAU$. Thus T/U is abelian. \square

7) In the group \mathbb{Z}_{24} , let $H = \langle 4 \rangle$ and $N = \langle 6 \rangle$.

a) List the elements in $H + N$ and $H \cap N$.

Solution: We have that

$$\begin{aligned} H + N &= \{h + n \mid h \in H \text{ and } n \in N\} \\ &= \{h + n \mid h \in \langle 4 \rangle \text{ and } n \in \langle 6 \rangle\} \\ &= \{0, 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22\} \\ &= \langle 2 \rangle \end{aligned}$$

We also see that $H \cap N = \{0, 12\}$.

Remarks: Note that $H + N = \langle \gcd(4, 6) \rangle$ in \mathbb{Z}_{24} and also that $H \cap N = \langle \text{lcm}(4, 6) \rangle \mathbb{Z}$ in \mathbb{Z}_{24} . □

b) List the cosets in HN/N , showing the elements in each coset.

Solution: We know that

$$(H + N)/N := \{g + n \mid g \in H + N\}.$$

So we see that the cosets that partition $H + N$ are the following:

$$\begin{aligned} 0 + N &= \{0, 6, 12, 18\} \\ 2 + N &= \{2, 8, 14, 20\} \\ 4 + N &= \{4, 10, 16, 22\} \end{aligned}$$

c) List the cosets in $H/(H \cap N)$, showing the elements in each coset.

Solution: We know that

$$H/(H \cap N) := \{aH \cap N \mid a \in H\}.$$

So we see that the cosets that partition the group H are the following:

$$\begin{aligned} 0 + H \cap N &= \{0, 12\} \\ 4 + H \cap N &= \{4, 16\} \\ 8 + H \cap N &= \{8, 20\} \end{aligned}$$

d) Give the correspondence between $(H + N)/N$ and $H/(H \cap N)$ described in the proof of the Second Isomorphism Theorem.

Solution: Recall that all subgroups of an abelian group are normal. Since \mathbb{Z}_{24} is an abelian group, we have that H and N are normal in \mathbb{Z}_{24} . So we can apply the Second Isomorphism Theorem which tells us

$$H/(H \cap N) \cong (H + N)/N.$$

□

16) If H and K are normal subgroups of G and $H \cap K = \{e\}$, prove that G is isomorphic to a subgroup of $G/H \times G/K$.

Proof: To show that G is isomorphic to a subgroup of $G/H \times G/K$, we need to define a function ϕ and show it is a group homomorphism. Then we will show that $\ker \phi = \{e\}$ and use the First Isomorphism Theorem.

Let $\phi : G \rightarrow G/H \times G/K$ be defined as $\phi(g) = (gH, gK)$. Clearly the function ϕ is well-defined. We now need to show this is indeed a group homomorphism. Let $a, b \in G$. Then

$$\begin{aligned} \phi(ab) &= (abH, abK) \\ &= (aH, aK)(bH, bK) \\ &= \phi(a)\phi(b). \end{aligned}$$

So ϕ is a homomorphism. From the First Isomorphism Theorem, we know that

$$G/\ker \phi \cong \phi(G).$$

Since $\phi(G)$ is a subgroup of $G/H \times G/K$ and we want to show $G \cong \phi(G)$, it suffices for us to show that $\ker \phi = \{e\}$, i.e. $\ker \phi = H \cap K$.

Let $g \in \ker \phi$. Then $\phi(g) = (gH, gK) = (H, K)$. That is, $gH = H$ and $gK = K$. Thus $g \in H \cap K$ and we have that $\ker \phi \subset H \cap K$. Contrarily, if $g \in H \cap K$ then we clearly see that $\phi(g) = (gH, gK) = (H, K)$ so that $H \cap K \subset \ker \phi$. Thus $\ker \phi = H \cap K$.

Since we have shown that $\ker \phi = H \cap K$, i.e. ϕ is one-to-one, we have proven that

$$G \cong \phi(G)$$

which is a subgroup of $G/H \times G/K$. □

17) Let $\phi : G_1 \mapsto G_2$ be a surjective group homomorphism. Let H_1 be a normal subgroup of G_1 and suppose that $\phi(H_1) = H_2$. Prove or disprove that $G_1/H_1 \cong G_2/H_2$.

Proof: We will disprove that $G_1/H_1 \cong G_2/H_2$ by giving a counterexample.

Let $G_1 = \mathbb{Z}, G_2 = \mathbb{Z}, H_1 = 2\mathbb{Z}, H_2 = 3\mathbb{Z}$. Then we see that obviously $G_1 \cong G_2$ and $H_1 \cong \mathbb{Z} \cong H_2$. However, for the quotient groups, we have that

$$G_1/H_1 = \mathbb{Z}/2\mathbb{Z} \cong \mathbb{Z}_2$$

and

$$G_2/H_2 = \mathbb{Z}/3\mathbb{Z} \cong \mathbb{Z}_3.$$

So $G_1/H_1 \not\cong G_2/H_2$. We have not explicitly defined our surjective group homomorphism ϕ yet.

All possible homomorphisms from $\mathbb{Z} \mapsto \mathbb{Z}$ are determined by the image of the generator, i.e. 1. So suppose then that $\phi(1) = n$ for $n \in \mathbb{Z}$. Then the homomorphism is

$$\begin{aligned}\phi(k) &= \phi(k \cdot 1) \\ &= k\phi(1) \\ &= kn\end{aligned}$$

for all $k \in \mathbb{Z}$. Thus, *all* of the possible homomorphisms from $\mathbb{Z} \mapsto \mathbb{Z}$ are of the form $\phi_n : \mathbb{Z} \mapsto \mathbb{Z}$ defined by $\phi_n(k) = kn$ for all $k, n \in \mathbb{Z}$. However, for the purposes of this problem, we are only interested in homomorphisms of \mathbb{Z} onto \mathbb{Z} . Since \mathbb{Z} only has two generators: 1 and -1 , we have that the two possible onto homomorphisms are

$$\begin{aligned}\phi_1 : \mathbb{Z} \mapsto \mathbb{Z}; \phi_1(k) &= k \\ \phi_{-1} : \mathbb{Z} \mapsto \mathbb{Z}; \phi_{-1}(k) &= -k.\end{aligned}$$

Since we have constructed a specific example showing that $G_1/H_1 \cong G_2/H_2$ does not always hold, we are done. □

5) Let G be a group and let i_g be an inner automorphism of G and define a map $G \mapsto \text{Aut}(G)$ by $g \mapsto i_g$. Prove that this map is a homomorphism with image $\text{Inn}(G)$ and kernel $Z(G)$. Use this result to conclude that

$$G/Z(G) \cong \text{Inn}(G)$$

Proof: Recall that $i_g(x) := gxg^{-1}$. We first show that the map is a homomorphism. Let $a, b \in G$. Then

$$\begin{aligned}i_{ab}(x) &= (ab)x(ab)^{-1} \\ &= abxb^{-1}a^{-1} \\ &= a(bxb^{-1})a^{-1} \\ &= i_a(i_b(x)) \\ &= (i_a i_b)(x)\end{aligned}$$

Since $i_{ab} = i_a i_b$ our map is a homomorphism.

By definition, we know that $\text{Inn}(G)$ is the set of inner automorphisms:

$$\text{Inn}(G) := \{i_g \mid g \in G\}.$$

So clearly the image for our homomorphism is $\text{Inn}(G)$ by definition. Next, we show that the kernel of this homomorphism is $Z(G)$, the center of G .

$$\begin{aligned} \{a \in G \mid i_a(x) = x \quad \forall x \in G\} &= \{a \in G \mid axa^{-1} = x \quad \forall x \in G\} \\ &= \{a \in G \mid ax = xa \quad \forall x \in G\} \\ &= Z(G) \end{aligned}$$

By the First Isomorphism Theorem, we conclude that

$$G/Z(G) \cong \text{Inn}(G).$$

□