

Abstract Algebra Homework 10

Joe Loser

April 24, 2016

This problem set includes problems 3b, 5b, 12, and an extra problem from section 17.4.

3) Use the division algorithm to find $q(x)$ and $r(x)$ such that $a(x) = q(x)b(x) + r(x)$ with $\deg r(x) < \deg b(x)$.

3b) $a(x) = 6x^4 - 2x^3 + x^2 - 3x + 1, b(x) = x^2 + x - 2$ in $\mathbb{Z}_7[x]$.

Solution: Performing long division, we have

$$\begin{array}{r}
 \overline{6x^2 - 8x + 21} \\
 x^2 + x - 2 \overline{) } \\
 \underline{-6x^4 - 6x^3 + 12x^2} \\
 -8x^3 + 13x^2 - 3x \\
 \underline{8x^3 + 8x^2 - 16x} \\
 21x^2 - 19x + 1 \\
 \underline{-21x^2 - 21x + 42} \\
 -40x + 43
 \end{array}$$

So we have that

$$\begin{aligned}
 a(x) &= (6x^2 - 8x + 21)(x^2 + x - 2) - 40x + 43 \\
 &\equiv (6x^2 + 6x)(x^2 + x - 2) + 2x + 1.
 \end{aligned}$$

Thus $q(x) = 6x^2 + 6x$ and $r(x) = 2x + 1$. □

5b) Find all of the zeros of $p(x) := 3x^3 - 4x^2 - x + 4$ in \mathbb{Z}_5 .

Solution: We first find a root of $p(x)$ by looking at $p(0), p(1), p(2), p(3)$, and $p(4)$. We see that

$$p(0) = 4 \neq 0$$

$$p(1) = 3 - 4 + 4 = 2 \neq 0$$

$$p(2) = 24 - 16 + 2 = 10 \equiv 0$$

$$p(3) = 81 - 36 + 1 \equiv 1 - 1 + 1 \equiv 1 \neq 0 \text{ and}$$

$$p(4) = 3(64) - 64 = 2(64) = 128 \equiv 3 \neq 0.$$

So $x = 2$ is a root of $p(x)$. By Corollary R21, 2 is a root of $p(x) \iff x - 2 \mid p(x)$. Performing the long division, we have that

$$\begin{array}{r}
 \overline{3x^2 + 2x + 3} \\
 x - 2 \overline{) } \\
 \underline{-3x^3 + 6x^2} \\
 2x^2 - x \\
 \underline{-2x^2 + 4x} \\
 3x + 4 \\
 \underline{-3x + 6} \\
 10
 \end{array}$$

Note that $10 \equiv 0$ so $r(x) = 0$. Thus

$$p(x) = (3x^2 + 2x + 3)(x - 2) \tag{1}$$

To further see that we cannot factorize this any more, we look to see if there are any roots of $q(x) := (3x^2 + 2x + 3)$ in \mathbb{Z}_5 . We have that

$$\begin{aligned}
q(0) &= 3 \neq 0 \\
q(1) &= 8 \equiv 3 \neq 0 \\
q(2) &= 12 + 4 + 3 \equiv 19 \equiv 4 \neq 0 \\
q(3) &= 3(9) + 6 + 3 = 27 + 9 = 36 \equiv 1 \neq 0 \\
q(4) &= 3(16) + 8 + 3 \equiv 3(1) + 3 + 3 \equiv 4 \neq 0
\end{aligned}$$

So $q(x)$ cannot be factorized any more. Thus we see that the only factorization of $p(x)$ in \mathbb{Z}_5 is as seen in equation (1). Hence the only root of $p(x)$ is $x = 2$. \square

12) If F is a field, show that $F[x_1, \dots, x_n]$ is an integral domain.

Proof: Note that since F is a field, F is also an integral domain. Recall by definition we have that

$$F[x_1, x_2] = (F[x_1])[x_2].$$

Without loss of generality, we will assume that $F[x_1, x_2]$ is the same as $F[x_2, x_1]$. We now proceed by induction.

i) Base Case: $n = 1$

By Corollary R19 in class, we immediately have that $F[x_1]$ is an integral domain.

ii) Induction Step

Assume that $F[x_1, \dots, x_{n-1}]$ is an integral domain. Then we show that $F[x_1, \dots, x_n]$ is an integral domain. We have that

$$F[x_1, \dots, x_n] = (F[x_1, \dots, x_{n-1}])[x_n].$$

Note that $(F[x_1, \dots, x_{n-1}])$ is an integral domain and adjoining another variable, x_n , still makes $F[x_1, \dots, x_n]$ an integral domain by the induction hypothesis. \square

E1) Write $\mathbb{Z}[\sqrt{7}]$ as a quotient ring of the polynomial ring $\mathbb{Z}[x]$ and then use this to find a familiar ring isomorphic to

$$R := \frac{\mathbb{Z}[\sqrt{7}]}{\langle 8 - \sqrt{7} \rangle}.$$

Solution: First recall that for any integer $n \in \mathbb{Z}$ we have that

$$\mathbb{Z}[\sqrt{n}] := \{a + b\sqrt{n} \mid a, b \in \mathbb{Z}\}.$$

Note that $x^2 - 7 = 0$ i.e. $x = \sqrt{7}$. Define $\phi : \mathbb{Z}[x] \rightarrow \mathbb{Z}[\sqrt{7}]$ by $\phi(f(x)) = f(\sqrt{7})$. We now show that ϕ is a ring homomorphism and is onto.

Let $a, b \in \mathbb{Z}[x]$. Then we have that

$$\begin{aligned}
\phi(a + b) &= (a + b)(\sqrt{7}) \\
&= a(\sqrt{7}) + b(\sqrt{7}) \\
&= \phi(a) + \phi(b).
\end{aligned}$$

Also,

$$\begin{aligned}
\phi(a \cdot b) &= (a \cdot b)(\sqrt{7}) \\
&= a(\sqrt{7}) \cdot b(\sqrt{7}) \\
&= \phi(a) \cdot \phi(b).
\end{aligned}$$

So ϕ is a ring homomorphism.

Next we determine the kernel of ϕ . We see that

$$\begin{aligned}
\ker \phi &= \{f(x) \in \mathbb{Z}[x] \mid f(\sqrt{7}) = 0\} \\
&= \{g(x) \cdot (x^2 - 7) \mid g(x) \in \mathbb{Z}[x]\} \\
&= (x^2 - 7)\mathbb{Z}[x].
\end{aligned}$$

To show ϕ is onto, note every element of $\mathbb{Z}[\sqrt{7}]$ is of the form $a + b\sqrt{7}$ where $a, b \in \mathbb{Z}$ and $a + b\sqrt{7} = \phi(a + bx)$. Thus ϕ is onto.

Applying the First Isomorphism Theorem, we see that

$$\mathbb{Z}[\sqrt{7}] \cong \frac{\mathbb{Z}[x]}{(x^2 - 7)\mathbb{Z}[x]}.$$

Using this expression, we now have that

$$R \cong \frac{\frac{\mathbb{Z}[x]}{(x^2 - 7)\mathbb{Z}[x]}}{(8 - x)\mathbb{Z}[x]}.$$

Trying to get things into a form where we can use the Third Isomorphism Theorem, we get

$$R \cong \frac{\frac{\mathbb{Z}[x]}{(x^2 - 7)\mathbb{Z}[x]}}{\frac{(8 - x, x^2 - 7)\mathbb{Z}[x]}{(x^2 - 7)\mathbb{Z}[x]}}.$$

Applying the Third Isomorphism now yields

$$R \cong \frac{\mathbb{Z}[x]}{(8 - x, x^2 - 7)\mathbb{Z}[x]}.$$

Now we try to make $8 - x = 0$ i.e. $x = 8$. Define a map $\psi : \mathbb{Z}[x] \rightarrow \mathbb{Z}[8]$ by $\psi(p(x)) = p(8)$. First note that adjoining by 8 does not give us anything new. So $\mathbb{Z}[8] = \mathbb{Z}$. We now show that ψ is indeed a ring homomorphism.

Let $a, b \in \mathbb{Z}[x]$. Then

$$\begin{aligned}\psi(a + b) &= (a + b)(8) \\ &= a(8) + b(8) \\ &= \psi(a) + \psi(b).\end{aligned}$$

Also,

$$\begin{aligned}\psi(a \cdot b) &= (a \cdot b)(8) \\ &= a(8) \cdot b(8) \\ &= \psi(a) \cdot \psi(b).\end{aligned}$$

So ψ is a ring homomorphism.

We can find the kernel of ψ by:

$$\begin{aligned}\ker \psi &= \{f(x) \in \mathbb{Z}[x] \mid f(8) = 0\} \\ &= \{g(x) \cdot (8 - x) \mid g(x) \in \mathbb{Z}[x]\} \\ &= (8 - x)\mathbb{Z}[x].\end{aligned}$$

To show ψ is onto, note every element of $\mathbb{Z}[8]$ is of the form $a + b(8)$ where $a, b \in \mathbb{Z}$ and $a + b(8) = \psi(a + bx)$. Thus ψ is onto.

Note that

$$(8 - x, x^2 - 7)\mathbb{Z}[x] = \{(8 - x)a(x) + (x^2 - 7)b(x) \mid a(x), b(x) \in \mathbb{Z}[x]\}.$$

Applying our mapping ψ now yields

$$\begin{aligned}R &\cong \frac{\mathbb{Z}}{(8 - 8, 8^2 - 7)\mathbb{Z}} \\ &= \frac{\mathbb{Z}}{(0, 57)\mathbb{Z}} \\ &\cong \frac{\mathbb{Z}}{57\mathbb{Z}} \\ &\cong \mathbb{Z}_{57}.\end{aligned}$$

□