

Abstract Algebra Homework 11

Joe Loser

April 30, 2016

This problem set includes problems 3c, 4b, 24, and an extra problem from section 17.4.

3) Use the division algorithm to find $q(x)$ and $r(x)$ such that $a(x) = q(x)b(x) + r(x)$ with $\deg r(x) < \deg b(x)$.

3c) $a(x) = 4x^5 - x^3 + x^2 + 4$ and $b(x) = x^3 - 2$ where $a(x), b(x) \in \mathbb{Z}_5[x]$.

Solution: Performing long division, we have

$$\begin{array}{r}
 4x^2 - 1 \\
 x^3 - 2 \overline{) 4x^5 - x^3 + x^2 + 4} \\
 \underline{-4x^5} + 8x^2 \\
 -x^3 + 9x^2 + 4 \\
 \underline{x^3} -2 \\
 9x^2 + 2
 \end{array}$$

Thus,

$$\begin{aligned}
 a(x) &= (4x^2 - 1) \cdot (x^3 - 2) + (9x^2 + 2) \\
 &\equiv (4x^2 - 1) \cdot (x^3 - 2) + (4x^2 + 2).
 \end{aligned}$$

□

4) Find the greatest common divisor of each of the following pairs $p(x)$ and $q(x)$ of polynomials. If $d(x) = \gcd(p(x), q(x))$, find two polynomials $a(x)$ and $b(x)$ such that $d(x) = a(x)p(x) + b(x)q(x)$.

4b) $p(x) = x^3 + x^2 - x + 1$ and $q(x) = x^3 + x - 1$ where $p(x), q(x) \in \mathbb{Z}_2[x]$.

Performing the first stage of long division, we have that

$$\begin{array}{r}
 1 \\
 x^3 + x^2 - x + 1 \overline{) x^3 + x - 1} \\
 \underline{-x^3 - x^2} + x - 1 \\
 -x^2 + 2x - 2
 \end{array}$$

Note that in $\mathbb{Z}_2[x]$, $-x^2 + 2x - 2 \equiv -x^2$. So

$$x^3 + x - 1 = 1 \cdot (x^3 + x^2 - x + 1) + (-x^2). \quad (1)$$

In the second stage of long division, we get

$$\begin{array}{r}
 -x - 1 \\
 -x^2 \overline{) x^3 + x^2 - x + 1} \\
 \underline{-x^3} \\
 x^2 \\
 \underline{-x^2} \\
 -x - 1
 \end{array}$$

So

$$x^3 + x^2 - x + 1 = (-x - 1) \cdot (-x^2) + (-x + 1). \quad (2)$$

In the third stage, we get

$$\begin{array}{r} x+1 \\ -x+1 \overline{) -x^2} \\ \underline{x^2-x} \\ -x \\ \underline{x-1} \\ -1 \end{array}$$

So

$$-x^2 = (-x + 1) \cdot (x + 1) + (-1). \quad (3)$$

In the last stage of long division, we have

$$\begin{array}{r} x-1 \\ -1 \overline{) -x+1} \\ \underline{x} \\ 1 \\ \underline{-1} \\ 0 \end{array}$$

So

$$-x + 1 = (-1) \cdot (x - 1) + 0. \quad (4)$$

Thus $\gcd(p(x), q(x)) = -1$. Performing the back substitution, we have the following

$$\begin{aligned} -1 &= -x^2 - (x + 1)(-x + 1) \\ &= -x^2 - (x + 1)((x^3 + x^2 - x + 1) - (-x - 1)(-x^2)) \\ &= (x^3 + x - 1) - (x^3 + x^2 - x + 1) - (x + 1)[(x^3 + x^2 - x + 1) - (-x - 1)((x^3 + x - 1) - (x^3 + x^2 - x + 1))] \end{aligned}$$

finish back
sub

24) Show that $x^p - x$ has p distinct zeros in \mathbb{Z}_p for any prime p . Conclude that

$$x^p - x = x(x - 1)(x - 2) \cdots (x - (p - 1)).$$

Proof: By Fermat's Little Theorem, for all $a \in \mathbb{Z}_p$ we have that $a^p = a$. So $a^p - a = 0$. Thus every $a \in \mathbb{Z}_p$ is a zero of the polynomial $x^p - x$. Note that the polynomial has degree p and p zeros in \mathbb{Z}_p . The numbers $0, 1, \dots, p - 1$ are the roots of the equation $x^p - x$, i.e. the p distinct roots. Hence it must split into p distinct linear factors in $\mathbb{Z}_p[x]$ as follows:

$$x^p - x = x(x - 1)(x - 2) \cdots (x - (p - 1)).$$

□

E1) Construct a field with 8 elements.

Solution: Since $8 = 2^3$ we start with a field \mathbb{Z}_2 of characteristic 2 and look for an irreducible polynomial of degree 3 in $\mathbb{Z}_2[x]$. Such a polynomial is $p(x) = x^3 + x + 1$.

We will show that

$$K := \frac{\mathbb{Z}_2[x]}{\langle x^3 + x + 1 \rangle}$$

is a field of 8 elements.

To see why $p(x)$ is irreducible in $\mathbb{Z}_2[x]$, since it of degree 3 or lower, we can look at all of the roots in \mathbb{Z}_2 . We have that $g(0) = 1$ and $g(1) = 3 \equiv 1$. So neither 0 or 1 are a root of $p(x)$. Hence we see that $p(x)$ is irreducible over $\mathbb{Z}_2[x]$.

By