

MAT 4381

Bayesian Inference

Study Guide

Winter 2024

$$P(A|B) = \frac{P(B|A) P(A)}{P(B)}$$

Bayesian vs. Frequentist

Frequentist: The parameter is not a random variable

Bayesian: The parameter is a random variable

Bayes Theorem

Ω be the sample space
 E_1, E_2, \dots sequence of exhaustive events

$A \subset \Omega$ another event

$$P(E_k|A) = \frac{P(A|E_k)P(E_k)}{\sum_i P(A|E_i)P(E_i)}$$

Bernoulli Trials

X takes value 1 if event occurs and 0 if event do not occur

Bernoulli Distribution: $Ber(\theta)$

$$P(X=1) = \theta$$

$$P(X=0) = 1-\theta$$

$$P(X=x) = f(x|\theta) = \theta^x(1-\theta)^{1-x} I(x \in \{0,1\}, \theta)$$

$$E[X] = \sum_{x=0}^1 x f(x|\theta) = \theta$$

$$\text{Var}(X) = \theta(1-\theta)$$

Binomial Distribution

$$X_1, \dots, X_n \stackrel{iid}{\sim} Ber(\theta)$$

$$Y = \sum_{i=1}^n X_i \in \{0, 1, 2, \dots, n\} \sim \text{Bin}(n, \theta)$$

$$P(Y=y) = \binom{n}{y} \theta^y (1-\theta)^{n-y} I(y \in \{0, 1, \dots, n\})$$

$$E[Y] = n\theta \quad \text{Var}(Y) = n\theta(1-\theta)$$

Bayes Formula for Distributions

$$f(\theta|y) = \frac{f(y|\theta)g(\theta)}{f_y(y)}$$

If θ is discrete with p.m.f. $g(\theta)$

$$f(\theta|y) = \frac{f(y|\theta)g(\theta)}{f_y(y)} = \frac{f(y|\theta)g(\theta)}{\sum_j f(y|\theta_j)g(\theta_j)}$$

If $\theta \sim g(\theta)$ is continuous

$$f(\theta|y) = \frac{f(y|\theta)g(\theta)}{f_y(y)} = \int f(y|\theta)g(\theta)d\theta$$

The denominator is sometimes called the proportionality constant

$$f(\theta|y) = C f(y|\theta) g(\theta)$$

or $f(\theta|y) \propto f(y|\theta)g(\theta)$ = likelihood \times prior

C some constant and \propto means "proportional to"

$g(\theta)$: prior distribution for θ

$f(\theta|y)$: posterior distribution

Conjugate Prior

- A conjugate prior is also a convenient prior, since it allows one to easily calculate the posterior distribution


Beta Distribution

$$g(\theta) = \frac{\Gamma(a+\beta)}{\Gamma(a)\Gamma(\beta)} \theta^{a-1} (1-\theta)^{\beta-1} I(0 < \theta < 1)$$

The distribution is the Beta(a, β) distribution

We use the fact that

$$\frac{\Gamma(a)}{x^a} = \int_0^\infty x^{a-1} e^{-bx} dx, a, b > 0$$



The Beta distribution is a conjugate prior for binomial distribution

If $(Y|\theta=B) \sim f(y|\theta)$
 $f(y|\theta) = \binom{n}{y} \theta^y (1-\theta)^{n-y} I(y \in \{0, 1, \dots, n\})$
 and $(\theta | \alpha, \beta) \sim \text{Beta}(\alpha, \beta)$
 we have

$$f(\theta|y) \propto f(y|\theta)g(\theta)$$

In other words,
 $f(\theta|y) \propto \theta^y (1-\theta)^{n-y} \theta^{a-1} (1-\theta)^{\beta-1}$

The posterior is:
 $f(\theta|y) = \frac{\Gamma(a+\beta)}{\Gamma(a)\Gamma(\beta)} \theta^{a+y-1} (1-\theta)^{\beta+n-y-1}$

If the prior is Beta(α, β), then the posterior is Beta($a+y, \beta+n-y$)

If $\theta \sim \text{Beta}(\alpha, \beta)$, then

$$E[\theta] = \frac{\alpha}{\alpha+\beta}$$

$$\text{Var}(\theta) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$$

$$\text{Mode: } \hat{\theta} = \frac{\alpha-1}{\alpha+\beta-2}$$

i) For large values of α and β mode and mean are roughly the same.

ii) Median for Beta(α, β) cannot be written in a closed form

iii) Peak of mode of beta density gets sharper as $\alpha+\beta$ increases

iv) We need to use numerical methods to calculate the median for different values of α and β

Credible Region

- We want to find I s.t. $P(\theta \in I) = 1-\alpha$, where $1-\alpha$ is the confidence level
- We can also use simulation method

Choosing Beta function

- First, consider priors that don't express a strong opinion.
- Later, consider priors which incorporate a strong prior belief. If there are no preferences on the value for θ , then one might consider using a

"flat prior". That is, the uniform distribution between 0 and 1 ($\text{Beta}(1,1)$)

In some sense, this implies that each value of θ is equally likely

Since the prior is flat, the posterior distribution has the same shape as the likelihood multiplied by a constant C

$$h(\theta|x) = C f(x|\theta)g(\theta) \propto f(x|\theta)$$

Since the uniform distribution is Beta($1,1$), then the posterior is Beta($y+1, n-y+1$)

Making Bayesian Inference

The posterior contains the results of the analysis

To communicate the result, we can either present the entire posterior distribution, or give summary statistics of the posterior

For credible regions, there are 2 common types:

- Smallest interval
- Equal tail area

Jeffreys Prior

Jeffreys proposed that an acceptable "non-informative prior finding principle" should be invariant under monotone transformations of the parameter

The Jeffreys prior satisfying this invariance is proportional to $I(\theta)$, where $I(\theta)$ is the Fisher information on θ . Can be generalized when θ is k-dimensional to $\sqrt{\det(I(\theta))}$

$$I(\theta) = -E\left[\frac{\partial^2 \ln f(x|\theta)}{\partial \theta^2}\right]$$

For k-dimensional, we have

$$I(\theta) = E\left[\left(\frac{\partial \ln f(x|\theta)}{\partial \theta_i \partial \theta_j}\right)_{(i,j)}\right]$$

Multinomial Distribution

$$(X_1, \dots, X_k) \sim \text{Multinomial}(n, P_1, \dots, P_k)$$

$$P_k = P_1 + P_2 + \dots + P_{k-1}$$

Models probability of counts for each side of an experiment with k exhaustive outcomes

$$f(X_1, X_2, \dots, X_k) = \binom{n}{X_1, X_2, \dots, X_k} P_1^{X_1} P_2^{X_2} \dots P_k^{X_k} (1-P_1-P_2-\dots-P_{k-1})^{X_k}$$

where $X_i = 0, 1, 2, \dots, n$, $X_1 + X_2 + \dots + X_k = n$

Simple Case: Trinomial Distribution

$$(X, Y) \sim \text{Multinomial}(n, P_1, P_2)$$

$$f(x,y) = \binom{n}{x,y} P_1^x P_2^y (1-P_1-P_2)^{n-x-y}$$

$$X, Y = 0, 1, 2, \dots, n, X+Y \leq n$$

We can also get that

$$X \sim \text{Bin}(n, P_1)$$

$$Y \sim \text{Bin}(n, P_2)$$

$$Y|X=x \sim \text{Bin}(n-x, \frac{P_2}{1-P_1})$$

$$E[Y|X=x] = \frac{P_2}{1-P_1} (n-x)$$

$$\text{Var}[Y|X=x] = \frac{P_2}{1-P_1} (1-\frac{P_2}{1-P_1})(n-x)$$

$$E[X] = \frac{P_1}{1-P_1} n, E[Y] = \frac{P_2}{1-P_1} n$$

$$\text{Cov}(X, Y) = -n(n+1) P_1 P_2$$

$$P(X, Y) = \frac{-n(n+1) P_1 P_2}{\sqrt{P_1(1-P_1)P_2(1-P_2)}} = \frac{-(n+1) P_1 P_2}{\sqrt{P_1 P_2 (1-P_1)(1-P_2)}}$$

Jeffreys Prior for Trinomial Distribution

$$g(P_1, P_2) \propto \frac{1}{\sqrt{P_1 P_2 (1-P_1)(1-P_2)}}$$

Special case of Dirichlet distribution
 $\text{Dir}(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

Dirichlet Distribution

$$(X_1, X_2, \dots, X_k) \sim \text{Dir}(\alpha_1, \alpha_2, \dots, \alpha_k)$$

$$f(x_1, \dots, x_k) = \frac{\prod_{i=1}^k x_i^{\alpha_i-1}}{\prod_{i=1}^k \Gamma(\alpha_i)} \left(\prod_{i=1}^k x_i\right)^{1-\sum_{i=1}^k \alpha_i}$$

$$\text{Defined on } S = \{(x_1, \dots, x_k) : \sum_{i=1}^k x_i \leq 1, x_i \geq 0, i=1, \dots, k\}$$

Posterior for Trinomial

$$(\theta_1, \dots, \theta_{k-1} | X_1=x_1, \dots, X_k=x_{k-1}) \sim \text{Dir}(\alpha_1+x_1, \dots, \alpha_{k-1}+x_{k-1}, n-\sum x_i)$$

$$E[\theta_i | X_1=x_1, \dots, X_{k-1}=x_{k-1}] = \frac{\alpha_i + x_i}{\sum_{j=1}^k \alpha_j + n}$$

$$\theta = [\theta_1, \theta_2, \dots, \theta_{k-1}, \theta_k]^T$$

$$\theta_k = 1 - \theta_1 - \dots - \theta_{k-1} \quad x_k = n - x_1 - \dots - x_{k-1}$$

$$\text{Notice: } (\theta_1, \theta_2) \sim \text{Dir}(\alpha_1, \alpha_2, \alpha_3 + \dots + \alpha_k)$$

Basic Properties of Dirichlet Distribution

$$(Y_1, \dots, Y_{k-1}) \sim \text{Dir}(\alpha_1, \dots, \alpha_{k-1}) \quad \alpha_0 = \sum_{i=1}^k \alpha_i$$

$$E[Y_i] = \frac{\alpha_i}{\alpha_0} \quad \text{Var}[Y_i] = \frac{\alpha_i(\alpha_0 - \alpha_i)}{\alpha_0^2(\alpha_0 + 1)}$$

$$\text{Cov}(X_i, X_j) = -\frac{\alpha_i \alpha_j}{\alpha_0^2(\alpha_0 + 1)}$$

Conjugate Prior and Posterior for Poisson

$$X_1, \dots, X_n \stackrel{iid}{\sim} f(x|\theta) = \frac{\exp(-\theta)x^x}{x!} I(x=0, 1, \dots)$$

$$\text{Likelihood: } \prod_{i=1}^n f(x_i|\theta) = \prod_{i=1}^n \frac{\exp(-\theta)x_i^x}{x_i!} \propto \exp(-n\theta) \theta^n$$

$$y = \sum_{i=1}^n x_i$$

$$\text{Conjugate Prior: } g(\theta) = \frac{\theta^x}{\Gamma(x)} \theta^{x-1} \exp(-\theta) I(\theta > 0)$$

$$\text{Posterior: } h(\theta|y) \propto \theta^{y+x+1} \exp(-(\theta+n)\theta) I(\theta > 0)$$

$$h(\theta|y) = \frac{(\theta+n)^{y+x}}{\Gamma(y+x)} \theta^{y+x+1} \exp(-(\theta+n)\theta) I(\theta > 0)$$

$$E[\theta|Y=y] = \frac{y+x}{n+x}$$

$$\text{Var}[\theta|Y=y] = \frac{y+x}{(n+x)^2}$$

$$\text{Mode}(\theta|Y=y) = \frac{y+x-1}{n+x}$$

Jeffrey's Prior for Poisson

$$f(x|\lambda) = \frac{\exp(-\lambda)\lambda^{\sum_{i=1}^n x_i}}{\prod_{i=1}^n x_i!}$$

$$-\mathbb{E}\left[\frac{\partial^2}{\partial \theta^2} \log f(x|\theta)\right] = \frac{n}{\lambda}$$

Jeffrey's prior: $g(\lambda) \propto \frac{1}{\lambda}$

This leads to an improper prior since we have

$$\int_0^\infty \frac{d\lambda}{\sqrt{\lambda}} = \infty$$

No normalizing constant will turn it into a prior p.d.f.

We can still use it to calculate the posterior

$$\lambda|x \propto \exp(-n\lambda)^{\frac{n}{2} + \frac{1}{2}}$$

The use of improper prior leads to Gamma posterior

$$\lambda|x \sim \frac{1}{\Gamma(\frac{n}{2} + \frac{1}{2})} \lambda^{\frac{n}{2} + \frac{1}{2}} \exp(-n\lambda)$$

$$\mathbb{E}[\lambda|x_1, \dots, x_n] = \frac{\sum_{i=1}^n x_i + 0.5}{n}$$

$$\text{Var}[\lambda|x_1, \dots, x_n] = \frac{\sum_{i=1}^n x_i + 0.5}{n^2}$$

Normal Distribution

$$(y_1, \dots, y_n|\mu) \sim f(y_1, \dots, y_n|\mu) \propto \exp\left(-\sum_{i=1}^n \frac{(y_i - \mu)^2}{2\sigma^2}\right)$$

σ^2 known

$$f(y_1, \dots, y_n|\mu) \propto \exp\left(-\frac{n(\mu - \bar{y})^2}{2\sigma^2}\right)$$

The Jeffrey's prior is

$$g(\mu) \propto 1$$

This constitutes an improper prior

However, we can still calculate the posterior

$$(\mu|Y=y_1, \dots, Y_n=y_n) \sim N(\bar{y}, \frac{\sigma^2}{n})$$

The normal prior

$$Y_1, \dots, Y_n \stackrel{iid}{\sim} N(\mu, \sigma^2)$$

σ^2 known

$$\text{Prior: } g(\theta) \propto \exp(-(\theta - \mu)^2/\sigma^2)$$

$$\text{Likelihood: } \prod_{i=1}^n f(y_i, \dots, y_n|\mu) \propto \exp\left(-\frac{(y_i - \mu)^2}{2\sigma^2}\right)$$

Posterior for θ

$$\begin{aligned} h(\theta|y_1, \dots, y_n) &\propto \exp\left(-\frac{(\theta - \mu)^2}{\sigma^2} - \frac{(y_i - \mu)^2}{2\sigma^2}\right) \\ &\propto \exp\left(-\frac{(\theta - \mu)^2}{\sigma^2} - \frac{(y_i - \mu)^2}{2\sigma^2} - \frac{2(\theta - \mu)(y_i - \mu)}{2\sigma^2}\right) \\ &\propto \exp\left(-\frac{(\theta - \mu)^2}{\sigma^2} - \frac{2(\theta - \mu)(y_i - \mu)}{2\sigma^2} - \frac{(y_i - \mu)^2}{2\sigma^2}\right) \end{aligned}$$

$$(y_1, \dots, y_n) \sim N\left(\frac{\sigma^2 \mu + n\bar{y}}{\sigma^2 + n}, \frac{\sigma^2}{\sigma^2 + n}\right)$$

$$\mathbb{E}[\theta|Y=y] = \frac{\sigma^2 \mu + n\bar{y}}{\sigma^2 + n}$$

$$\text{MAP} = \mathbb{E}[\theta|Y=y] = \text{Median}(\theta|Y=y)$$

$$\text{Posterior precision: } \frac{\tau^2 \sigma^2}{\tau^2 \sigma^2 + n} = \frac{1}{\sigma^2 n} + \frac{1}{\tau^2}$$

Posterior precision is equal to prior precision plus observation precision

Since $E[\theta|y_1, \dots, y_n] = \frac{\sigma^2 \mu + n\bar{y}}{\sigma^2 + n} + \frac{\tau^2}{\sigma^2 + n} \bar{y}$, the posterior mean is the weighted average of the prior mean and the mean of observations, where the weights are the proportions of the precision to the posterior precision.

Credible region for θ :

$$\frac{\tau^2 \bar{y} + n\mu^2/n}{\tau^2 + \sigma^2/n} \pm \frac{\tau^2}{\sqrt{\tau^2 + \sigma^2/n}}$$

If σ^2 unknown, we can use

$$\hat{\sigma}^2 = \frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2$$

We change $Z_{\alpha/2}$ to $t_{\alpha/2}(n-1)$

Credible region becomes

$$\frac{\tau^2 \bar{y} + n\mu^2/n}{\tau^2 + \hat{\sigma}^2/n} \pm t_{\alpha/2}(n-1) \frac{\sqrt{\hat{\sigma}^2/n}}{\sqrt{\tau^2 + \hat{\sigma}^2/n}}$$

Jeffrey's prior for σ and σ^2

Suppose μ is fixed

Then we get the Jeffrey's prior for σ and σ^2 are

$$g(\sigma) \propto \frac{1}{\sigma} \text{ and } g(\sigma^2) \propto \frac{1}{\sigma^2}$$

Bayes Factor and Relative Evidence

In model selection, say 2 models.

The Bayes factor is defined by

$$BF(1,2) = \frac{P(M_1|\text{Data})P(M_2)}{P(M_2|\text{Data})P(M_1)}$$

With a prior on model, the Bayes factor can be calculated. With equal probability on models, we get

$$BF(1,2) = \frac{P(M_1, \text{Data})}{P(M_2, \text{Data})}$$

Inverse χ^2 Distribution

$V \sim \chi^2(r)$. The p.d.f. for V is

$$f(v) = \frac{1}{2^{r/2} \Gamma(r/2)} v^{\frac{r}{2}-1} I(v>0)$$

$$\text{Define } U = \frac{1}{V}$$

We say $U \sim \text{Inv-}\chi^2(v)$

The p.d.f. for inverse- χ^2 is

$$g(u) = \frac{1}{2^{r/2} \Gamma(r/2)} u^{-\frac{r}{2}-1} I(u>0)$$

$$E[U] = \frac{1}{r-2} \quad \text{Mode}(U) = \frac{1}{r+2}$$

$$\text{Var}(U) = \frac{2}{(r-2)^2(r-4)}$$

Mean exists if $r > 2$ and $r > 4$

Inverse Gamma Distribution

$$X \sim \text{Gamma}(\alpha, \beta)$$

$$f(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} \exp(-\beta x) x^{\alpha-1} I(x>0)$$

Define $U = X^{-1}$. The p.d.f. for U is

$$g(u) = \frac{\beta^\alpha}{\Gamma(\alpha)} \exp(-\beta/u) u^{-\alpha-1} I(u>0)$$

Denote by $U \sim \text{IG}(\alpha, \beta)$

$$E[U] = \frac{\beta}{\alpha-1} \quad \text{Mode}(U) = \frac{\beta}{\alpha+1}$$

$$\text{Var}(U) = \frac{\beta^2}{(\alpha-1)^2(\alpha-2)}$$

Inverse Gamma prior on σ^2 , μ fixed

$$(y_1, \dots, y_n|\mu, \sigma^2) \stackrel{iid}{\sim} N(\mu, \sigma^2)$$

$$g(\sigma^2) = \frac{\beta^\alpha}{\Gamma(\alpha)} \exp(-\frac{\beta}{\sigma^2}) (\sigma^2)^{-\alpha-1}$$

$$h(\sigma^2|\mu, y_1, \dots, y_n) \propto (\sigma^2)^{-\alpha-1} \exp\left(-\frac{\sum (y_i - \mu)^2}{2\sigma^2}\right) \cdot \exp(-\beta/\sigma^2) \cdot (\sigma^2)^{-\alpha-1}$$

$$\propto (\sigma^2)^{-(\alpha+\frac{1}{2})} \exp\left(-\frac{\beta + \sum (y_i - \mu)^2}{2\sigma^2}\right)$$

Therefore

$$(\sigma^2|\mu, y_1, \dots, y_n) \sim \text{IG}\left(\alpha + \frac{n}{2}, \beta + \frac{\sum (y_i - \mu)^2}{2}\right)$$

In this case

$$E[\sigma^2|\mu, y_1, \dots, y_n] = \frac{\beta + \sum_{i=1}^n (y_i - \mu)^2 / 2}{\alpha + \frac{n}{2} - 1}$$

$$\text{Mode} = \frac{\beta + \sum_{i=1}^n (y_i - \mu)^2 / 2}{\alpha + \frac{n}{2} + 1}$$

If $X \sim \chi^2(v)$, then $\frac{1}{X} \sim \text{Inv-}\chi^2(v)$

Inverse chi-squared distribution is a special case of gamma distribution with parameters $\alpha = v/2$, $\beta = 1/2$, or $\alpha = v/2$, $\beta = v^2/2$ for scaled inverse- χ^2 distribution

Posterior Predictive Distribution

Let y_{n+1} be the next observations drawn after the random sample $Y_n = (y_1, y_2, \dots, y_n)$.

The predictive accuracy for $(y_{n+1}|y_1, y_2, \dots, y_n)$ is the conditional density

$$f(y_{n+1} | Y_n) = \int f(y_{n+1}|\mu) g(\mu|Y_n) d\mu$$

Jeffrey's Prior for normal family with μ and σ^2 unknown

$$X \sim N(\mu, \sigma^2) \quad \Theta = [\mu \ \sigma^2]^T$$

Then we get

$$-E\left[\frac{\partial^2 \ln f}{\partial \mu^2}\right] = \frac{1}{\sigma^2} \quad -E\left[\frac{\partial^2 \ln f}{\partial \sigma^2}\right] = 0$$

$$-E\left[\frac{\partial^2 \ln f}{\partial \mu \partial \sigma^2}\right] = \frac{1}{\sigma^4}$$

The Jeffrey's prior is

$$g(\mu, \sigma^2) \propto (\sigma^2)^{-\frac{3}{2}}$$

To calculate posterior, we get

$$h(\mu, \sigma^2 | y_1, \dots, y_n) \propto (\sigma^2)^{-\alpha-\frac{n}{2}} \exp\left(-\frac{(n-1)\sigma^2 + (\mu - \bar{y})^2}{2\sigma^2}\right)$$

This shows that

$$(\mu, \sigma^2 | y_1, \dots, y_n) \sim N-\text{IG}(\mu = \bar{y}, \lambda = n, \alpha = n, \beta = \sum_{i=1}^n (y_i - \bar{y})^2)$$

Normal Inverse Gamma Distribution

Let $\lambda > 0$ and

$$(y|\mu, \sigma^2, \lambda) \sim N(\mu, \sigma^2/\lambda)$$

$$(\sigma^2 | \mu, \lambda) \sim \text{IG}(\mu, \lambda)$$

Then we say (y, σ^2) has normal inverse gamma distribution with parameters $(\mu, \lambda, \alpha, \beta)$ which is denoted by

$$(y, \sigma^2) \sim \text{NIG}(\mu, \lambda, \alpha, \beta)$$

$$(y, \sigma^2) \sim N-\text{IG}(\mu, \lambda, \alpha, \beta)$$

We can write the joint p.d.f. for (y, σ^2) as follows

$$f(y, \sigma^2 | \mu, \lambda, \alpha, \beta) = \frac{1}{\sqrt{2\pi}} \frac{\beta^\alpha}{\Gamma(\alpha)} (\sigma^2)^{-\alpha-1} \exp\left(-\frac{\beta + (y - \mu)^2}{2\sigma^2}\right)$$

year, $\sigma^2 > 0$

By definition, $(\sigma^2 | \mu, \lambda) \sim \text{IG}(\mu, \lambda)$

We can get that $y \sim t(2\alpha, \mu, \beta/\lambda\alpha)$

Student t distribution

Let $W \sim N(0,1)$ and $V \sim \chi^2(r)$ and W and V are mutually independent. The joint distribution for W and V is

$$f(v, w) = \frac{1}{\sqrt{2\pi}} \frac{1}{2^{r/2} \Gamma(r/2)} \exp\left(-\frac{w^2}{2}\right) V^{\frac{r}{2}-1} \exp(-v^2/2)$$

for $w \in R$, $v > 0$

If we define $T = \frac{W}{\sqrt{V}}$ and $U = V$,

we get

$$g(u, t) = \frac{1}{\sqrt{\pi t}} \frac{1}{2^{(r+1)/2}} u^{(r+1)/2-1} \exp(-u(u+\frac{t}{u}))$$

$$g(t) = \frac{1}{\sqrt{\pi t}} \frac{\Gamma((r+1)/2)}{\Gamma(r/2)(1+\frac{t}{u})^{(r+1)/2}}$$

when $t > 0$

This is known as the student t distribution

Notes:

- $g(t)$ is a Cauchy distribution if $r=1$

- $g(t) \rightarrow N(0,1)$ as $r \rightarrow \infty$

If we use the transformation $V = \sigma^2 t + \mu$ when $\sigma > 0$, we get

$$h(v) = \frac{1}{\sigma \sqrt{\pi r}} \frac{\Gamma((r+1)/2)}{\Gamma(r/2)} \frac{1}{(1 + \frac{v-\mu}{\sigma})^{(r+1)/2}}$$

Denote this distribution by $t(r, \mu, \sigma)$

If we let $U = \sigma \left(\frac{W}{\sqrt{V}} \right) + \mu$, we get

$$\text{Median}(U) = \text{Mode}(U) = E[U] = \sigma E[W] \frac{1}{\sqrt{V}} + \mu$$

If $E[\frac{1}{\sqrt{V}}]$ exists then $E[U] = \mu$

$$\text{Var}(U) = \frac{r\sigma^2}{r-2}$$

Properties of Normal Inverse Gamma Distribution

$$(y, \sigma^2) \sim N-\text{IG}(\mu, \lambda, \alpha, \beta) \Rightarrow (\sigma^2 | \mu, \lambda) \sim \text{IG}(\mu, \lambda)$$

$$E[\sigma^2 | \alpha, \beta] = \frac{\beta}{\alpha-1}$$

$$\text{Var}[\sigma^2 | \alpha, \beta] = \frac{\beta^2}{(\alpha-1)^2(\alpha-2)}$$

$$\text{Var}[y] = E\left[\frac{\sigma^2}{\lambda}\right] = \frac{\beta}{\lambda(\alpha-1)}$$

Normal-Inverse Gamma Prior

$$(y_1, \dots, y_n | \mu, \sigma^2) \stackrel{\text{iid}}{\sim} N(\mu, \sigma^2)$$

$$\mu | \sigma^2 \sim N(\mu_0, \sigma^2/n_0)$$

$$\sigma^2 \sim IG(\alpha, \beta)$$

n_0 : equivalent sample size of the prior
 n : equivalent sample size we would obtain for n observations

The posterior is

$$f(\mu, \sigma^2 | y_1, \dots, y_n) \propto (\sigma^2)^{-\frac{n}{2}} \exp\left(-\frac{\sum_{i=1}^n (y_i - \mu)^2}{2\sigma^2}\right)$$

$$\times (\sigma^2)^{-\frac{1}{2}} \exp\left(-\frac{n_0(\mu - \mu_0)^2}{2\sigma^2}\right)$$

$$\times (\sigma^2)^{-(\alpha+\beta)} \exp\left(\frac{\beta}{\sigma^2}\right)$$

Therefore,

$$f(\mu, \sigma^2 | y_1, \dots, y_n) \propto (\sigma^2)^{-(n+2\alpha+3)/2}$$

$$\times \exp\left(-\frac{\sum_{i=1}^n (y_i - \mu)^2}{2\sigma^2} - \frac{n_0(\mu - \mu_0)^2}{2\sigma^2} - \frac{\beta}{\sigma^2}\right)$$

Expand to get

$$f(\mu, \sigma^2 | y_1, \dots, y_n) \propto \exp\left(-\alpha\left(\frac{n+n_0}{2\sigma^2}\right) + \left(\frac{n\bar{y} + n_0\mu_0}{\sigma^2}\right)\mu\right)$$

$$\times (\sigma^2)^{-(n+2\alpha+3)/2} \exp\left(-\frac{\sum_{i=1}^n (y_i - \bar{y})^2 + n_0(\mu - \mu_0)^2 + 2\beta}{2\sigma^2}\right)$$

Therefore

$$f(\mu, \sigma^2 | y_1, \dots, y_n) \propto \exp\left(-\frac{\mu^2 - 2\mu\left(\frac{n\bar{y} + n_0\mu_0}{n+n_0}\right)}{2\sigma^2}\right)$$

$$\times (\sigma^2)^{-(n+2\alpha+3)/2} \exp\left(-\frac{\sum_{i=1}^n (y_i - \bar{y})^2 + n_0(\mu - \mu_0)^2 + 2\beta}{2\sigma^2}\right)$$

In other words,

$$f(\mu, \sigma^2 | y_1, \dots, y_n) \propto \exp\left(-\frac{(\mu - \frac{n\bar{y} + n_0\mu_0}{n+n_0})^2}{2\sigma^2}\right)$$

$$\times (\sigma^2)^{-(n+2\alpha+3)/2} \exp\left(-\frac{\sum_{i=1}^n (y_i - \bar{y})^2 + n_0(\mu - \mu_0)^2 + 2\beta}{2\sigma^2} + \frac{(n\bar{y} + n_0\mu_0)^2}{2\sigma^2(n+n_0)}\right)$$

The posterior for σ^2 is

$$f(\sigma^2 | y_1, \dots, y_n) \propto (\sigma^2)^{-(\alpha+2)} \exp\left(-\frac{\sum_{i=1}^n (y_i - \bar{y})^2}{2\sigma^2} + \frac{(n\bar{y} + n_0\mu_0)^2}{2\sigma^2(n+n_0)}\right)$$

$$\sigma^2 | y_1, \dots, y_n \sim IG\left(\alpha + \frac{n}{2}, \beta + \frac{\sum_{i=1}^n (y_i - \bar{y})^2}{2} + \frac{n_0(n\bar{y} + n_0\mu_0)^2}{2(n+n_0)}\right)$$

For the posterior of μ , we have

$$h(\mu | \sigma^2, \bar{y}) = \frac{f(\mu, \sigma^2 | \bar{y})}{f(\sigma^2 | \bar{y})}$$

$$\mu | \sigma^2, \bar{y} \sim N\left(\frac{n\bar{y} + n_0\mu_0}{n+n_0}, \frac{\sigma^2}{n+n_0}\right)$$

$$h(\mu | \bar{y}) = \int_0^\infty h(\mu | \sigma^2, \bar{y}) h(\sigma^2 | \bar{y}) d\sigma^2$$

Simulating Posterior for μ and σ^2 given data

$$\sigma^2 | y_1, \dots, y_n \sim IG\left(\alpha + \frac{n}{2}, \beta + \frac{\sum_{i=1}^n (y_i - \bar{y})^2}{2} + \frac{n_0(n\bar{y} + n_0\mu_0)^2}{2(n+n_0)}\right)$$

$$\mu | \sigma^2, y_1, \dots, y_n \sim N\left(\frac{n\bar{y} + n_0\mu_0}{n+n_0}, \frac{\sigma^2}{n+n_0}\right)$$

We can first simulate σ^2 from the inverse gamma distribution. Then we use it to simulate μ from the normal distribution.

Algorithm:

- Load data y_1, \dots, y_n
- Find \bar{y} , α , and β parameters
- Simulate σ^2 a number of times
- Use σ^2 to simulate μ

We have μ is symmetrically distributed. We can find a credible region the following way:

- Sort the simulated μ values
- Find 2.5 and 97.5 percentiles
- Then we get the credible region

This method is more tricky with σ^2 since it is not symmetrically distributed.

Computational Methods

We always have

$$h(\theta | y) = g(\theta) f(y|\theta) / k$$

$$k = \int g(\theta) f(y|\theta) d\theta$$

We need to calculate the integral numerically.