

There are several competing **interpretations of quantum mechanics** on the market. These interpretations try to make sense out of the theory and explain what the theory tells us about the world.

There are several competing **interpretations of quantum mechanics** on the market. These interpretations try to make sense out of the theory and explain what the theory tells us about the world.

Here are two well-known interpretations:

- ▶ The Copenhagen Interpretation (defenders: N. Bohr and W. Heisenberg)
- ▶ Bohmian Mechanics (defenders: D. Dürr, S. Goldstein and N. Zanghi. Note, however, that the defenders of Bohmian Mechanics do not consider it as an interpretation, but as a theory.)



(Niels Bohr
Image in the public domain)



(Werner Heisenberg
Source: Bundesarchiv, Bild183-R57262 / CC-BY-SA)



(Prof. Dethlef Dürr
used by permission)

There are several competing **interpretations of quantum mechanics** on the market. These interpretations try to make sense out of the theory and explain what the theory tells us about the world.

Here are two well-known interpretations:

- ▶ The Copenhagen Interpretation (defenders: N. Bohr and W. Heisenberg)
- ▶ Bohmian Mechanics (defenders: D. Dürr, S. Goldstein and N. Zanghi. Note, however, that the defenders of Bohmian Mechanics do not consider it as an interpretation, but as a theory.)

Different interpretations give different answers to various philosophical questions, such as the problem of determinism and the interpretation of probability in quantum mechanics.

A **random variable** is a variable that can take on a set of possible values, each with a specific probability.

A **random variable** is a variable that can take on a set of possible values, each with a specific probability.

Examples:

- ▶ the weight of a randomly picked person
- ▶ the spin of an electron

A **random variable** is a variable that can take on a set of possible values, each with a specific probability.

Examples:

- ▶ the weight of a randomly picked person (this is a **continuous variable** with values in the positive real numbers (on a given scale))
- ▶ the spin of an electron (this is a **binary variable** with values $\pm 1/2$)

Probabilities and Correlations

Let us consider two binary random variables A and B with the values: $a_1, b_1 = +1$ and $a_2, b_2 = -1$.

Probabilities and Correlations

Let us consider two binary random variables A and B with the values: $a_1, b_1 = +1$ and $a_2, b_2 = -1$.

To specify the joint probability distribution P , we have to fix the values of

$$\begin{aligned} &P(A = 1, B = 1), \\ &P(A = 1, B = -1), \\ &P(A = -1, B = 1), \\ &P(A = -1, B = -1). \end{aligned}$$

Probabilities and Correlations

Let us consider two binary random variables A and B with the values: $a_1, b_1 = +1$ and $a_2, b_2 = -1$.

To specify the joint probability distribution P , we have to fix the values of

$$P(A = 1, B = 1) =: P(1, 1),$$

$$P(A = 1, B = -1) =: P(1, -1),$$

$$P(A = -1, B = 1) =: P(-1, 1),$$

$$P(A = -1, B = -1) =: P(-1, -1).$$

Probabilities and Correlations

Let us consider two binary random variables A and B with the values: $a_1, b_1 = +1$ and $a_2, b_2 = -1$.

To specify the joint probability distribution P , we have to fix the values of

$$P(A = 1, B = 1) =: P(1, 1),$$

$$P(A = 1, B = -1) =: P(1, -1),$$

$$P(A = -1, B = 1) =: P(-1, 1),$$

$$P(A = -1, B = -1) =: P(-1, -1).$$

Please note:

- ▶ $P(a, b) \equiv P(a \wedge b)$

Probabilities and Correlations

Let us consider two binary random variables A and B with the values: $a_1, b_1 = +1$ and $a_2, b_2 = -1$.

To specify the joint probability distribution P , we have to fix the values of

$$P(A = 1, B = 1) =: P(1, 1),$$

$$P(A = 1, B = -1) =: P(1, -1),$$

$$P(A = -1, B = 1) =: P(-1, 1),$$

$$P(A = -1, B = -1) =: P(-1, -1).$$

Please note:

- ▶ $P(a, b) \equiv P(a \wedge b)$
- ▶ $P(1, 1) + P(1, -1) + P(-1, 1) + P(-1, -1) = 1$

The **expectation** of the random variable A :

$$\begin{aligned} E(A) &= \sum_{i,j=1}^2 a_i \cdot P(a_i, b_j) \\ &= 1 \cdot P(1, 1) + 1 \cdot P(1, -1) + (-1) \cdot P(-1, 1) \\ &\quad + (-1) \cdot P(-1, -1) \\ &= P(1, 1) + P(1, -1) - P(-1, 1) - P(-1, -1) \end{aligned}$$

The **expectation** of the random variable A :

$$\begin{aligned} E(A) &= \sum_{i,j=1}^2 a_i \cdot P(a_i, b_j) \\ &= 1 \cdot P(1, 1) + 1 \cdot P(1, -1) + (-1) \cdot P(-1, 1) \\ &\quad + (-1) \cdot P(-1, -1) \\ &= P(1, 1) + P(1, -1) - P(-1, 1) - P(-1, -1) \end{aligned}$$

The **expectation** of the random variable B :

$$\begin{aligned} E(B) &= \sum_{i,j=1}^2 b_j \cdot P(a_i, b_j) \\ &= 1 \cdot P(1, 1) + (-1) \cdot P(1, -1) + 1 \cdot P(-1, 1) \\ &\quad + (-1) \cdot P(-1, -1) \\ &= P(1, 1) - P(1, -1) + P(-1, 1) - P(-1, -1) \end{aligned}$$

The **expectation** of the random variable AB :

$$\begin{aligned} E(AB) &= \sum_{i,j=1}^2 a_i \cdot b_j \cdot P(a_i, b_j) \\ &= 1 \cdot P(1, 1) + (-1) \cdot P(1, -1) + (-1) \cdot P(-1, 1) \\ &\quad + (-1) \cdot (-1) \cdot P(-1, -1) \\ &= P(1, 1) - P(1, -1) - P(-1, 1) + P(-1, -1) \end{aligned}$$

The **variance** of the random variable A :

$$V(A) = E((A - E(A))^2)$$

The **variance** of the random variable A :

$$\begin{aligned} V(A) &= E((A - E(A))^2) \\ &= E(A^2) - (E(A))^2 \end{aligned}$$

The **covariance** of the random variables A and B :

$$\text{cov}(A, B) = E((A - E(A)) \cdot (B - E(B)))$$

The **covariance** of the random variables A and B :

$$\begin{aligned} cov(A, B) &= E((A - E(A)) \cdot (B - E(B))) \\ &= E(AB) - E(A) \cdot E(B) \end{aligned}$$

If A and B are probabilistically independent, then $P(a_i, b_j) = P(a_i) \cdot P(b_j)$ (remember Lecture 5!).

If A and B are probabilistically independent, then $P(a_i, b_j) = P(a_i) \cdot P(b_j)$ (remember Lecture 5!).

Hence,

$$\begin{aligned} E(AB) &= \sum_{i,j=1}^2 a_i \cdot b_j \cdot P(a_i) P(b_j) \\ E(AB) &= \sum_{i,j=1}^2 a_i P(a_i) \cdot b_j P(b_j) \\ &= \left(\sum_{i=1}^2 a_i P(a_i) \right) \cdot \left(\sum_{j=1}^2 b_j P(b_j) \right) \\ &= E(A) \cdot E(B). \end{aligned}$$

If A and B are probabilistically independent, then $P(a_i, b_j) = P(a_i) \cdot P(b_j)$ (remember Lecture 5!).

Hence,

$$\begin{aligned} E(AB) &= \sum_{i,j=1}^2 a_i \cdot b_j \cdot P(a_i) P(b_j) \\ E(AB) &= \sum_{i,j=1}^2 a_i P(a_i) \cdot b_j P(b_j) \\ &= \left(\sum_{i=1}^2 a_i P(a_i) \right) \cdot \left(\sum_{j=1}^2 b_j P(b_j) \right) \\ &= E(A) \cdot E(B). \end{aligned}$$

Hence, $\text{cov}(A, B) = 0$.

Example

$$P(1, 1) = P(-1, -1) = 0.4$$

$$P(1, -1) = P(-1, 1) = 0.1$$

An Example

$$P(1, 1) = P(-1, -1) = 0.4$$

$$P(1, -1) = P(-1, 1) = 0.1$$

$$P(A = 1) = P(1, 1) + P(1, -1) = 0.5$$

An Example

$$P(1, 1) = P(-1, -1) = 0.4$$

$$P(1, -1) = P(-1, 1) = 0.1$$

$$P(A = 1) = P(1, 1) + P(1, -1) = 0.5$$

$$P(B = 1) = P(1, 1) + P(-1, 1) = 0.5$$

An Example

$$P(1, 1) = P(-1, -1) = 0.4$$

$$P(1, -1) = P(-1, 1) = 0.1$$

$$P(A = 1) = P(1, 1) + P(1, -1) = 0.5$$

$$P(B = 1) = P(1, 1) + P(-1, 1) = 0.5$$

$$P(A = -1) = P(B = -1) = 0.5$$

An Example

$$P(1, 1) = P(-1, -1) = 0.4$$

$$P(1, -1) = P(-1, 1) = 0.1$$

$$P(A = 1) = P(1, 1) + P(1, -1) = 0.5$$

$$P(B = 1) = P(1, 1) + P(-1, 1) = 0.5$$

$$P(A = -1) = P(B = -1) = 0.5$$

$$E(A) = 1 \cdot 0.5 + (-1) \cdot 0.5 = 0$$

$$E(B) = 1 \cdot 0.5 + (-1) \cdot 0.5 = 0$$

An Example

$$\begin{aligned} E(AB) &= \sum_{i,j=1}^2 a_i b_j \cdot P(a_i, b_j) \\ &= 0.4 - 0.1 + 0.4 - 0.1 \\ &= 0.6 \end{aligned}$$

An Example

$$\begin{aligned} E(AB) &= \sum_{i,j=1}^2 a_i b_j \cdot P(a_i, b_j) \\ &= 0.4 - 0.1 + 0.4 - 0.1 \\ &= 0.6 \end{aligned}$$

$$\begin{aligned} \text{cov}(A, B) &= E(AB) - E(A) \cdot E(B) \\ &= E(AB) \\ &= 0.6 \end{aligned}$$

$$E(A) = E(B) = 0, \operatorname{cov}(A, B) = x \in [-1, 1] \rightarrow E(AB) = x$$

$$E(A) = E(B) = 0, \text{cov}(A, B) = x \in [-1, 1] \rightarrow E(AB) = x$$

1. $E(A) = 0$ implies

$$P(1, 1) + P(1, -1) - P(-1, 1) - P(-1, -1) = 0.$$

$$E(A) = E(B) = 0, \text{cov}(A, B) = x \in [-1, 1] \rightarrow E(AB) = x$$

1. $E(A) = 0$ implies

$$P(1, 1) + P(1, -1) - P(-1, 1) - P(-1, -1) = 0.$$

2. $E(B) = 0$ implies

$$P(1, 1) - P(1, -1) + P(-1, 1) - P(-1, -1) = 0.$$

$$E(A) = E(B) = 0, \text{cov}(A, B) = x \in [-1, 1] \rightarrow E(AB) = x$$

1. $E(A) = 0$ implies

$$P(1, 1) + P(1, -1) - P(-1, 1) - P(-1, -1) = 0.$$

2. $E(B) = 0$ implies

$$P(1, 1) - P(1, -1) + P(-1, 1) - P(-1, -1) = 0.$$

3. $E(AB) = x$ implies

$$P(1, 1) - P(1, -1) - P(-1, 1) + P(-1, -1) = x.$$

$$E(A) = E(B) = 0, \text{cov}(A, B) = x \in [-1, 1] \rightarrow E(AB) = x$$

1. $E(A) = 0$ implies

$$P(1, 1) + P(1, -1) - P(-1, 1) - P(-1, -1) = 0.$$

2. $E(B) = 0$ implies

$$P(1, 1) - P(1, -1) + P(-1, 1) - P(-1, -1) = 0.$$

3. $E(AB) = x$ implies

$$P(1, 1) - P(1, -1) - P(-1, 1) + P(-1, -1) = x.$$

4. Probability Theory implies

$$P(1, 1) + P(1, -1) + P(-1, 1) + P(-1, -1) = 1.$$

Solving this system of four linear equations leads to:

$$P(1, 1) = P(-1, -1) = (1 + x)/4$$

$$P(1, -1) = P(-1, 1) = (1 - x)/4$$

Remember: x is the covariance of A and B .

$$E(AB) = E(AB') = E(A'B) = x, E(A'B') = -x, x = 1/\sqrt{2}$$

$$E(AB) = E(AB') = E(A'B) = x, E(A'B') = -x, x = 1/\sqrt{2}$$

$$P(A = 1, B = 1) = P(A = -1, B = -1) = (1 + x)/4$$

$$P(A = 1, B = -1) = P(A = -1, B = 1) = (1 - x)/4$$

$$P(A = 1, B' = 1) = P(A = -1, B' = -1) = (1 + x)/4$$

$$P(A = 1, B' = -1) = P(A = -1, B' = 1) = (1 - x)/4$$

$$P(A' = 1, B = 1) = P(A' = -1, B = -1) = (1 + x)/4$$

$$P(A' = 1, B = -1) = P(A' = -1, B = 1) = (1 - x)/4$$

$$E(AB) = E(AB') = E(A'B) = x, E(A'B') = -x, x = 1/\sqrt{2}$$

$$P(A = 1, B = 1) = P(A = -1, B = -1) = (1 + x)/4$$

$$P(A = 1, B = -1) = P(A = -1, B = 1) = (1 - x)/4$$

$$P(A = 1, B' = 1) = P(A = -1, B' = -1) = (1 + x)/4$$

$$P(A = 1, B' = -1) = P(A = -1, B' = 1) = (1 - x)/4$$

$$P(A' = 1, B = 1) = P(A' = -1, B = -1) = (1 + x)/4$$

$$P(A' = 1, B = -1) = P(A' = -1, B = 1) = (1 - x)/4$$

$$P(A' = 1, B' = 1) = P(A' = -1, B' = -1) = (1 - x)/4$$

$$P(A' = 1, B' = -1) = P(A' = -1, B' = 1) = (1 + x)/4$$

The Clauser-Horne-Shimony-Holt Theorem

There is a joint probability distribution over the random variables A, A', B , and B' if and only if

$$|E(AB) + E(AB') + E(A'B) - E(A'B')| \leq 2.$$

The Clauser-Horne-Shimony-Holt Theorem

There is a joint probability distribution over the random variables A, A', B , and B' if and only if

$$|E(AB) + E(AB') + E(A'B) - E(A'B')| \leq 2.$$

Note: Here are two examples for the **modulus** or **absolute value** of a number:

- ▶ $|5| = 5$
- ▶ $|-5| = 5$

The consequence of the CHSH Theorem

Insert

$$\begin{aligned} E(AB) &= E(AB') = E(A'B) = 1/\sqrt{2} \\ E(A'B') &= -1/\sqrt{2} \end{aligned}$$

The consequence of the CHSH Theorem

Insert

$$\begin{aligned}E(AB) &= E(AB') = E(A'B) = 1/\sqrt{2} \\ E(A'B') &= -1/\sqrt{2}\end{aligned}$$

in the CHSH-inequality:

$$|1/\sqrt{2} + 1/\sqrt{2} + 1/\sqrt{2} + 1/\sqrt{2}| = 2\sqrt{2} \approx 2.828 > 2.$$

The consequence of the CHSH Theorem

Insert

$$\begin{aligned}E(AB) &= E(AB') = E(A'B) = 1/\sqrt{2} \\E(A'B') &= -1/\sqrt{2}\end{aligned}$$

in the CHSH-inequality:

$$|1/\sqrt{2} + 1/\sqrt{2} + 1/\sqrt{2} + 1/\sqrt{2}| = 2\sqrt{2} \approx 2.828 > 2.$$

Hence, there is no joint probability distribution over A, A', B , and B' .

$$E(A) = E(A') = E(B) = E(B') = 0$$

$$E(AB) = E(AB') = E(A'B) = 1/\sqrt{2}$$

$$E(A'B') = -1/\sqrt{2}$$

$$\begin{aligned}
E(A) &= E(A') = E(B) = E(B') = 0 \\
E(AB) &= E(AB') = E(A'B) = 1/\sqrt{2} \\
E(A'B') &= -1/\sqrt{2}
\end{aligned}$$

Some “probabilities”, such as $P(1, 1, -1, 1)$, are negative, but, for example,

$$\begin{aligned}
P(A = 1, B = 1) &= \sum_{A', B'} P(A = 1, A', B = 1, B') \\
&= \frac{1 + \sqrt{2}}{4\sqrt{2}} \geq 0.
\end{aligned}$$

Boolean algebra

Operations:

- ▶ Conjunction: \wedge
- ▶ Disjunction: \vee
- ▶ Negation: \neg

Some laws that hold in a Boolean algebra:

Associativity of \vee : $x \vee (y \vee z) = (x \vee y) \vee z$

Some laws that hold in a Boolean algebra:

Associativity of \vee : $x \vee (y \vee z) = (x \vee y) \vee z$

Associativity of \wedge : $x \wedge (y \wedge z) = (x \wedge y) \wedge z$

Some laws that hold in a Boolean algebra:

Associativity of \vee : $x \vee (y \vee z) = (x \vee y) \vee z$

Associativity of \wedge : $x \wedge (y \wedge z) = (x \wedge y) \wedge z$

Commutativity of \vee : $x \vee y = y \vee x$

Some laws that hold in a Boolean algebra:

Associativity of \vee : $x \vee (y \vee z) = (x \vee y) \vee z$

Associativity of \wedge : $x \wedge (y \wedge z) = (x \wedge y) \wedge z$

Commutativity of \vee : $x \vee y = y \vee x$

Commutativity of \wedge : $x \wedge y = y \wedge x$

Some laws that hold in a Boolean algebra:

Associativity of \vee : $x \vee (y \vee z) = (x \vee y) \vee z$

Associativity of \wedge : $x \wedge (y \wedge z) = (x \wedge y) \wedge z$

Commutativity of \vee : $x \vee y = y \vee x$

Commutativity of \wedge : $x \wedge y = y \wedge x$

Distributivity of \wedge over \vee : $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$

A surprising feature involving spin

- ▶ Electrons can be prepared in a way that their spin has the value $+1/2$ in the z-direction.

A surprising feature involving spin

- ▶ Electrons can be prepared in a way that their spin has the value $+1/2$ in the z -direction.
- ▶ That is, if we measure the spin of electrons which are prepared in this way in the z -direction, then we will always obtain the result $+1/2$.

A surprising feature involving spin

- ▶ Electrons can be prepared in a way that their spin has the value $+1/2$ in the z -direction.
- ▶ That is, if we measure the spin of electrons which are prepared in this way in the z -direction, then we will always obtain the result $+1/2$.
- ▶ If we subsequently measure the spin of such electrons in the orthogonal x -direction, then we will find that 50% of the electrons have spin $+1/2$ and 50% of the electrons have spin $-1/2$.

A surprising feature involving spin

- ▶ Electrons can be prepared in a way that their spin has the value $+1/2$ in the z -direction.
- ▶ That is, if we measure the spin of electrons which are prepared in this way in the z -direction, then we will always obtain the result $+1/2$.
- ▶ If we subsequently measure the spin of such electrons in the orthogonal x -direction, then we will find that 50% of the electrons have spin $+1/2$ and 50% of the electrons have spin $-1/2$.
- ▶ If we next consider the electrons that have spin $+1/2$ in the x -direction and measure their spin in the z -direction again, then we will find that 50% of them have spin $+1/2$ and 50% have spin $-1/2$.

Distributivity is violated!

Let:

- ▶ X : The electron has spin $+1/2$ in the z -direction.

Distributivity is violated!

Let:

- ▶ X : The electron has spin $+1/2$ in the z -direction.
- ▶ Y : The electron has spin $+1/2$ in the x -direction.
- ▶ Z : The electron has spin $-1/2$ in the x -direction.

Distributivity is violated!

Let:

- ▶ X : The electron has spin $+1/2$ in the z -direction.
- ▶ Y : The electron has spin $+1/2$ in the x -direction.
- ▶ Z : The electron has spin $-1/2$ in the x -direction.

We note:

- ▶ $Y \vee Z$ is true (as $+1/2$ and $-1/2$ are the only possible spin-values).

Distributivity is violated!

Let:

- ▶ X : The electron has spin $+1/2$ in the z -direction.
- ▶ Y : The electron has spin $+1/2$ in the x -direction.
- ▶ Z : The electron has spin $-1/2$ in the x -direction.

We note:

- ▶ $Y \vee Z$ is true (as $+1/2$ and $-1/2$ are the only possible spin-values).
- ▶ Hence $X \wedge (Y \vee Z)$ is true.

Distributivity is violated!

Let:

- ▶ X : The electron has spin $+1/2$ in the z -direction.
- ▶ Y : The electron has spin $+1/2$ in the x -direction.
- ▶ Z : The electron has spin $-1/2$ in the x -direction.

We note:

- ▶ $Y \vee Z$ is true (as $+1/2$ and $-1/2$ are the only possible spin-values).
- ▶ Hence $X \wedge (Y \vee Z)$ is true.
- ▶ $(X \wedge Y) \vee (X \wedge Z)$ is false.

Distributivity is violated!

Let:

- ▶ X : The electron has spin $+1/2$ in the z -direction.
- ▶ Y : The electron has spin $+1/2$ in the x -direction.
- ▶ Z : The electron has spin $-1/2$ in the x -direction.

We note:

- ▶ $Y \vee Z$ is true (as $+1/2$ and $-1/2$ are the only possible spin-values).
- ▶ Hence $X \wedge (Y \vee Z)$ is true.
- ▶ $(X \wedge Y) \vee (X \wedge Z)$ is false.
- ▶ Hence, $X \wedge (Y \vee Z) \neq (X \wedge Y) \vee (X \wedge Z)$.



(John von Neumann

Source: The United States Department of Energy, image in the public domain)

A Short History of Quantum Logic

- ▶ 1936: John von Neumann and Garrett Birkhoff published “The Logic of Quantum Mechanics”.

A Short History of Quantum Logic

- ▶ 1936: John von Neumann and Garrett Birkhoff published “The Logic of Quantum Mechanics”.
- ▶ 1960ies and 70ies: George Mackey formulated a set of axioms for the quantum propositional system as a so-called **ortho-complemented lattice**.

A Short History of Quantum Logic

- ▶ 1936: John von Neumann and Garrett Birkhoff published “The Logic of Quantum Mechanics”.
- ▶ 1960ies and 70ies: George Mackey formulated a set of axioms for the quantum propositional system as a so-called **ortho-complemented lattice**.
- ▶ 1964: Constantin Piron proved a representation theorem that aims at motivating the Hilbert-space structure of quantum mechanics from quantum logical axioms.

A Short History of Quantum Logic

- ▶ 1936: John von Neumann and Garrett Birkhoff published “The Logic of Quantum Mechanics”.
- ▶ 1960ies and 70ies: George Mackey formulated a set of axioms for the quantum propositional system as a so-called **ortho-complemented lattice**.
- ▶ 1964: Constantin Piron proved a representation theorem that aims at motivating the Hilbert-space structure of quantum mechanics from quantum logical axioms.
- ▶ Today many quantum logicians explore the connection to the theory of quantum computation and prefer an operational approach.