Rapid one-pot assembly of lentiviral vectors

A. Researcher[1], B. Scientist[1,2] and C. Professor[2]

Abstract. We describe a method for assembling lentiviral transfer vectors in a single reaction. The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium. Transfection efficiency was determined by flow cytometry 48 h after transfection. All experiments were performed in biological triplicates unless stated otherwise.

2 Methods

2.1 Cell culture

HEK293T cells were maintained in DMEM supplemented with 10% fetal bovine serum.[3] The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium. Transfection efficiency was determined by flow cytometry 48 h after transfection. All experiments were performed in biological triplicates unless stated otherwise. Statistical significance was assessed with a two-tailed Student's t-test. The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium. Transfection efficiency was determined by flow cytometry 48 h after transfection.

The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium. Transfection efficiency was determined by flow cytometry 48 h after transfection. All experiments were performed in biological triplicates unless stated otherwise. Statistical significance was assessed with a two-tailed Student's t-test. The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium.

2.2 Lentivirus production

The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium. Transfection efficiency was determined by flow cytometry 48 h after transfection. All experiments were performed in biological triplicates unless stated otherwise. Statistical significance was assessed with a two-tailed Student's t-test. The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium. Transfection efficiency was determined by flow cytometry 48 h after transfection. All experiments were performed in biological triplicates unless stated otherwise. Statistical significance was assessed with a two-tailed Student's t-test.

The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium. Transfection efficiency was determined by flow cytometry 48 h after transfection. All experiments were performed in biological triplicates unless stated otherwise. Statistical significance was assessed with a two-tailed Student's t-test. The cells were washed twice in phosphate-buffered saline and resuspended in fresh medium. Transfection efficiency was determined by flow cytometry 48 h after transfection.