The OEIS is supported by the many generous donors to the OEIS Foundation.

OF INTEGER SEQUENCES ®

founded in 1964 by N. J. A. Sloane

Search

(Greetings from <u>The On-Line Encyclopedia of Integer Sequences!</u>)

A062693 Squarefree n such that the elliptic curve $n*y^2 = x^3 - x$ arising in ⁴ the "congruent number" problem has rank 3.

1254, 2605, 2774, 3502, 4199, 4669, 4895, 6286, 6671, 7230, 7766, 8005, 9015, 9430, 9654, 10199, 10549, 11005, 11029, 12166, 12270, 12534, 12935, 13317, 14965, 15655, 16151, 16206, 16887, 17958, 18221, 19046, 19726, 20005, 20366

(<u>list</u>; <u>graph</u>; <u>refs</u>; <u>listen</u>; <u>history</u>; <u>text</u>; <u>internal format</u>)

OFFSET 0,1

COMMENTS Conjectural, as detailed in the pages from which it is extracted (see the first few links at the web site mentioned for details), but the conjecture is supported by much numerical and theoretical evidence.

LINKS Table of n, a(n) for n=0...34.

A. Dujella, A. S.Janfeda, S. Salami, <u>A Search for High</u>
<u>Rank Congruent Number Elliptic Curves</u>, JIS 12 (2009)
09.5.8.

N. D. Elkies, <u>Algorithmic (a.k.a. Computational) Number Theory: Tables, Links, etc.</u>

Fidel Ronquillo Nemenzo, <u>All congruent numbers less than</u> 40000, Proc. Japan Acad. Ser. A Math. Sci., Volume 74, Number 1 (1998), 29-31. See Table IV p. 31.

PROG (PARI) r(n)=ellanalyticrank(ellinit([0, 0, 0, -n^2, 0]))
[1]

for(n=1, 1e4, if(r(n)==3, print1(n", "))) \\ Charles R
Greathouse IV, Sep 01 2011

CROSSREFS Cf. A062694, A062695.

Sequence in context: <u>A252157</u> <u>A252150</u> <u>A023068</u> * <u>A067203</u>

A230544 A280928

Adjacent sequences: <u>A062690</u> <u>A062691</u> <u>A062692</u> * <u>A062694</u>

<u>A062695</u> <u>A062696</u>

KEYWORD nonn

AUTHOR Noam D. Elkies, Jul 04 2001

STATUS approved

The OEIS Community

Maintained by <u>The OEIS Foundation Inc.</u>

Last modified August 20 12:51 EDT 2025. Contains 386856 sequences.

<u>License Agreements, Terms of Use, Privacy Policy</u>