

Key tropical crops at risk from pollinator biodiversity loss due to climate change and land use

Joseph Millard^{1,2*}, Charlotte L. Outhwaite³, Lynn V. Dicks⁴, Jeff Ollerton⁵, Silvia Ceausu³,
Tim Newbold³

¹ Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

² Leverhulme Centre for Demographic Science, Nuffield College, University of Oxford, Oxford OX1 3UQ, United Kingdom

³ Centre for Biodiversity and Environment Research, Department of Genetics, Evolution and Environment, University College London, London, WC1E 6BT, UK

⁴ Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

⁵ Faculty of Arts, Science & Technology, University of Northampton. University Drive, Northampton, NN1 5PH UK

*Corresponding author. Email: joseph.millard@nhm.ac.uk

One sentence summary

Pollinator abundance losses from agriculture and climate change risk production of key crops in the tropics, especially cocoa.

Short title

Pollinator abundance changes and pollination risk.

Abstract

Recent studies have highlighted rapid ongoing changes in insect biodiversity, in part driven by the interaction of land-use and climate change. Insect pollinator biodiversity specifically has been shown to be changing rapidly, with potential consequences for the provision of crop pollination. The role of land-use-climate interactions in pollinator biodiversity changes remains poorly understood. Here, we present a global assessment of the interactive effects of climate change and land use on pollinator abundance, and predictions of the risk to crop pollination worldwide from the inferred abundance changes. We show that the interactive combination of agricultural land use and recent climate change is associated with particularly large reductions in the abundance of insect pollinators. As a result, it is expected that the tropics will likely experience the greatest risk to crop production from pollinator biodiversity losses. The risk to these crops is highest, and predicted to increase most rapidly, in regions of sub-Saharan Africa and northern South America, especially in areas used to produce cocoa, mango, pumpkin, melon, watermelon, and coffee. The health, well-being, and livelihoods of billions of people rely to some extent upon the availability and affordability of crops dependent on animal pollination. Climate change and agricultural land use could put these people's wellbeing at high risk, if loss of pollinator abundance translates into crop pollination shortfalls.

Introduction

Recent studies have highlighted rapid ongoing changes in terrestrial insect biodiversity (1–3), including among pollinating species (4–8). Some of these studies have reported net declines (1, 3), while others have shown mixtures of gains and losses (2). Pollinator biodiversity changes have potential consequences for the provision of pollination to wild plants and crops. Importantly, evidence for insect biodiversity trends has been biased toward Western Europe and North America (9, 10), with little coverage of tropical and subtropical regions (11, 12). Although a few studies have shown steep declines of insects in the tropics (3), evidence about insect biodiversity trends there often remains anecdotal (13), with global syntheses (1, 2) having strong geographic biases towards non-tropical regions.

Among the drivers of insect and pollinator biodiversity changes, human-driven land-use changes and climate change are prominent (5, 10, 14–17). Climate change, in particular, is emerging as an increasingly important driver (8, 14, 18–21), while synergistic interactive effects of land-use and climate change are associated with further reductions in insect biodiversity compared to if the pressures acted in isolation (22–26). A key mechanism underpinning interactive land use and climate effects is the altered microclimatic conditions in areas where vegetation has been modified for human land use (22). Tropical insects are expected to be more susceptible to climate change, including interactive effects with land use, given their narrower physiological tolerance compared to non-tropical species (27). Indeed, recent studies show greater effects in tropical than non-tropical insect biodiversity (26).

Changes in the biodiversity and composition of pollinator communities are expected to have large effects on the provision of pollination services. Pollen limitation from animal pollinator losses has already been shown to reduce the reproductive success of wild plants (4, 28), and the productivity of certain crops (29–32), although there is no clear evidence that pollen limitation is yet causing wholesale reductions in yields of crops that rely on animal pollination (33). Evidence that insect biodiversity responds to human pressures more strongly in the tropics than elsewhere (17, 26) is noteworthy, given that the majority of animal-pollination-dependent crops are grown there (34). However, it is not only tropical countries that will experience the effects of pollinator losses and subsequent pollen limitation, with high income countries benefiting from imports of animal-pollination-dependent foods from tropical areas (35). Abundance, species diversity, and functional diversity of pollinators have all been implicated as determinants of the delivery of pollination service (32, 36–40). Previous attempts to model the provision of crop pollination service are often based on predictions of pollinator abundance, which bears a direct relation to pollination of plants, and has been shown to give a reasonable approximation of pollen deposition in at least some study systems (41). A key uncertainty however relates to the shape of the functional relationship between pollinator abundance and crop production (41).

Here, we present a global assessment of the interactive effects of climate change and land use on pollinator abundance, and predictions of how the inferred abundance changes might translate into risk to crop pollination worldwide based on a range of possible abundance-pollination relationships. Our underlying analyses are based on the PREDICTS database of biodiversity comparisons in different land uses (42), together with a list of species identified in the literature as likely pollinators (17). We use mixed-effects models to fit total pollinator community abundance as a function of land use (primary vegetation versus cropland) and a measure of historical temperature change between a baseline period (1901–1930) and the year prior to biodiversity sampling, standardized by monthly temperature variability in the baseline period (26). We standardised temperature changes in this way to capture where temperatures have exceeded ordinary seasonal variation, a consideration that has previously been identified as important for insects in general (26). We then apply these models to predict which locations and crops are likely to be exposed to the greatest losses of pollinator abundance, and thus to face the greatest risk of pollination shortfalls. We moderate estimates of risk according to estimates of where crops are grown (43), how dependent these crops are on animal pollination (34), projections of historical and future climate change, and a set of assumptions for the relationship between local pollinator abundance and crop pollination (from linear to highly convex or concave). We focus on future climate projections under two Representative Concentration Pathways (RCP) scenarios: RCP 2.6, which has a multi-model-median predicted 1.5°C increase in global average temperatures by 2100 compared to the pre-industrial climate; and RCP 6.0, which has a multi-model-median predicted 3°C increase in global

85 average temperatures (44). Finally, we combine projected pollination risk with estimates of the trade in
86 pollination-dependent crop production (35), to predict regions of the world that may be vulnerable to the
87 indirect consequences of crop pollination risk via trade connections.

88 Results and Discussion

89 The abundance of insect pollinators responded strongly to the interaction of recent climate change and land
90 use (Figure 1). Within cropland experiencing novel temperatures (Standardized Temperature Anomaly =
91 1), pollinator abundance is 38.9% that in natural habitat that has not experienced temperature increases.
92 The causal mechanism underpinning this interaction is unclear, but the alteration and homogenisation of
93 microclimatic conditions in croplands (45) is likely to be partly responsible. Our results are qualitatively
94 consistent with recent results for a sample of all insects (26), but importantly we show that responses to the
95 interactive effects of climate change and land use are stronger for pollinating than non-pollinating insects
96 (Figure 1). We also show that the strength of the interactive effect varies among taxonomic groups, with
97 the strongest effects seen in dipteran and hymenopteran pollinators (Figure S1). Whether the sensitivity of
98 pollinating insects to the interaction of climate change and land use relates directly to their reliance on floral
99 resources, or to other correlated traits typical of pollinators, is unclear, and a combination of both factors
100 is likely to operate. For example, selection of animal pollinated plants is thought to be highly sensitive to
101 climatic conditions (46), suggesting that plant-pollinator interactions are highly sensitive to thermal changes.
102 Pollinator pilosity (i.e. hairiness), on the other hand, likely affects an insect’s ability to adapt to changes
103 in climate (47), and given its nature as a trait typical of bees and hoverflies, tends to be correlated with a
104 reliance on floral resources (48).




Figure 1: Response of pollinating (A) and non-pollinating (B) insect total abundance to the interactive effect of standardised temperature anomaly and land use (primary vegetation versus cropland) (A: $F = 22.4068$, $p < 0.001$; B: $F = 10.7520$, $p < 0.01$). Note that abundance is plotted on a log_e scale. Results are based on linear mixed-effects models. Site numbers are as follows (also see Table S1): insects known to pollinate (primary vegetation = 1166, cropland = 1507); insects not known to pollinate (primary vegetation = 1747, cropland = 922). See Table S2 for the number of species represented in both the pollinating and non-pollinating groups, and Table S3 for AIC and R^2 values for each model. Shading represents 95% confidence intervals around the mean fitted effect. Green = primary vegetation; yellow = cropland.

105 We predict that total pollination production risk will increase under all climate scenarios (Figure 2). For all
 106 scenarios, our projections use change in pollinating insect abundance as a proxy for the relative risk to the
 107 production of crops dependent on insects, incorporating information on where crops are grown worldwide
 108 (43) as well as the fractional dependence of crops on animal pollination (34). Our estimates of both relative
 109 risk as well as absolute production risk should be interpreted as indices of risk, rather than predictions of the
 110 actual amount of production likely to be lost, given the very high uncertainty in how pollinator abundance
 111 changes will translate into actual production losses. Increases in risk are seen for all assumed relationships
 112 between abundance loss and production risk, although the magnitude of changes in relative risk and especially
 113 absolute production risk varied widely (Figure 2). The predicted rate of increase in average production risk
 114 was substantially higher under RCP (Representative Concentration Pathway) 6.0 than RCP 2.6, suggesting
 115 that efforts to mitigate climate change will reduce the risk to future crop production, alongside the many
 116 other benefits (49). Relative production risk varied strongly between years under the convex abundance-
 117 production relationships (Figure 2). This volatility may be explained by way in which the non-linearity of

118 abundance/production relationships interacts with inter-annual climate variability caused by the El Nino
119 Southern Oscillation (50). While increasingly convex assumed relationships between insect abundance loss
120 and crop production risk led to steeper increases in relative production risk with future climate change,
121 absolute production risk was markedly lower, owing to a lower baseline in the present day (Figure 2). Our
122 estimates of risk are based on the distribution of crops as grown in 2000 (see Table S4 for the full list of
123 crops), meaning we do not account for changes in the distribution of crops over time, which are likely to occur
124 as a result of the direct impacts of climate change, indirect effects through the loss of pollinator biodiversity,
125 and socio-economic factors such as price changes in the global markets for particular crops. The modelled
126 effects of land-use-climate interactions on pollinator abundance are robust to dropping individual taxonomic
127 families from the model dataset (Figure S2), and to using threshold temperatures to restrict the months
128 considered to those in which insects are likely to be active (Figure S3). The projections of crop production
129 risk are robust to variation in climate predictions under different individual climate models (Figure S4), do
130 not change markedly when abundance loss is capped at the maximum model-fitted value (Figure S5), and
131 do not change markedly when tested for sensitivity to the quality of mapped estimates of crop production
132 (Figure S6).




Figure 2: Projected change in total production risk under two RCP scenarios (2.6 and 6.0, see Figure S7 for 8.5) and a set of hypothetical relationships between pollinator abundance and crop production (linear and varying degrees of convexity, see Figure S7 for a set of concave relationships). Results are shown both for an index of change in relative risk (A and B) and for the total production potentially at risk (in tonnes) (C and D). For each year into the future, the standardised temperature anomaly was projected globally for all cells with production of crops dependent on animal pollination, using a 3-year rolling average of temperature anomaly estimates in each cell, from an ensemble of different climate models. We used data on crop production from the year 2000 (the latest year when such data are available for all crops). For each annual projection of standardised temperature anomaly, insect pollinator abundance on cropland was predicted according to the model shown in Figure 1A, and then expressed as proportional abundance loss compared to cropland that has experienced no warming (i.e. standardised temperature anomaly of 0). In each cell, the total production of each crop (43) was multiplied by dependence on animal pollination (34), and then adjusted for the predicted percentage reduction in insect pollinator abundance in that cell. These estimates of crop production at risk were summed across crops, and then across all terrestrial cells globally. Lines show different assumed relationships between insect pollinator abundance and crop production: dashed = convex relationships (of differing degrees, indicated by different colours: yellow = most convex; purple/blue = least convex); and solid = linear relationship.

133 Projected risk to crop production in 2050 from insect pollinator abundance losses, as a proportion of all
134 production in a given location, is highest in the tropical regions of sub-Saharan Africa, South America, and
135 southeast Asia (Figure 3A; see Figure S8 for underlying maps). In terms of total production potentially at
136 risk, China, India, Indonesia, Brazil, and the Philippines emerge as being most at risk (Figure 3B). Among

137 crops, cocoa is estimated to be at highest risk, by a large margin, especially in Africa, followed by mango
138 (particularly in India) and watermelon (notably in China) (Figure 3B). The risk to cocoa production is
139 particularly significant in light of the social and economic context, as most cocoa is produced on small farms
140 (two to four hectares) that provide income to between 40 and 50 million people globally (51). Coffee is also
141 expected to have a combination of relatively high production risk (Figure 3B) and high value, suggesting
142 that regions in which it is grown may experience economic difficulties, unless the pollination service can
143 be replaced cost effectively. Similarly to cocoa, coffee production provides income to millions of small-
144 scale farmers and their families in the tropics (52). Therefore, the increased production risk due to loss
145 of pollinators could lead to increased income insecurity for some of the most vulnerable people globally.
146 Projections of local crop production risk are sensitive to the assumed abundance-production relationship
147 (Figure S9), with the exception of South East Asia, which is consistently projected to have high risk, and
148 the temperate realm, which is consistently projected to have low risk (Figure S10).

149 It is impossible to predict exactly how our estimates of production risk measure will translate into actual
150 crop production losses. There are multiple uncertainties associated with predicting pollinator biodiversity
151 changes and how this impacts crop production, some of which we explore here (e.g. the relationship between
152 pollinator abundance and crop production), but many of which we do not or cannot (e.g. the changing
153 distribution of crops, the economic viability of hand pollination, the buffering effects of managed pollinators,
154 the effects of climate change alone, and other technological solutions such as the breeding or engineering
155 of pollinator-independent cultivars). Regardless, at the global scale, the changes to pollinator biodiversity
156 that we estimate are likely to be reflected in changes in the risk to production of crops dependent on animal
157 pollination. For cocoa, recent research has focused on the direct effects of climate change on crop production
158 (53, 54), often overlooking those that might be pollinator-mediated, probably because the direct effects
159 of climate change are easier to capture. Importantly, solutions to direct and pollinator-mediated effects
160 of climate change may differ. For example, shade trees might protect crops from extreme temperatures
161 (53), but might not for the ceratopogonid fly pollinators on which cocoa pollination depends. The focus in
162 previous research on direct climate impacts rather than pollinator-mediated effects is a key gap given that
163 some cocoa varieties are limited more by pollination availability than resource availability (55).




Figure 3: Projected change globally in crop production estimated to be at risk in 2050 under the RCP 6.0 climate scenario, assuming a linear relationship between insect pollinator abundance loss and production loss for crops dependent on animal pollination. All projections are based on mean projections of the standardized temperature anomaly based on temperature estimates from an ensemble of individual climate models. (A) The sum of crop production at risk across all crops with some dependence on animal pollination, as a proportion of the production of all crops grown in a location ('proportional production risk'). (B) The total crop production at risk for the 7 crops with the highest total pollination dependent production value globally (see Figure S11 for the top 20 crops by pollination dependent production alone, Figure S12 for country level proportional production risk, and Figure S13 for crop level proportional production risk), in million US\$ per annum, broken down into four main geographic regions. Each coloured bar represents a pollination dependent crop group: Grey, cocoa; black, tropical fruits (not recorded elsewhere); red, fruits (not recorded elsewhere); orange, mango; dark blue, coffee; yellow, watermelon; green, coconut; and other crops, light blue. Per tonne values of each crop are for the years 2015-2019 (US\$ in 2015-2019 values) taken from (56), and total pollination dependent production according to (43) and (34).

Countries besides those with high production risk will feel the impact of losses of pollinator biodiversity and the crops that depend on them, through disruption to imports, especially as the most vulnerable crops tend to be valuable export products such as coffee and cocoa. In absolute terms, large countries such as China and the United States have the highest total import risks (Figure 4B). The Netherlands emerges as having surprisingly high risk given its size, the third largest overall import risk (Figure 4B), consistent with its status as the greatest importer of cocoa beans worldwide (57). Import risk per capita (Figure 4A) highlights the challenges that could be faced by nations with limited agricultural production capacity, such as many island countries (e.g. Cayman Islands, Aruba, Singapore, the first, second and fourth highest import risk per capita, respectively) or countries with unfavourable environmental conditions for agriculture (e.g., Mongolia, with the 19th highest import risk per capita). Total import risk per capita tends to be high also in northern and high-income countries, particularly continental western Europe, which has large processing industries for crops such as coffee and cocoa. High income and unfavourable environment for agriculture could both account for high import risk per capita for some countries in the Middle East (United Arab Emirates, Kuwait, Saudi Arabia, which have the 5th, 8th and 27th highest estimated pollination import risk per capita). Our predictions of import risk are based on trade patterns in the present day (35), meaning we do not account for changes in trade flows that will likely occur in the future.




Figure 4: Projected total import risk for 2050 under the RCP 6.0 scenario, assuming a linear relationship between insect pollinator abundance loss and production loss. Projections are based on changes in the mean standardized temperature anomaly based on projections from an ensemble of individual climate models. Import risk is a measure of how the effects of localised production risk might be distributed among other countries, calculated using trade flow data from (35). For each yearly time step into the future, local production risk in each spatial cell is attributed to importers according to the quantity of pollination trade imported from each country. For example, if an importer is dependent on 3 countries for imports, at a proportion of 30%, 50%, and 20%, then any change in import risk should scale as a function of the local production risk aggregated at those same proportions. (A) The geographic distribution of total import risk adjusted for country population size and then converted to a percentile. Colours represent each percentile grouping: yellow, 0-25th percentile; orange, 25-50th; purple, 50-75th; dark blue, 75-100th. Inset plot represents the absolute import risk values on a log10 scale, with the same percentile breakpoints. (B) The total import crop production at risk in thousand tonnes per annum, for the 10 countries with the highest import production at risk.

180 Interacting effects of climate change and anthropogenic land use are rapidly restructuring biodiversity (26,
 181 58). Here we show that the combination of agricultural land use and recent climate change is associated
 182 with particularly large reductions in the abundance of insect pollinators. As a result, we predict that the
 183 tropics will likely experience the greatest risk of future pollination shortfalls, putting at risk the production
 184 of crops that depend on insect pollination. Future pollination risk is estimated to be highest in areas used to
 185 produce cocoa, mango, watermelon, and coffee. Given the many factors that determine crop production and
 186 crop price (59), the likely effects of insect pollinator losses on crop production are unclear, and even if they
 187 do occur, conclusive attribution is likely to be challenging. Such complications likely in part explain why
 188 identifying a strong effect of pollinator losses on global crop yield and price has thus far been so difficult (33,
 189 60, 61). Regardless, there is sufficient evidence to suggest that declining insect pollinator abundance will
 190 influence crop production risk, particularly for those in the global south (62). Such risk could manifest in
 191 the form of direct and immediate losses to crop production through pollinator shortfall, fluctuations in the
 192 stability of production, or through decreased resilience to changes that will happen in conjunction (e.g. the
 193 effects of extreme temperature and drought on crop growth). The health, well-being, and livelihoods of a
 194 high proportion of the global population, from small farmers to consumers, relies to some extent upon the
 195 availability and affordability of crops dependent on animal pollination, which is likely to be put at greater
 196 risk as a result of future pollinator losses as land-use and climate changes intensify.

197 References

- 198 1. R. van Klink, D. E. Bowler, K. B. Gongalsky, A. B. Swengel, A. Gentile, J. M. Chase, Meta-analysis
 199 reveals declines in terrestrial but increases in freshwater insect abundances. *Science*. **368**, 417–420
 (2020).
- 200 2. M. Dornelas, N. J. Gotelli, H. Shimadzu, F. Moyes, A. E. Magurran, B. J. McGill, A balance of
 201 winners and losers in the Anthropocene. *Ecology Letters*. **22**, 847–854 (2019).
- 202 3. B. C. Lister, A. Garcia, Climate-driven declines in arthropod abundance restructure a rainforest food
 203 web. *Proceedings of the National Academy of Sciences*. **115**, E10397–E10406 (2018).
- 204 4. J. C. Biesmeijer, S. P. M. Roberts, M. Reemer, R. Ohlemüller, M. Edwards, T. Peeters, A. P. Schaffers,
 205 S. G. Potts, R. Kleukers, C. D. Thomas, J. Settele, W. E. Kunin, Parallel declines in pollinators and
 insect-pollinated plants in Britain and the Netherlands. *Science*. **313**, 351–354 (2006).
- 206 5. S. G. Potts, J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, W. E. Kunin, Global pollinator
 207 declines: Trends, impacts and drivers. *Trends in Ecology & Evolution*. **25**, 345–353 (2010).
- 208 6. T. H. Oliver, N. J. B. Isaac, T. A. August, B. A. Woodcock, D. B. Roy, J. M. Bullock, Declining
 209 resilience of ecosystem functions under biodiversity loss. *Nature Communications*. **6**, 10122 (2015).
- 210 7. G. D. Powney, C. Carvell, M. Edwards, R. K. A. Morris, H. E. Roy, B. A. Woodcock, N. J. B. Isaac,
 211 Widespread losses of pollinating insects in Britain. *Nature Communications*. **10**, 1018 (2019).
- 212 8. P. Soroye, T. Newbold, J. Kerr, Climate change contributes to widespread declines among bumble
 213 bees across continents. *Science*. **367**, 685–688 (2020).
- 214 9. Y. Basset, G. P. A. Lamarre, Toward a world that values insects. *Science*. **364**, 1230–1231 (2019).
 215
- 216 10. D. L. Wagner, Insect declines in the Anthropocene. *Annual Review of Entomology*. **65**, 457–480
 217 (2020).
- 218 11. S. A. Khan, M. Tanveer, S. Ahmad, M. Mars, M. Naeem, Z. Naveed, W. Schuett, C. Drees, D.
 Goulson, Declining abundance of pollinating insects drives falls in loquat (*Eriobotrya japonica*) fruit
 yields in the Pothwar region of Pakistan. *Agriculture, Ecosystems & Environment*. **339**, 108138
 (2022).

- 220 12. T. M. Lewinsohn, K. Agostini, A. V. Lucci Freitas, A. S. Melo, Insect decline in Brazil: an appraisal
221 of current evidence. *Biology Letters*. **18**, 20220219 (2022).
- 222 13. D. H. Janzen, W. Hallwachs, Perspective: Where might be many tropical insects? *Biological Conser-*
223 *vation*. **233**, 102–108 (2019).
- 224 14. M. S. Warren, J. K. Hill, J. A. Thomas, J. Asher, R. Fox, B. Huntley, D. B. Roy, M. G. Telfer, S.
Jeffcoate, P. Harding, G. Jeffcoate, S. G. Willis, J. N. Greatorex-Davies, D. Moss, C. D. Thomas,
225 Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature*. **414**,
65–69 (2001).
- 226 15. A. De Palma, S. Abrahamczyk, M. A. Aizen, M. Albrecht, Y. Basset, A. Bates, R. J. Blake, C.
Boutin, R. Bugter, S. Connop, L. Cruz-López, S. A. Cunningham, B. Darvill, T. Diekötter, S. Dorn,
N. Downing, M. H. Entling, N. Farwig, A. Felicioli, S. J. Fonte, R. Fowler, M. Franzén, D. Goulson,
I. Grass, M. E. Hanley, S. D. Hendrix, F. Herrmann, F. Herzog, A. Holzschuh, B. Jauker, M. Kessler,
M. E. Knight, A. Kruess, P. Lavelle, V. Le Féon, P. Lentini, L. A. Malone, J. Marshall, E. M. Pachón,
Q. S. McFrederick, C. L. Morales, S. Mudri-Stojnic, G. Nates-Parra, S. G. Nilsson, E. Öckinger,
L. Osgathorpe, A. Parra-H, C. A. Peres, A. S. Persson, T. Petanidou, K. Poveda, E. F. Power,
M. Quaranta, C. Quintero, R. Rader, M. H. Richards, T. Roulston, L. Rousseau, J. P. Sadler, U.
Samnegård, N. A. Schellhorn, C. Schüepp, O. Schweiger, A. H. Smith-Pardo, I. Steffan-Dewenter,
J. C. Stout, R. K. Tonietto, T. Tschardtke, J. M. Tylianakis, H. A. F. Verboven, C. H. Vergara,
J. Verhulst, C. Westphal, H. J. Yoon, A. Purvis, Predicting bee community responses to land-use
227 changes: Effects of geographic and taxonomic biases. *Scientific Reports*. **6**, 31153 (2016).
- 228 16. I. Oliver, J. Dorrough, H. Doherty, N. R. Andrew, Additive and synergistic effects of land cover, land
229 use and climate on insect biodiversity. *Landscape Ecology*. **31**, 2415–2431 (2016).
- 230 17. J. Millard, C. L. Outhwaite, R. Kinnerley, R. Freeman, R. D. Gregory, O. Adedaja, S. Gavini, E.
Kioko, M. Kuhlmann, J. Ollerton, Z.-X. Ren, T. Newbold, Global effects of land-use intensity on local
231 pollinator biodiversity. *Nature Communications*. **12**, 2902 (2021).
- 232 18. R. Hickling, D. B. Roy, J. K. Hill, R. Fox, C. D. Thomas, The distributions of a wide range of
233 taxonomic groups are expanding polewards. *Global Change Biology*. **12**, 450–455 (2006).
- 234 19. B. Martay, M. J. Brewer, D. A. Elston, J. R. Bell, R. Harrington, T. M. Brereton, K. E. Barlow,
M. S. Botham, J. W. Pearce-Higgins, Impacts of climate change on national biodiversity population
235 trends. *Ecography*. **40**, 1139–1151 (2017).
- 236 20. J. Lenoir, R. Bertrand, L. Comte, L. Bourgeaud, T. Hattab, J. Murienne, G. Grenouillet, Species
better track climate warming in the oceans than on land. *Nature Ecology & Evolution*. **4**, 1044–1059
237 (2020).
- 238 21. C. A. Halsch, A. M. Shapiro, J. A. Fordyce, C. C. Nice, J. H. Thorne, D. P. Waetjen, M. L. Forister, In-
sects and recent climate change. *Proceedings of the National Academy of Sciences*. **118**, e2002543117
239 (2021).
- 240 22. A. J. Suggitt, R. J. Wilson, N. J. B. Isaac, C. M. Beale, A. G. Auffret, T. August, J. J. Bennie, H.
Q. P. Crick, S. Duffield, R. Fox, J. J. Hopkins, N. A. Macgregor, M. D. Morecroft, K. J. Walker, I.
M. D. Maclean, Extinction risk from climate change is reduced by microclimatic buffering. *Nature*
241 *Climate Change*. **8**, 713–717 (2018).
- 242 23. P. J. Platts, S. C. Mason, G. Palmer, J. K. Hill, T. H. Oliver, G. D. Powney, R. Fox, C. D. Thomas,
Habitat availability explains variation in climate-driven range shifts across multiple taxonomic groups.
243 *Scientific Reports*. **9**, 15039 (2019).

- 244 24. T. H. Oliver, S. Gillings, J. W. Pearce-Higgins, T. Brereton, H. Q. P. Crick, S. J. Duffield, M. D.
245 Morecroft, D. B. Roy, Large extents of intensive land use limit community reorganization during
climate warming. *Global Change Biology*. **23**, 2272–2283 (2017).
- 246 25. M. K. Peters, A. Hemp, T. Appelhans, J. N. Becker, C. Behler, A. Classen, F. Detsch, A. Ensslin, S.
W. Ferger, S. B. Frederiksen, F. Gebert, F. Gerschlauser, A. Gütlein, M. Helbig-Bonitz, C. Hemp, W.
J. Kindeketa, A. Kühnel, A. V. Mayr, E. Mwangomo, C. Ngereza, H. K. Njovu, I. Otte, H. Pabst,
M. Renner, J. Röder, G. Rutten, D. Schellenberger Costa, N. Sierra-Cornejo, M. G. R. Vollstädt, H.
I. Dulle, C. D. Eardley, K. M. Howell, A. Keller, R. S. Peters, A. Ssymank, V. Kakengi, J. Zhang,
C. Bogner, K. Böhning-Gaese, R. Brandl, D. Hertel, B. Huwe, R. Kiese, M. Kleyer, Y. Kuzyakov, T.
Nauss, M. Schleuning, M. Tschapka, M. Fischer, I. Steffan-Dewenter, Climate–land-use interactions
247 shape tropical mountain biodiversity and ecosystem functions. *Nature*. **568**, 88–92 (2019).
- 248 26. C. L. Outhwaite, P. McCann, T. Newbold, Agriculture and climate change are reshaping insect
249 biodiversity worldwide. *Nature*. **605**, 97–102 (2022).
- 250 27. C. A. Deutsch, J. J. Tewksbury, R. B. Huey, K. S. Sheldon, C. K. Ghalambor, D. C. Haak, P. R.
251 Martin, Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the
National Academy of Sciences*. **105**, 6668–6672 (2008).
- 252 28. J. G. Rodger, J. M. Bennett, M. Razanajatovo, T. M. Knight, M. van Kleunen, T.-L. Ashman, J. A.
Steets, C. Hui, G. Arceo-Gómez, M. Burd, L. A. Burkle, J. H. Burns, W. Durka, L. Freitas, J. E.
Kemp, J. Li, A. Pauw, J. C. Vamosi, M. Wolowski, J. Xia, A. G. Ellis, Widespread vulnerability of
253 flowering plant seed production to pollinator declines. *Science Advances*. **7**, eabd3524 (2021).
- 254 29. A. Klein, I. Steffan-Dewenter, T. Tschardtke, Fruit set of highland coffee increases with the diversity
of pollinating bees. *Proceedings of the Royal Society of London. Series B: Biological Sciences*. **270**,
255 955–961 (2003).
- 256 30. T. H. Ricketts, G. C. Daily, P. R. Ehrlich, C. D. Michener, Economic value of tropical forest to coffee
257 production. *Proceedings of the National Academy of Sciences*. **101**, 12579–12582 (2004).
- 258 31. A. Holzschuh, C. F. Dormann, T. Tschardtke, I. Steffan-Dewenter, Expansion of mass-flowering crops
leads to transient pollinator dilution and reduced wild plant pollination. *Proceedings of the Royal
259 Society B: Biological Sciences*. **278**, 3444–3451 (2011).
- 260 32. B. A. Woodcock, M. P. D. Garratt, G. D. Powney, R. F. Shaw, J. L. Osborne, J. Soroka, S. A. M.
Lindström, D. Stanley, P. Ouyard, M. E. Edwards, F. Jauker, M. E. McCracken, Y. Zou, S. G. Potts,
M. Rundlöf, J. A. Noriega, A. Greenop, H. G. Smith, R. Bommarco, W. van der Werf, J. C. Stout,
I. Steffan-Dewenter, L. Morandin, J. M. Bullock, R. F. Pywell, Meta-analysis reveals that pollinator
261 functional diversity and abundance enhance crop pollination and yield. *Nature Communications*. **10**,
1481 (2019).
- 262 33. M. A. Aizen, L. A. Garibaldi, S. A. Cunningham, A. M. Klein, Long-term global trends in crop yield
and production reveal no current pollination shortage but increasing pollinator dependency. *Current
263 Biology*. **18**, 1572–1575 (2008).
- 264 34. A.-M. Klein, B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C. Kremen, T.
Tschardtke, Importance of pollinators in changing landscapes for world crops. *Proceedings of the
265 Royal Society B: Biological Sciences*. **274**, 303–313 (2007).
- 266 35. F. D. S. Silva, L. G. Carvalheiro, J. Aguirre-Gutiérrez, M. Lucotte, K. Guidoni-Martins, F. Mertens,
Virtual pollination trade uncovers global dependence on biodiversity of developing countries. *Science
267 Advances*. **7**, eabe6636 (2021).
- 268 36. M. Albrecht, B. Schmid, Y. Hautier, C. B. Müller, Diverse pollinator communities enhance plant
269 reproductive success. *Proceedings of the Royal Society B: Biological Sciences*. **279**, 4845–4852 (2012).

- 270 37. C. Brittain, C. Kremen, A.-M. Klein, Biodiversity buffers pollination from changes in environmental
271 conditions. *Global Change Biology*. **19**, 540–547 (2013).
- 272 38. M. A. Genung, J. Fox, N. M. Williams, C. Kremen, J. Ascher, J. Gibbs, R. Winfree, The relative
273 importance of pollinator abundance and species richness for the temporal variance of pollination
services. *Ecology*. **98**, 1807–1816 (2017).
- 274 39. R. Winfree, J. R. Reilly, I. Bartomeus, D. P. Cariveau, N. M. Williams, J. Gibbs, Species turnover
275 promotes the importance of bee diversity for crop pollination at regional scales. *Science*. **359**, 791–793
(2018).
- 276 40. M. Dainese, E. A. Martin, M. A. Aizen, M. Albrecht, I. Bartomeus, R. Bommarco, L. G. Carneiro,
R. Chaplin-Kramer, V. Gagic, L. A. Garibaldi, J. Ghazoul, H. Grab, M. Jonsson, D. S. Karp, C. M.
Kennedy, D. Kleijn, C. Kremen, D. A. Landis, D. K. Letourneau, L. Marini, K. Poveda, R. Rader,
H. G. Smith, T. Tscharntke, G. K. S. Andersson, I. Badenhausser, S. Baensch, A. D. M. Bezerra, F.
J. J. A. Bianchi, V. Boreux, V. Bretagnolle, B. Caballero-Lopez, P. Cavigliasso, A. Četković, N. P.
Chacoff, A. Classen, S. Cusser, F. D. da Silva e Silva, G. A. de Groot, J. H. Dudenhöffer, J. Ekroos,
T. Fijen, P. Franck, B. M. Freitas, M. P. D. Garratt, C. Gratton, J. Hipólito, A. Holzschuh, L.
Hunt, A. L. Iverson, S. Jha, T. Keasar, T. N. Kim, M. Kishinevsky, B. K. Klatt, A.-M. Klein, K. M.
Krewenka, S. Krishnan, A. E. Larsen, C. Lavigne, H. Liere, B. Maas, R. E. Mallinger, E. M. Pachon,
A. Martínez-Salinas, T. D. Meehan, M. G. E. Mitchell, G. A. R. Molina, M. Nesper, L. Nilsson, M.
E. O'Rourke, M. K. Peters, M. Plečáš, S. G. Potts, D. de L. Ramos, J. A. Rosenheim, M. Rundlöf, A.
Rusch, A. Sáez, J. Scheper, M. Schleuning, J. M. Schmack, A. R. Sciligo, C. Seymour, D. A. Stanley,
R. Stewart, J. C. Stout, L. Sutter, M. B. Takada, H. Taki, G. Tamburini, M. Tschumi, B. F. Viana,
C. Westphal, B. K. Willcox, S. D. Wratten, A. Yoshioka, C. Zaragoza-Trello, W. Zhang, Y. Zou,
I. Steffan-Dewenter, A global synthesis reveals biodiversity-mediated benefits for crop production.
277 *Science Advances*. **5**, eaax0121 (2019).
- 278 41. E. Lonsdorf, C. Kremen, T. Ricketts, R. Winfree, N. Williams, S. Greenleaf, Modelling pollination
279 services across agricultural landscapes. *Annals of Botany*. **103**, 1589–1600 (2009).
- 280 42. L. N. Hudson, T. Newbold, S. Contu, S. L. L. Hill, I. Lysenko, A. De Palma, B. Collen, R. M. Ewers,
G. M. Mace, D. W. Purves, J. P. W. Scharlemann, A. Purvis, The database of the PREDICTS
(Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project. *Ecology and*
281 *Evolution*. **7**, 145–188 (2017).
- 282 43. C. Monfreda, N. Ramankutty, J. A. Foley, Farming the planet: 2. Geographic distribution of crop
areas, yields, physiological types, and net primary production in the year 2000. *Global Biogeochemical*
283 *Cycles*. **22**, GB1022 (2008).
- 284 44. J. Rogelj, M. Meinshausen, R. Knutti, Global warming under old and new scenarios using IPCC
285 climate sensitivity range estimates. *Nature Climate Change*. **2**, 248–253 (2012).
- 286 45. T. H. Oliver, M. D. Morecroft, Interactions between climate change and land use change on biodi-
versity: Attribution problems, risks, and opportunities. *Wiley Interdisciplinary Reviews: Climate*
287 *Change*. **5**, 317–335 (2014).
- 288 46. A. R. Rech, B. Dalsgaard, B. Sandel, J. Sonne, J.-C. Svenning, N. Holmes, J. Ollerton, The macroecol-
ogy of animal versus wind pollination: Ecological factors are more important than historical climate
289 stability. *Plant Ecology & Diversity*. **9**, 253–262 (2016).
- 290 47. M. K. Peters, J. Peisker, I. Steffan-Dewenter, B. Hoiss, Morphological traits are linked to the cold
performance and distribution of bees along elevational gradients. *Journal of Biogeography*. **43**, 2040–
291 2049 (2016).
- 292 48. J. R. Stavert, G. Liñán-Cembrano, J. R. Beggs, B. G. Howlett, D. E. Pattemore, I. Bartomeus,
293 Hairiness: The missing link between pollinators and pollination. *PeerJ*. **4**, e2779 (2016).

- 294 49. D. Ürge-Vorsatz, S. T. Herrero, N. K. Dubash, F. Lecocq, Measuring the co-benefits of climate change
295 mitigation. *Annual Review of Environment and Resources*. **39**, 549–582 (2014).
- 296 50. A. Timmermann, S.-I. An, J.-S. Kug, F.-F. Jin, W. Cai, A. Capotondi, K. M. Cobb, M. Lengaigne,
297 M. J. McPhaden, M. F. Stuecker, others, El Niño–southern oscillation complexity. *Nature*. **559**,
535–545 (2018).
- 298 51. M. S. Beg, S. Ahmad, K. Jan, K. Bashir, Status, supply chain and processing of cocoa—a review.
299 *Trends in food science & technology*. **66**, 108–116 (2017).
- 300 52. Y. Pham, K. Reardon-Smith, S. Mushtaq, G. Cockfield, The impact of climate change and variability
301 on coffee production: A systematic review. *Climatic Change*. **156**, 609–630 (2019).
- 302 53. G. Schroth, P. Läderach, A. I. Martinez-Valle, C. Bunn, L. Jassogne, Vulnerability to climate change
303 of cocoa in West Africa: Patterns, opportunities and limits to adaptation. *Science of the Total
Environment*. **556**, 231–241 (2016).
- 304 54. K. Ofori-Boateng, B. Insah, The impact of climate change on cocoa production in West Africa.
305 *International Journal of Climate Change Strategies and Management* (2014).
- 306 55. J. H. Groeneveld, T. Tschardtke, G. Moser, Y. Clough, Experimental evidence for stronger cacao
307 yield limitation by pollination than by plant resources. *Perspectives in Plant Ecology, Evolution and
Systematics*. **12**, 183–191 (2010).
- 308 56. Food and Agriculture Organization of the United Nations (FAO), Producer prices & crops and live-
309 stock products (2022).
- 310 57. A. J. G. Simoes, C. A. Hidalgo, in *Workshops at the twenty-fifth AAAI conference on artificial intel-
311 ligence* (2011).
- 312 58. C. Ganuza, S. Redlich, J. Uhler, C. Tobisch, S. Rojas-Botero, M. K. Peters, J. Zhang, C. S. Benjamin,
313 J. Englmeier, J. Ewald, others, Interactive effects of climate and land use on pollinator diversity differ
among taxa and scales. *Science advances*. **8**, eabm9359 (2022).
- 314 59. S. F. Gaetano, L. Emilia, C. Francesco, N. Gianluca, S. Antonio, Drivers of grain price volatility: A
315 cursory critical review. *Agricultural Economics*. **64**, 347–356 (2018).
- 316 60. M. A. Aizen, L. A. Garibaldi, S. A. Cunningham, A. M. Klein, How much does agriculture depend
317 on pollinators? Lessons from long-term trends in crop production. *Annals of botany*. **103**, 1579–1588
(2009).
- 318 61. L. A. Garibaldi, M. A. Aizen, A. M. Klein, S. A. Cunningham, L. D. Harder, Global growth and sta-
319 bility of agricultural yield decrease with pollinator dependence. *Proceedings of the National Academy
of Sciences*. **108**, 5909–5914 (2011).
- 320 62. L. V. Dicks, T. D. Breeze, H. T. Ngo, D. Senapathi, J. An, M. A. Aizen, P. Basu, D. Buchori, L.
321 Galetto, L. A. Garibaldi, others, A global-scale expert assessment of drivers and risks associated with
pollinator decline. *Nature Ecology & Evolution*. **5**, 1453–1461 (2021).

322 Acknowledgements

323 Thanks to Richard Gregory, Robin Freeman, and colleagues on the GLITRS project for useful discussions
324 and comments. Thanks to Andy Purvis and Melinda Mills, who both provided support and guidance.
325 Thanks also to Opeyemi Adedaja, Sabrina Gavini, Esther Kioko, Michael Kuhlmann, Zong-Xin Ren, and
326 Manu Saunders, who provided input on our set of likely pollinating species, and to all contributors of the
327 PREDICTS database.

328 **Funding**

329 JM received PhD funding from the London NERC DTP and the RSPB. JM, TN and CLO receive
330 funding from the UK Natural Environment Research Council (NERC; grant references: NE/R010811/1,
331 NE/V006533/1 and NE/V006800/1). TN and SC receive funding from the UK Economic and Social
332 Research Council (reference: ES/S008160/1). TN is additionally supported by a University Research
333 Fellowship from the Royal Society.

334 **Author contributions**

335 Conceptualization: JM, TN
336 Methodology: JM, CO, TN, LD, JO
337 Investigation: JM, SC
338 Visualization: JM
339 Supervision: TN
340 Writing – original draft: JM, TN
341 Writing – review & editing: JM, TN, CO, LD, JO, SC

342 **Competing interests**

343 All authors declare that they have no competing interests.

344 **Data and materials availability**

345 All data will be made available on FigShare, and all code on both Zenodo and GitHub.

346 **Supplementary Materials**

347 Materials and Methods
348 Figs. S1-S13
349 Tables 1-4
350 References