

# Key tropical crops at risk from pollinator biodiversity loss due to climate change and land use

Joseph Millard<sup>1,2\*</sup>, Charlotte L. Outhwaite<sup>3</sup>, Lynn V. Dicks<sup>4</sup>, Jeff Ollerton<sup>5</sup>, Silvia Ceausu<sup>3</sup>,  
Tim Newbold<sup>3</sup>

<sup>1</sup> Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

<sup>2</sup> Leverhulme Centre for Demographic Science, Nuffield College, University of Oxford, Oxford OX1 3UQ, United Kingdom

<sup>3</sup> Centre for Biodiversity and Environment Research, Department of Genetics, Evolution and Environment, University College London, London, WC1E 6BT, UK

<sup>4</sup> Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

<sup>5</sup> Faculty of Arts, Science & Technology, University of Northampton. University Drive, Northampton, NN1 5PH UK

\*Corresponding author. Email: joseph.millard@nhm.ac.uk

## One sentence summary

Pollinator abundance losses from agriculture and climate change risk production of key crops in the tropics, especially cocoa.

## Short title

Pollinator abundance changes and pollination risk.

## Abstract

Recent studies have highlighted rapid ongoing changes in insect biodiversity, in part driven by the interaction of land-use and climate change. Insect pollinator biodiversity specifically has been shown to be changing rapidly, with potential consequences for the provision of crop pollination. The role of land-use-climate interactions in pollinator biodiversity changes remains poorly understood. Here, we present a global assessment of the interactive effects of climate change and land use on pollinator abundance, and predictions of the risk to crop pollination worldwide from the inferred abundance changes. We show that the interactive combination of agricultural land use and recent climate change is associated with particularly large reductions in the abundance of insect pollinators. As a result, it is expected that the tropics will likely experience the greatest risk to crop production from pollinator biodiversity losses. The risk to these crops is highest, and predicted to increase most rapidly, in regions of sub-Saharan Africa and northern South America, especially in areas used to produce cocoa, mango, pumpkin, melon, watermelon, and coffee. The health, well-being, and livelihoods of billions of people rely to some extent upon the availability and affordability of crops dependent on animal pollination. Climate change and agricultural land use could put these people's wellbeing at high risk, if loss of pollinator abundance translates into crop pollination shortfalls.

## Introduction

Recent studies have highlighted rapid ongoing changes in terrestrial insect biodiversity (1–3), including among pollinating species (4–8). Some of these studies have reported net declines (1, 3), while others have shown mixtures of gains and losses (2). Pollinator biodiversity changes have potential consequences for the provision of pollination to wild plants and crops. Importantly, evidence for insect biodiversity trends has been biased toward Western Europe and North America (9, 10), with little coverage of tropical and subtropical regions (11, 12). Although a few studies have shown steep declines of insects in the tropics (3), evidence about insect biodiversity trends there often remains anecdotal (13), with global syntheses (1, 2) having strong geographic biases towards non-tropical regions.

Among the drivers of insect and pollinator biodiversity changes, human-driven land-use changes and climate change are prominent (5, 10, 14–17). Climate change, in particular, is emerging as an increasingly important driver (8, 14, 18–21), while synergistic interactive effects of land-use and climate change are associated with further reductions in insect biodiversity compared to if the pressures acted in isolation (22–26). A key mechanism underpinning interactive land use and climate effects is the altered microclimatic conditions in areas where vegetation has been modified for human land use (22). Tropical insects are expected to be more susceptible to climate change, including interactive effects with land use, given their narrower physiological tolerance compared to non-tropical species (27). Indeed, recent studies show greater effects in tropical than non-tropical insect biodiversity (26).

Changes in the biodiversity and composition of pollinator communities are expected to have large effects on the provision of pollination services. Pollen limitation from animal pollinator losses has already been shown to reduce the reproductive success of wild plants (4, 28), and the productivity of certain crops (29–32), although there is no clear evidence that pollen limitation is yet causing wholesale reductions in yields of crops that rely on animal pollination (33). Evidence that insect biodiversity responds to human pressures more strongly in the tropics than elsewhere (17, 26) is noteworthy, given that the majority of animal-pollination-dependent crops are grown there (34). However, it is not only tropical countries that will experience the effects of pollinator losses and subsequent pollen limitation, with high income countries benefiting from imports of animal-pollination-dependent foods from tropical areas (35). Abundance, species diversity, and functional diversity of pollinators have all been implicated as determinants of the delivery of pollination service (32, 36–40). Previous attempts to model the provision of crop pollination service are often based on predictions of pollinator abundance, which bears a direct relation to pollination of plants, and has been shown to give a reasonable approximation of pollen deposition in at least some study systems (41). A key uncertainty however relates to the shape of the functional relationship between pollinator abundance and crop production (41).

Here, we present a global assessment of the interactive effects of climate change and land use on pollinator abundance, and predictions of how the inferred abundance changes might translate into risk to crop pollination worldwide based on a range of possible abundance-pollination relationships. Our underlying analyses are based on the PREDICTS database of biodiversity comparisons in different land uses (42), together with a list of species identified in the literature as likely pollinators (17). We use mixed-effects models to fit total pollinator community abundance as a function of land use (primary vegetation versus cropland) and a measure of historical temperature change between a baseline period (1901–1930) and the year prior to biodiversity sampling, standardized by monthly temperature variability in the baseline period (26). We standardised temperature changes in this way to capture where temperatures have exceeded ordinary seasonal variation, a consideration that has previously been identified as important for insects in general (26). We then apply these models to predict which locations and crops are likely to be exposed to the greatest losses of pollinator abundance, and thus to face the greatest risk of pollination shortfalls. We moderate estimates of risk according to estimates of where crops are grown (43), how dependent these crops are on animal pollination (34), projections of historical and future climate change, and a set of assumptions for the relationship between local pollinator abundance and crop pollination (from linear to highly convex or concave). We focus on future climate projections under two Representative Concentration Pathways (RCP) scenarios: RCP 2.6, which has a multi-model-median predicted 1.5°C increase in global average temperatures by 2100 compared to the pre-industrial climate; and RCP 6.0, which has a multi-model-median predicted 3°C increase in global

85 average temperatures (44). Finally, we combine projected pollination risk with estimates of the trade in  
86 pollination-dependent crop production (35), to predict regions of the world that may be vulnerable to the  
87 indirect consequences of crop pollination risk via trade connections.

## 88 Results and Discussion

89 The abundance of insect pollinators responded strongly to the interaction of recent climate change and land  
90 use (Figure 1). Within cropland experiencing novel temperatures (Standardized Temperature Anomaly =  
91 1), pollinator abundance is 38.9% that in natural habitat that has not experienced temperature increases.  
92 The causal mechanism underpinning this interaction is unclear, but the alteration and homogenisation of  
93 microclimatic conditions in croplands (45) is likely to be partly responsible. Our results are qualitatively  
94 consistent with recent results for a sample of all insects (26), but importantly we show that responses to the  
95 interactive effects of climate change and land use are stronger for pollinating than non-pollinating insects  
96 (Figure 1). We also show that the strength of the interactive effect varies among taxonomic groups, with  
97 the strongest effects seen in dipteran and hymenopteran pollinators (Figure S1). Whether the sensitivity of  
98 pollinating insects to the interaction of climate change and land use relates directly to their reliance on floral  
99 resources, or to other correlated traits typical of pollinators, is unclear, and a combination of both factors  
100 is likely to operate. For example, selection of animal pollinated plants is thought to be highly sensitive to  
101 climatic conditions (46), suggesting that plant-pollinator interactions are highly sensitive to thermal changes.  
102 Pollinator pilosity (i.e. hairiness), on the other hand, likely affects an insect’s ability to adapt to changes  
103 in climate (47), and given its nature as a trait typical of bees and hoverflies, tends to be correlated with a  
104 reliance on floral resources (48).




Figure 1: Response of pollinating (A) and non-pollinating (B) insect total abundance to the interactive effect of standardised temperature anomaly and land use (primary vegetation versus cropland) (A:  $F = 22.4068$ ,  $p < 0.001$ ; B:  $F = 10.7520$ ,  $p < 0.01$ ). Note that abundance is plotted on a  $\log_e$  scale. Results are based on linear mixed-effects models. Site numbers are as follows (also see Table S1): insects known to pollinate (primary vegetation = 1166, cropland = 1507); insects not known to pollinate (primary vegetation = 1747, cropland = 922). See Table S2 for the number of species represented in both the pollinating and non-pollinating groups, and Table S3 for AIC and  $R^2$  values for each model. Shading represents 95% confidence intervals around the mean fitted effect. Green = primary vegetation; yellow = cropland.

105 We predict that total pollination production risk will increase under all climate scenarios (Figure 2). For all  
 106 scenarios, our projections use change in pollinating insect abundance as a proxy for the relative risk to the  
 107 production of crops dependent on insects, incorporating information on where crops are grown worldwide (43)  
 108 as well as the fractional dependence of crops on animal pollination (34). Our estimates of both relative risk as  
 109 well as absolute production risk should be interpreted as indices of risk, rather than predictions of the actual  
 110 amount of production likely to be lost, given the very high uncertainty in how pollinator abundance changes  
 111 will translate into actual production losses (49). Increases in risk are seen for all assumed relationships  
 112 between abundance loss and production risk, although the magnitude of changes in relative risk and especially  
 113 absolute production risk varied widely (Figure 2). The predicted rate of increase in average production risk  
 114 was substantially higher under RCP (Representative Concentration Pathway) 6.0 than RCP 2.6, suggesting  
 115 that efforts to mitigate climate change will reduce the risk to future crop production, alongside the many  
 116 other benefits (50). Relative production risk varied strongly between years under the convex abundance-  
 117 production relationships (Figure 2). This volatility may be explained by way in which the non-linearity of

118 abundance/production relationships interacts with inter-annual climate variability caused by the El Nino  
119 Southern Oscillation (51). While increasingly convex assumed relationships between insect abundance loss  
120 and crop production risk led to steeper increases in relative production risk with future climate change,  
121 absolute production risk was markedly lower, owing to a lower baseline in the present day (Figure 2). Our  
122 estimates of risk are based on the distribution of crops as grown in 2000 (see Table S4 for the full list of  
123 crops), meaning we do not account for changes in the distribution of crops over time, which are likely to occur  
124 as a result of the direct impacts of climate change, indirect effects through the loss of pollinator biodiversity,  
125 and socio-economic factors such as price changes in the global markets for particular crops. The modelled  
126 effects of land-use-climate interactions on pollinator abundance are robust to dropping individual taxonomic  
127 families from the model dataset (Figure S2), and to using threshold temperatures to restrict the months  
128 considered to those in which insects are likely to be active (Figure S3). The projections of crop production  
129 risk are robust to variation in climate predictions under different individual climate models (Figure S4), do  
130 not change markedly when abundance loss is capped at the maximum model-fitted value (Figure S5), and  
131 do not change markedly when tested for sensitivity to the quality of mapped estimates of crop production  
132 (Figure S6).




Figure 2: Projected change in total production risk under two RCP scenarios (2.6 and 6.0, see Figure S7 for 8.5) and a set of hypothetical relationships between pollinator abundance and crop production (linear and varying degrees of convexity, see Figure S7 for a set of concave relationships). Results are shown both for an index of change in relative risk (A and B) and for the total production potentially at risk (in tonnes) (C and D). For each year into the future, the standardised temperature anomaly was projected globally for all cells with production of crops dependent on animal pollination, using a 3-year rolling average of temperature anomaly estimates in each cell, from an ensemble of different climate models. We used data on crop production from the year 2000 (the latest year when such data are available for all crops). For each annual projection of standardised temperature anomaly, insect pollinator abundance on cropland was predicted according to the model shown in Figure 1A, and then expressed as proportional abundance loss compared to cropland that has experienced no warming (i.e. standardised temperature anomaly of 0). In each cell, the total production of each crop (43) was multiplied by dependence on animal pollination (34), and then adjusted for the predicted percentage reduction in insect pollinator abundance in that cell. These estimates of crop production at risk were summed across crops, and then across all terrestrial cells globally. Lines show different assumed relationships between insect pollinator abundance and crop production: dashed = convex relationships (of differing degrees, indicated by different colours: yellow = most convex; purple/blue = least convex); and solid = linear relationship.

133 Projected risk to crop production in 2050 from insect pollinator abundance losses, as a proportion of all  
 134 production in a given location, is highest in the tropical regions of sub-Saharan Africa, South America, and  
 135 southeast Asia (Figure 3A; see Figure S8 for underlying maps). In terms of total production potentially at  
 136 risk, China, India, Indonesia, Brazil, and the Philippines emerge as being most at risk (Figure 3B). Among

137 crops, cocoa is estimated to be at highest risk, by a large margin, especially in Africa, followed by mango  
138 (particularly in India) and watermelon (notably in China) (Figure 3B). The risk to cocoa production is  
139 particularly significant in light of the social and economic context, as most cocoa is produced on small farms  
140 (two to four hectares) that provide income to between 40 and 50 million people globally (52). Coffee is also  
141 expected to have a combination of relatively high production risk (Figure 3B) and high value, suggesting  
142 that regions in which it is grown may experience economic difficulties, unless the pollination service can  
143 be replaced cost effectively. Similarly to cocoa, coffee production provides income to millions of small-  
144 scale farmers and their families in the tropics (53). Therefore, the increased production risk due to loss  
145 of pollinators could lead to increased income insecurity for some of the most vulnerable people globally.  
146 Projections of local crop production risk are sensitive to the assumed abundance-production relationship  
147 (Figure S9), with the exception of South East Asia, which is consistently projected to have high risk, and  
148 the temperate realm, which is consistently projected to have low risk (Figure S10).

149 It is impossible to predict exactly how our estimates of production risk measure will translate into actual  
150 crop production losses. There are multiple uncertainties associated with predicting pollinator biodiversity  
151 changes and how this impacts crop production, some of which we explore here (e.g. the relationship between  
152 pollinator abundance and crop production), but many of which we do not or cannot (e.g. the changing  
153 distribution of crops, the economic viability of hand pollination, the buffering effects of managed pollinators,  
154 the effects of climate change alone, and other technological solutions such as the breeding or engineering  
155 of pollinator-independent cultivars). Regardless, at the global scale, the changes to pollinator biodiversity  
156 that we estimate are likely to be reflected in changes in the risk to production of crops dependent on animal  
157 pollination. For cocoa, recent research has focused on the direct effects of climate change on crop production  
158 (54, 55), often overlooking those that might be pollinator-mediated, probably because the direct effects of  
159 climate change are easier to capture. Importantly, solutions to direct and pollinator-mediated effects of  
160 climate change may differ. For example, shade trees might protect crops from the detrimental effects of  
161 extreme temperatures (54), but might not for the ceratopogonid fly pollinators on which cocoa pollination  
162 depends. The focus in previous research on direct climate impacts rather than pollinator-mediated effects is a  
163 key gap given that some cocoa varieties are limited more by pollination availability than resource availability  
164 (56).




Figure 3: Projected change globally in crop production estimated to be at risk in 2050 under the RCP 6.0 climate scenario, assuming a linear relationship between insect pollinator abundance loss and production loss for crops dependent on animal pollination. All projections are based on mean projections of the standardized temperature anomaly based on temperature estimates from an ensemble of individual climate models. (A) The sum of crop production at risk across all crops with some dependence on animal pollination, as a proportion of the production of all crops grown in a location ('proportional production risk'). (B) The total crop production at risk for the 7 crops with the highest total pollination dependent production value globally (see Figure S11 for the top 20 crops by pollination dependent production alone, Figure S12 for country level proportional production risk, and Figure S13 for crop level proportional production risk), in million US\$ per annum, broken down into four main geographic regions. Each coloured bar represents a pollination dependent crop group: Grey, cocoa; black, tropical fruits (not recorded elsewhere); red, fruits (not recorded elsewhere); orange, mango; dark blue, coffee; yellow, watermelon; green, coconut; and other crops, light blue. Per tonne values of each crop are for the years 2015-2019 (US\$ in 2015-2019 values) taken from (57), and total pollination dependent production according to (43) and (34).



Countries besides those with high production risk will feel the impact of losses of pollinator biodiversity and the crops that depend on them, through disruption to imports, especially as the most vulnerable crops tend to be valuable export products such as coffee and cocoa. In absolute terms, large countries such as China and the United States have the highest total import risks (Figure 4B). The Netherlands emerges as having surprisingly high risk given its size, the third largest overall import risk (Figure 4B), consistent with its status as the greatest importer of cocoa beans worldwide (58). Import risk per capita (Figure 4A) highlights the challenges that could be faced by nations with limited agricultural production capacity, such as many island countries (e.g. Cayman Islands, Aruba, Singapore, the first, second and fourth highest import risk per capita, respectively) or countries with unfavourable environmental conditions for agriculture (e.g., Mongolia, with the 19th highest import risk per capita). Total import risk per capita tends to be high also in northern and high-income countries, particularly continental western Europe, which has large processing industries for crops such as coffee and cocoa. High income and unfavourable environment for agriculture could both account for high import risk per capita for some countries in the Middle East (United Arab Emirates, Kuwait, Saudi Arabia, which have the 5th, 8th and 27th highest estimated pollination import risk per capita). Our predictions of import risk are based on trade patterns in the present day (35), meaning we do not account for changes in trade flows that will likely occur in the future.




Figure 4: Projected total import risk for 2050 under the RCP 6.0 scenario, assuming a linear relationship between insect pollinator abundance loss and production loss. Projections are based on changes in the mean standardized temperature anomaly based on projections from an ensemble of individual climate models. Import risk is a measure of how the effects of localised production risk might be distributed among other countries, calculated using trade flow data from (35). For each yearly time step into the future, local production risk in each spatial cell is attributed to importers according to the quantity of pollination trade imported from each country. For example, if an importer is dependent on 3 countries for imports, at a proportion of 30%, 50%, and 20%, then any change in import risk should scale as a function of the local production risk aggregated at those same proportions. (A) The geographic distribution of total import risk adjusted for country population size and then converted to a percentile. Colours represent each percentile grouping: yellow, 0-25th percentile; orange, 25-50th; purple, 50-75th; dark blue, 75-100th. Inset plot represents the absolute import risk values on a log10 scale, with the same percentile breakpoints. (B) The total import crop production at risk in thousand tonnes per annum, for the 10 countries with the highest import production at risk.

181 Interacting effects of climate change and anthropogenic land use are rapidly restructuring biodiversity (26,  
 182 59). Here we show that the combination of agricultural land use and recent climate change is associated  
 183 with particularly large reductions in the abundance of insect pollinators. As a result, we predict that the  
 184 tropics will likely experience the greatest risk of future pollination shortfalls, putting at risk the production  
 185 of crops that depend on insect pollination. Future pollination risk is estimated to be highest in areas used to  
 186 produce cocoa, mango, watermelon, and coffee. Given the many factors that determine crop production and  
 187 crop price (60), the likely effects of insect pollinator losses on crop production are unclear, and even if they  
 188 do occur, conclusive attribution is likely to be challenging. Such complications likely in part explain why  
 189 identifying a strong effect of pollinator losses on global crop yield and price has thus far been so difficult (33,  
 190 61, 62). Regardless, there is sufficient evidence to suggest that declining insect pollinator abundance will  
 191 influence crop production risk, particularly for those in the global south (63). Such risk could manifest in  
 192 the form of direct and immediate losses to crop production through pollinator shortfall, fluctuations in the  
 193 stability of production, or through decreased resilience to changes that will happen in conjunction (e.g. the  
 194 effects of extreme temperature and drought on crop growth). The health, well-being, and livelihoods of a  
 195 high proportion of the global population, from small farmers to consumers, relies to some extent upon the  
 196 availability and affordability of crops dependent on animal pollination, which is likely to be put at greater  
 197 risk as a result of future pollinator losses as land-use and climate changes intensify.

## 198 References

- 199 1. R. van Klink, D. E. Bowler, K. B. Gongalsky, A. B. Swengel, A. Gentile, J. M. Chase, Meta-analysis  
 200 reveals declines in terrestrial but increases in freshwater insect abundances. *Science*. **368**, 417–420  
 (2020).
- 201 2. M. Dornelas, N. J. Gotelli, H. Shimadzu, F. Moyes, A. E. Magurran, B. J. McGill, A balance of  
 202 winners and losers in the Anthropocene. *Ecology Letters*. **22**, 847–854 (2019).
- 203 3. B. C. Lister, A. Garcia, Climate-driven declines in arthropod abundance restructure a rainforest food  
 204 web. *Proceedings of the National Academy of Sciences*. **115**, E10397–E10406 (2018).
- 205 4. J. C. Biesmeijer, S. P. M. Roberts, M. Reemer, R. Ohlemüller, M. Edwards, T. Peeters, A. P. Schaffers,  
 206 S. G. Potts, R. Kleukers, C. D. Thomas, J. Settele, W. E. Kunin, Parallel declines in pollinators and  
 insect-pollinated plants in Britain and the Netherlands. *Science*. **313**, 351–354 (2006).
- 207 5. S. G. Potts, J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, W. E. Kunin, Global pollinator  
 208 declines: Trends, impacts and drivers. *Trends in Ecology & Evolution*. **25**, 345–353 (2010).
- 209 6. T. H. Oliver, N. J. B. Isaac, T. A. August, B. A. Woodcock, D. B. Roy, J. M. Bullock, Declining  
 210 resilience of ecosystem functions under biodiversity loss. *Nature Communications*. **6**, 10122 (2015).
- 211 7. G. D. Powney, C. Carvell, M. Edwards, R. K. A. Morris, H. E. Roy, B. A. Woodcock, N. J. B. Isaac,  
 212 Widespread losses of pollinating insects in Britain. *Nature Communications*. **10**, 1018 (2019).
- 213 8. P. Soroye, T. Newbold, J. Kerr, Climate change contributes to widespread declines among bumble  
 214 bees across continents. *Science*. **367**, 685–688 (2020).
- 215 9. Y. Basset, G. P. A. Lamarre, Toward a world that values insects. *Science*. **364**, 1230–1231 (2019).  
 216
- 217 10. D. L. Wagner, Insect declines in the Anthropocene. *Annual Review of Entomology*. **65**, 457–480  
 218 (2020).
- 219 11. S. A. Khan, M. Tanveer, S. Ahmad, M. Mars, M. Naeem, Z. Naveed, W. Schuett, C. Drees, D.  
 Goulson, Declining abundance of pollinating insects drives falls in loquat (*Eriobotrya japonica*) fruit  
 yields in the Pothwar region of Pakistan. *Agriculture, Ecosystems & Environment*. **339**, 108138  
 (2022).

220

221 12. T. M. Lewinsohn, K. Agostini, A. V. Lucci Freitas, A. S. Melo, Insect decline in Brazil: an appraisal  
222 of current evidence. *Biology Letters*. **18**, 20220219 (2022).

223 13. D. H. Janzen, W. Hallwachs, Perspective: Where might be many tropical insects? *Biological Conser-*  
224 *vation*. **233**, 102–108 (2019).

225 14. M. S. Warren, J. K. Hill, J. A. Thomas, J. Asher, R. Fox, B. Huntley, D. B. Roy, M. G. Telfer, S.  
Jeffcoate, P. Harding, G. Jeffcoate, S. G. Willis, J. N. Greatorex-Davies, D. Moss, C. D. Thomas,  
226 Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature*. **414**,  
65–69 (2001).

227 15. A. De Palma, S. Abrahamczyk, M. A. Aizen, M. Albrecht, Y. Basset, A. Bates, R. J. Blake, C.  
Boutin, R. Bugter, S. Connop, L. Cruz-López, S. A. Cunningham, B. Darvill, T. Diekötter, S. Dorn,  
N. Downing, M. H. Entling, N. Farwig, A. Felicioli, S. J. Fonte, R. Fowler, M. Franzén, D. Goulson,  
I. Grass, M. E. Hanley, S. D. Hendrix, F. Herrmann, F. Herzog, A. Holzschuh, B. Jauker, M. Kessler,  
M. E. Knight, A. Kruess, P. Lavelle, V. Le Féon, P. Lentini, L. A. Malone, J. Marshall, E. M. Pachón,  
Q. S. McFrederick, C. L. Morales, S. Mudri-Stojnic, G. Nates-Parra, S. G. Nilsson, E. Öckinger,  
L. Osgathorpe, A. Parra-H, C. A. Peres, A. S. Persson, T. Petanidou, K. Poveda, E. F. Power,  
M. Quaranta, C. Quintero, R. Rader, M. H. Richards, T. Roulston, L. Rousseau, J. P. Sadler, U.  
Samnegård, N. A. Schellhorn, C. Schüepp, O. Schweiger, A. H. Smith-Pardo, I. Steffan-Dewenter,  
J. C. Stout, R. K. Tonietto, T. Tschardtke, J. M. Tylianakis, H. A. F. Verboven, C. H. Vergara,  
J. Verhulst, C. Westphal, H. J. Yoon, A. Purvis, Predicting bee community responses to land-use  
228 changes: Effects of geographic and taxonomic biases. *Scientific Reports*. **6**, 31153 (2016).

229 16. I. Oliver, J. Dorrough, H. Doherty, N. R. Andrew, Additive and synergistic effects of land cover, land  
230 use and climate on insect biodiversity. *Landscape Ecology*. **31**, 2415–2431 (2016).

231 17. J. Millard, C. L. Outhwaite, R. Kinnersley, R. Freeman, R. D. Gregory, O. Adedaja, S. Gavini, E.  
Kioko, M. Kuhlmann, J. Ollerton, Z.-X. Ren, T. Newbold, Global effects of land-use intensity on local  
232 pollinator biodiversity. *Nature Communications*. **12**, 2902 (2021).

233 18. R. Hickling, D. B. Roy, J. K. Hill, R. Fox, C. D. Thomas, The distributions of a wide range of  
234 taxonomic groups are expanding polewards. *Global Change Biology*. **12**, 450–455 (2006).

235 19. B. Martay, M. J. Brewer, D. A. Elston, J. R. Bell, R. Harrington, T. M. Brereton, K. E. Barlow,  
M. S. Botham, J. W. Pearce-Higgins, Impacts of climate change on national biodiversity population  
236 trends. *Ecography*. **40**, 1139–1151 (2017).

237 20. J. Lenoir, R. Bertrand, L. Comte, L. Bourgeaud, T. Hattab, J. Murienne, G. Grenouillet, Species  
better track climate warming in the oceans than on land. *Nature Ecology & Evolution*. **4**, 1044–1059  
238 (2020).

239 21. C. A. Halsch, A. M. Shapiro, J. A. Fordyce, C. C. Nice, J. H. Thorne, D. P. Waetjen, M. L. Forister, In-  
sects and recent climate change. *Proceedings of the National Academy of Sciences*. **118**, e2002543117  
240 (2021).

241 22. A. J. Suggitt, R. J. Wilson, N. J. B. Isaac, C. M. Beale, A. G. Auffret, T. August, J. J. Bennie, H.  
Q. P. Crick, S. Duffield, R. Fox, J. J. Hopkins, N. A. Macgregor, M. D. Morecroft, K. J. Walker, I.  
M. D. Maclean, Extinction risk from climate change is reduced by microclimatic buffering. *Nature*  
242 *Climate Change*. **8**, 713–717 (2018).

243 23. P. J. Platts, S. C. Mason, G. Palmer, J. K. Hill, T. H. Oliver, G. D. Powney, R. Fox, C. D. Thomas,  
Habitat availability explains variation in climate-driven range shifts across multiple taxonomic groups.  
244 *Scientific Reports*. **9**, 15039 (2019).

- 245 24. T. H. Oliver, S. Gillings, J. W. Pearce-Higgins, T. Brereton, H. Q. P. Crick, S. J. Duffield, M. D.  
246 Morecroft, D. B. Roy, Large extents of intensive land use limit community reorganization during  
climate warming. *Global Change Biology*. **23**, 2272–2283 (2017).
- 247 25. M. K. Peters, A. Hemp, T. Appelhans, J. N. Becker, C. Behler, A. Classen, F. Detsch, A. Ensslin, S.  
W. Ferger, S. B. Frederiksen, F. Gebert, F. Gerschlaier, A. Gütlein, M. Helbig-Bonitz, C. Hemp, W.  
J. Kindeketa, A. Kühnel, A. V. Mayr, E. Mwangomo, C. Ngereza, H. K. Njovu, I. Otte, H. Pabst,  
M. Renner, J. Röder, G. Rutten, D. Schellenberger Costa, N. Sierra-Cornejo, M. G. R. Vollstädt, H.  
I. Dulle, C. D. Eardley, K. M. Howell, A. Keller, R. S. Peters, A. Ssymank, V. Kakengi, J. Zhang,  
C. Bogner, K. Böhning-Gaese, R. Brandl, D. Hertel, B. Huwe, R. Kiese, M. Kleyer, Y. Kuzyakov, T.  
Nauss, M. Schleuning, M. Tschapka, M. Fischer, I. Steffan-Dewenter, Climate–land-use interactions  
248 shape tropical mountain biodiversity and ecosystem functions. *Nature*. **568**, 88–92 (2019).
- 249 26. C. L. Outhwaite, P. McCann, T. Newbold, Agriculture and climate change are reshaping insect  
250 biodiversity worldwide. *Nature*. **605**, 97–102 (2022).
- 251 27. C. A. Deutsch, J. J. Tewksbury, R. B. Huey, K. S. Sheldon, C. K. Ghalambor, D. C. Haak, P. R.  
252 Martin, Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the  
National Academy of Sciences*. **105**, 6668–6672 (2008).
- 253 28. J. G. Rodger, J. M. Bennett, M. Razanajatovo, T. M. Knight, M. van Kleunen, T.-L. Ashman, J. A.  
Steets, C. Hui, G. Arceo-Gómez, M. Burd, L. A. Burkle, J. H. Burns, W. Durka, L. Freitas, J. E.  
Kemp, J. Li, A. Pauw, J. C. Vamosi, M. Wolowski, J. Xia, A. G. Ellis, Widespread vulnerability of  
254 flowering plant seed production to pollinator declines. *Science Advances*. **7**, eabd3524 (2021).
- 255 29. A. Klein, I. Steffan-Dewenter, T. Tschardtke, Fruit set of highland coffee increases with the diversity  
of pollinating bees. *Proceedings of the Royal Society of London. Series B: Biological Sciences*. **270**,  
256 955–961 (2003).
- 257 30. T. H. Ricketts, G. C. Daily, P. R. Ehrlich, C. D. Michener, Economic value of tropical forest to coffee  
258 production. *Proceedings of the National Academy of Sciences*. **101**, 12579–12582 (2004).
- 259 31. A. Holzschuh, C. F. Dormann, T. Tschardtke, I. Steffan-Dewenter, Expansion of mass-flowering crops  
leads to transient pollinator dilution and reduced wild plant pollination. *Proceedings of the Royal  
260 Society B: Biological Sciences*. **278**, 3444–3451 (2011).
- 261 32. B. A. Woodcock, M. P. D. Garratt, G. D. Powney, R. F. Shaw, J. L. Osborne, J. Soroka, S. A. M.  
Lindström, D. Stanley, P. Ouyard, M. E. Edwards, F. Jauker, M. E. McCracken, Y. Zou, S. G. Potts,  
M. Rundlöf, J. A. Noriega, A. Greenop, H. G. Smith, R. Bommarco, W. van der Werf, J. C. Stout,  
I. Steffan-Dewenter, L. Morandin, J. M. Bullock, R. F. Pywell, Meta-analysis reveals that pollinator  
262 functional diversity and abundance enhance crop pollination and yield. *Nature Communications*. **10**,  
1481 (2019).
- 263 33. M. A. Aizen, L. A. Garibaldi, S. A. Cunningham, A. M. Klein, Long-term global trends in crop yield  
and production reveal no current pollination shortage but increasing pollinator dependency. *Current  
264 Biology*. **18**, 1572–1575 (2008).
- 265 34. A.-M. Klein, B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C. Kremen, T.  
Tschardtke, Importance of pollinators in changing landscapes for world crops. *Proceedings of the  
266 Royal Society B: Biological Sciences*. **274**, 303–313 (2007).
- 267 35. F. D. S. Silva, L. G. Carvalheiro, J. Aguirre-Gutiérrez, M. Lucotte, K. Guidoni-Martins, F. Mertens,  
Virtual pollination trade uncovers global dependence on biodiversity of developing countries. *Science  
268 Advances*. **7**, eabe6636 (2021).
- 269 36. M. Albrecht, B. Schmid, Y. Hautier, C. B. Müller, Diverse pollinator communities enhance plant  
270 reproductive success. *Proceedings of the Royal Society B: Biological Sciences*. **279**, 4845–4852 (2012).

37. C. Brittain, C. Kremen, A.-M. Klein, Biodiversity buffers pollination from changes in environmental conditions. *Global Change Biology*. **19**, 540–547 (2013).
38. M. A. Genung, J. Fox, N. M. Williams, C. Kremen, J. Ascher, J. Gibbs, R. Winfree, The relative importance of pollinator abundance and species richness for the temporal variance of pollination services. *Ecology*. **98**, 1807–1816 (2017).
39. R. Winfree, J. R. Reilly, I. Bartomeus, D. P. Cariveau, N. M. Williams, J. Gibbs, Species turnover promotes the importance of bee diversity for crop pollination at regional scales. *Science*. **359**, 791–793 (2018).
40. M. Dainese, E. A. Martin, M. A. Aizen, M. Albrecht, I. Bartomeus, R. Bommarco, L. G. Carvalheiro, R. Chaplin-Kramer, V. Gagic, L. A. Garibaldi, J. Ghazoul, H. Grab, M. Jonsson, D. S. Karp, C. M. Kennedy, D. Kleijn, C. Kremen, D. A. Landis, D. K. Letourneau, L. Marini, K. Poveda, R. Rader, H. G. Smith, T. Tscharntke, G. K. S. Andersson, I. Badenhausser, S. Baensch, A. D. M. Bezerra, F. J. J. A. Bianchi, V. Boreux, V. Bretagnolle, B. Caballero-Lopez, P. Cavigliasso, A. Četković, N. P. Chacoff, A. Classen, S. Cusser, F. D. da Silva e Silva, G. A. de Groot, J. H. Dudenhöffer, J. Ekroos, T. Fijen, P. Franck, B. M. Freitas, M. P. D. Garratt, C. Gratton, J. Hipólito, A. Holzschuh, L. Hunt, A. L. Iverson, S. Jha, T. Keasar, T. N. Kim, M. Kishinevsky, B. K. Klatt, A.-M. Klein, K. M. Krewenka, S. Krishnan, A. E. Larsen, C. Lavigne, H. Liere, B. Maas, R. E. Mallinger, E. M. Pachon, A. Martínez-Salinas, T. D. Meehan, M. G. E. Mitchell, G. A. R. Molina, M. Nesper, L. Nilsson, M. E. O'Rourke, M. K. Peters, M. Plečáš, S. G. Potts, D. de L. Ramos, J. A. Rosenheim, M. Rundlöf, A. Rusch, A. Sáez, J. Scheper, M. Schleuning, J. M. Schmack, A. R. Sciligo, C. Seymour, D. A. Stanley, R. Stewart, J. C. Stout, L. Sutter, M. B. Takada, H. Taki, G. Tamburini, M. Tschumi, B. F. Viana, C. Westphal, B. K. Willcox, S. D. Wratten, A. Yoshioka, C. Zaragoza-Trello, W. Zhang, Y. Zou, I. Steffan-Dewenter, A global synthesis reveals biodiversity-mediated benefits for crop production. *Science Advances*. **5**, eaax0121 (2019).
41. E. Lonsdorf, C. Kremen, T. Ricketts, R. Winfree, N. Williams, S. Greenleaf, Modelling pollination services across agricultural landscapes. *Annals of Botany*. **103**, 1589–1600 (2009).
42. L. N. Hudson, T. Newbold, S. Contu, S. L. L. Hill, I. Lysenko, A. De Palma, B. Collen, R. M. Ewers, G. M. Mace, D. W. Purves, J. P. W. Scharlemann, A. Purvis, The database of the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project. *Ecology and Evolution*. **7**, 145–188 (2017).
43. C. Monfreda, N. Ramankutty, J. A. Foley, Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000. *Global Biogeochemical Cycles*. **22**, GB1022 (2008).
44. J. Rogelj, M. Meinshausen, R. Knutti, Global warming under old and new scenarios using IPCC climate sensitivity range estimates. *Nature Climate Change*. **2**, 248–253 (2012).
45. T. H. Oliver, M. D. Morecroft, Interactions between climate change and land use change on biodiversity: Attribution problems, risks, and opportunities. *Wiley Interdisciplinary Reviews: Climate Change*. **5**, 317–335 (2014).
46. A. R. Rech, B. Dalsgaard, B. Sandel, J. Sonne, J.-C. Svenning, N. Holmes, J. Ollerton, The macroecology of animal versus wind pollination: Ecological factors are more important than historical climate stability. *Plant Ecology & Diversity*. **9**, 253–262 (2016).
47. M. K. Peters, J. Peisker, I. Steffan-Dewenter, B. Hoiss, Morphological traits are linked to the cold performance and distribution of bees along elevational gradients. *Journal of Biogeography*. **43**, 2040–2049 (2016).
48. J. R. Stavert, G. Liñán-Cembrano, J. R. Beggs, B. G. Howlett, D. E. Pattemore, I. Bartomeus, Hairiness: The missing link between pollinators and pollination. *PeerJ*. **4**, e2779 (2016).

- 295 49. M. A. Aizen, L. A. Garibaldi, L. D. Harder, Myth and reality of a global crisis for agricultural  
296 pollination. *Asociacion Argentina de Ecologia* (2022).
- 297 50. D. Ürge-Vorsatz, S. T. Herrero, N. K. Dubash, F. Lecocq, Measuring the co-benefits of climate change  
298 mitigation. *Annual Review of Environment and Resources*. **39**, 549–582 (2014).
- 299 51. A. Timmermann, S.-I. An, J.-S. Kug, F.-F. Jin, W. Cai, A. Capotondi, K. M. Cobb, M. Lengaigne,  
300 M. J. McPhaden, M. F. Stuecker, others, El Niño–southern oscillation complexity. *Nature*. **559**,  
535–545 (2018).
- 301 52. M. S. Beg, S. Ahmad, K. Jan, K. Bashir, Status, supply chain and processing of cocoa-a review.  
302 *Trends in food science & technology*. **66**, 108–116 (2017).
- 303 53. Y. Pham, K. Reardon-Smith, S. Mushtaq, G. Cockfield, The impact of climate change and variability  
304 on coffee production: A systematic review. *Climatic Change*. **156**, 609–630 (2019).
- 305 54. G. Schroth, P. Läderach, A. I. Martinez-Valle, C. Bunn, L. Jassogne, Vulnerability to climate change  
306 of cocoa in West Africa: Patterns, opportunities and limits to adaptation. *Science of the Total  
Environment*. **556**, 231–241 (2016).
- 307 55. K. Ofori-Boateng, B. Insah, The impact of climate change on cocoa production in West Africa.  
308 *International Journal of Climate Change Strategies and Management* (2014).
- 309 56. J. H. Groeneveld, T. Tschardtke, G. Moser, Y. Clough, Experimental evidence for stronger cacao  
310 yield limitation by pollination than by plant resources. *Perspectives in Plant Ecology, Evolution and  
Systematics*. **12**, 183–191 (2010).
- 311 57. Food and Agriculture Organization of the United Nations (FAO), Producer prices & crops and live-  
312 stock products (2022).
- 313 58. A. J. G. Simoes, C. A. Hidalgo, in *Workshops at the twenty-fifth AAAI conference on artificial intel-  
314 ligence* (2011).
- 315 59. C. Ganuza, S. Redlich, J. Uhler, C. Tobisch, S. Rojas-Botero, M. K. Peters, J. Zhang, C. S. Benjamin,  
316 J. Englmeier, J. Ewald, others, Interactive effects of climate and land use on pollinator diversity differ  
among taxa and scales. *Science advances*. **8**, eabm9359 (2022).
- 317 60. S. F. Gaetano, L. Emilia, C. Francesco, N. Gianluca, S. Antonio, Drivers of grain price volatility: A  
318 cursory critical review. *Agricultural Economics*. **64**, 347–356 (2018).
- 319 61. M. A. Aizen, L. A. Garibaldi, S. A. Cunningham, A. M. Klein, How much does agriculture depend  
320 on pollinators? Lessons from long-term trends in crop production. *Annals of botany*. **103**, 1579–1588  
(2009).
- 321 62. L. A. Garibaldi, M. A. Aizen, A. M. Klein, S. A. Cunningham, L. D. Harder, Global growth and sta-  
322 bility of agricultural yield decrease with pollinator dependence. *Proceedings of the National Academy  
of Sciences*. **108**, 5909–5914 (2011).
- 323 63. L. V. Dicks, T. D. Breeze, H. T. Ngo, D. Senapathi, J. An, M. A. Aizen, P. Basu, D. Buchori, L.  
324 Galetto, L. A. Garibaldi, others, A global-scale expert assessment of drivers and risks associated with  
pollinator decline. *Nature Ecology & Evolution*. **5**, 1453–1461 (2021).

## 325 Acknowledgements

326 Thanks to Richard Gregory, Robin Freeman, and colleagues on the GLITRS project for useful discussions  
327 and comments. Thanks to Andy Purvis and Melinda Mills, who both provided support and guidance.

328 Thanks also to Opeyemi Adedaja, Sabrina Gavini, Esther Kioko, Michael Kuhlmann, Zong-Xin Ren, and  
329 Manu Saunders, who provided input on our set of likely pollinating species, and to all contributors of the  
330 PREDICTS database.

## 331 **Funding**

332 JM received PhD funding from the London NERC DTP and the RSPB. JM, TN and CLO receive  
333 funding from the UK Natural Environment Research Council (NERC; grant references: NE/R010811/1,  
334 NE/V006533/1 and NE/V006800/1). TN and SC receive funding from the UK Economic and Social  
335 Research Council (reference: ES/S008160/1). TN is additionally supported by a University Research  
336 Fellowship from the Royal Society.

## 337 **Author contributions**

338 Conceptualization: JM, TN  
339 Data curation: JM  
340 Formal analysis: JM  
341 Methodology: JM, CO, TN, LD, JO  
342 Investigation: JM, SC  
343 Visualization: JM  
344 Supervision: TN  
345 Writing – original draft: JM, TN  
346 Writing – review & editing: JM, TN, CO, LD, JO, SC

## 347 **Competing interests**

348 All authors declare that they have no competing interests.

## 349 **Data and materials availability**

350 All data will be made available on FigShare, and all code on both Zenodo and GitHub.

## 351 **Supplementary Materials**

352 Materials and Methods  
353 Figs. S1-S13  
354 Tables 1-4  
355 References