

Key tropical crops at risk from pollinator biodiversity loss due to climate change and land use

Joseph Millard^{1,2*}, Charlotte L. Outhwaite³, Silvia Ceașu³, Luísa G. Carneiro^{4,5},
Felipe Deodato da Silva e Silva⁶, Lynn V. Dicks⁷, Jeff Ollerton⁸, Tim Newbold³

¹ Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

² Leverhulme Centre for Demographic Science and Nuffield College, University of Oxford, Oxford OX1 3UQ, United Kingdom

³ Centre for Biodiversity and Environment Research, Department of Genetics, Evolution and Environment, University College London, London, WC1E 6BT, UK

⁴ Department of Ecology, Federal University of Goiás, Goiânia-GO, 74690-900, Brazil.

⁵ Center for Ecology, Evolution and Environmental Change (CE3C), University of Lisbon, Lisbon, Portugal.

⁶ Federal Institute of Education, Science and Technology of Mato Grosso (IFMT)—Campus Barra do Garças, Barra do Garças-MT, 78600-000, Brazil.

⁷ Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

⁸ Faculty of Arts, Science & Technology, University of Northampton. University Drive, Northampton, NN1 5PH UK

*Corresponding author. Email: joseph.millard@nhm.ac.uk

One sentence summary

Pollinator abundance losses from agriculture and climate change risk production of key crops in the tropics, especially cocoa.

Short title

Pollinator abundance changes and pollination risk.

Abstract

Recent studies have highlighted rapid ongoing changes in insect biodiversity, in part driven by the interaction of land-use and climate change. Insect pollinator biodiversity specifically has been shown to be changing rapidly, with potential consequences for the provision of crop pollination. The role of land-use-climate interactions in pollinator biodiversity changes remains poorly understood. Here, we present a global assessment of the interactive effects of climate change and land use on pollinator abundance, and predictions of the risk to crop pollination worldwide from the inferred abundance changes. We show that the interactive combination of agricultural land use and recent climate change is associated with particularly large reductions in

the abundance of insect pollinators. As a result, it is expected that the tropics will likely experience the greatest risk to crop production from pollinator biodiversity losses. The risk to these crops is highest, and predicted to increase most rapidly, in regions of sub-Saharan Africa and northern South America, especially in areas used to produce cocoa, mango, pumpkin, melon, watermelon, and coffee. The health, well-being, and livelihoods of billions of people rely to some extent upon the availability and affordability of crops dependent on animal pollination. Climate change and agricultural land use could put these people’s wellbeing at high risk, if loss of pollinator abundance translates into crop pollination shortfalls.

Introduction

Recent studies have highlighted rapid ongoing changes in terrestrial insect biodiversity,^{1–3} including among pollinating species.^{4–8} Some of these studies have reported net declines,^{1,3} while others have shown mixtures of gains and losses.² Pollinator biodiversity changes have potential consequences for the provision of pollination to wild plants and crops. Importantly, evidence for insect biodiversity trends has been biased toward Western Europe and North America,^{9,10} with little coverage of tropical and subtropical regions.^{11,12} Although a few studies have shown steep declines of insects in the tropics,³ evidence about insect biodiversity trends there often remains anecdotal,¹³ with global syntheses^{1,2} having strong geographic biases towards non-tropical regions.

Among the drivers of insect and pollinator biodiversity changes, human-driven land-use changes and climate change are prominent.^{5,10,14–17} Climate change, in particular, is emerging as an increasingly important driver,^{8,14,18–21} while synergistic interactive effects of land-use and climate change are associated with further reductions in insect biodiversity compared to if the pressures acted in isolation.^{22–26} A key mechanism underpinning interactive land use and climate effects is the altered microclimatic conditions in areas where vegetation has been modified for human land use.²² Tropical insects are expected to be more susceptible to climate change, including interactive effects with land use, given their narrower physiological tolerance compared to non-tropical species.²⁷ Indeed, recent studies show greater effects in tropical than non-tropical insect biodiversity.²⁶

Changes in the biodiversity and composition of pollinator communities are expected to have large effects on the provision of pollination services. Pollen limitation from animal pollinator losses has already been shown to reduce the reproductive success of wild plants,^{4,28} and the productivity of certain crops,^{29–32} although there is no clear evidence that pollen limitation is yet causing wholesale reductions in yields of crops that rely on animal pollination.³³ Evidence that insect biodiversity responds to human pressures more strongly in the tropics than elsewhere^{17,26} is noteworthy, given that the majority of animal-pollination-dependent crops are grown in the tropics.³⁴ However, it is not only tropical countries that will experience the effects of pollinator losses and subsequent pollen limitation, with high income countries benefiting from imports of animal-pollination-dependent foods from tropical areas.³⁵ Abundance, species diversity, and functional diversity of pollinators have all been implicated as determinants of the delivery of pollination service.^{32,36–40} Previous attempts to model the provision of crop pollination service are often based on predictions of pollinator abundance, which bears a direct relation to pollination of plants, and has been shown to give a reasonable approximation of pollen deposition in at least some study systems.⁴¹ A key uncertainty however relates to the shape of the functional relationship between pollinator abundance and crop production.⁴¹

Here, we present a global assessment of the interactive effects of climate change and land use on pollinator abundance, and predictions of how the inferred abundance changes might translate into risk to crop pollination worldwide based on a range of possible abundance-pollination relationships. Our underlying analyses are based on the PREDICTS database of biodiversity recorded in different land uses,⁴² together with a list of species identified in the literature as likely pollinators.¹⁷ We use mixed-effects models to fit total pollinator community abundance as a function of land use (primary vegetation versus cropland) and a measure of historical temperature change between a baseline period (1901–1930) and the year prior to biodiversity sampling, standardized by monthly temperature variability in the baseline period.²⁶ We standardised temperature changes in this way to capture where temperatures have exceeded ordinary seasonal variation, a consideration that has previously been identified as important for insects in general.²⁶ We then apply these

models to predict which locations and crops are likely to be exposed to the greatest losses of pollinator abundance, and thus to face the greatest risk of crop pollination shortfalls. We moderate estimates of risk according to estimates of where crops are grown,⁴³ how dependent these crops are on animal pollination,³⁴ projections of historical and future climate change, and a set of assumptions for the relationship between local pollinator abundance and crop pollination (from linear to highly convex or concave). We focus on future climate projections under two Representative Concentration Pathways (RCP) scenarios: RCP 2.6, which has a multi-model-median predicted 1.5°C increase in global average temperatures by 2100 compared to the pre-industrial climate; and RCP 6.0, which has a multi-model-median predicted 3°C increase in global average temperatures.⁴⁴ Finally, we combine projected pollination risk with estimates of the trade in pollination-dependent crop production,³⁵ to predict regions of the world that may be vulnerable to the indirect consequences of crop pollination risk via trade connections.

Results and Discussion

The abundance of insect pollinators responded strongly to the interaction of recent climate change and land use (Figure 1). Within cropland experiencing novel temperatures (Standardized Temperature Anomaly = 1), pollinator abundance is 38.9% that in natural habitat that has not experienced temperature increases. The causal mechanism underpinning this interaction is unclear, but the moderation and homogenisation of microclimatic conditions in croplands⁴⁵ is likely to be partly responsible. Our results are qualitatively consistent with recent results for a sample of all insects,²⁶ but importantly we show that responses to the interactive effects of climate change and land use are stronger for pollinating than non-pollinating insects (Figure 1). We also show that the strength of the interactive effect varies among taxonomic groups, with the strongest effects seen in dipteran and hymenopteran pollinators (Figure S1). Whether the sensitivity of pollinating insects to the interaction of climate change and land use relates directly to their reliance on floral resources, or to other correlated traits typical of pollinators, is unclear, and a combination of both factors is likely to operate. For example, selection of animal pollinated plants is thought to be highly sensitive to climatic conditions,⁴⁶ suggesting that plant-pollinator interactions are highly sensitive to thermal changes. Pollinator pilosity (i.e. hairiness), on the other hand, likely affects an insect's ability to adapt to changes in climate,⁴⁷ and given its nature as a trait typical of bees and hoverflies, tends to be correlated with a reliance on floral resources.⁴⁸




Figure 1: Response of pollinating (A) and non-pollinating (B) insect total abundance to the interactive effect of standardised temperature anomaly and land use (primary vegetation versus cropland) (A: $F = 22.4068$, $p < 0.001$; B: $F = 10.7520$, $p < 0.01$). Note that abundance is plotted on a log_e scale. Results are based on linear mixed-effects models. Site numbers are as follows (also see Table S1 and Figure S14 for site spatial distribution): insects known to pollinate (primary vegetation = 1166, cropland = 1507); insects not known to pollinate (primary vegetation = 1747, cropland = 922). See Table S2 for the number of species represented in both the pollinating and non-pollinating groups, Table S3 for AIC and R^2 values for each model, and Figures S18 and S19 for the number of insect pollinating species. Shading represents 95% confidence intervals around the mean fitted effect. Green = primary vegetation; yellow = cropland.

108 We predict that total pollination production risk will increase under all climate scenarios (Figure 2). For all
109 scenarios, our projections use change in pollinating insect abundance as a proxy for the relative risk to the
110 production of crops dependent on insects, incorporating information on where crops are grown worldwide⁴³
111 as well as the fractional dependence of crops on animal pollination.³⁴ Our estimates of both relative risk
112 as well as absolute production risk should be interpreted as indices of risk, rather than predictions of the
113 actual amount of production likely to be lost, given the very high uncertainty in how pollinator abundance
114 changes will translate into actual production losses.⁴⁹ Increases in risk are seen for all assumed relationships
115 between abundance loss and production risk, although the magnitude of changes in relative risk and espe-
116 cially absolute production risk varied widely (Figure 2). The predicted rate of increase in average production
117 risk was substantially higher under RCP (Representative Concentration Pathway) 6.0 than RCP 2.6, sug-
118 gesting that efforts to mitigate climate change will reduce the risk to future crop production, alongside the
119 many other benefits.⁵⁰ Relative production risk varied strongly between years under the convex abundance-

120 production relationships (Figure 2). This volatility may be explained by way in which the non-linearity of
121 abundance/production relationships interacts with inter-annual climate variability caused by the El Nino
122 Southern Oscillation.⁵¹ While increasingly convex assumed relationships between insect abundance loss and
123 crop production risk led to steeper increases in relative production risk with future climate change, absolute
124 production risk was markedly lower, owing to a lower baseline in the present day (Figure 2). Our estimates
125 of risk are based on the distribution of crops as grown in 2000 (see Table S4 for the full list of crops),
126 meaning we do not account for changes in the distribution of crops over time, which are likely to occur as
127 a result of the direct impacts of climate change, indirect effects through the loss of pollinator biodiversity,
128 and socio-economic factors such as price changes in the global markets for particular crops. The modelled
129 effects of land-use-climate interactions on pollinator abundance are robust to dropping individual taxonomic
130 families from the model dataset (Figure S2), and to using threshold temperatures to restrict the months
131 considered to those in which insects are likely to be active (Figure S3), and likely hold for a causal inference
132 fixed effects approach (Figure S21), inspired by.⁵² The projections of crop production risk are robust to vari-
133 ation in climate predictions under different individual climate models (Figure S4), do not change markedly
134 when abundance loss is capped at the maximum model-fitted value (Figure S5), do not change markedly
135 when tested for sensitivity to the quality of mapped estimates of crop production (Figure S6), do not change
136 markedly when based on a measure of Chao species richness (Figures S15 and S16), or for risk estimated on
137 the basis of change in bee abundance alone (Figure S17).




Figure 2: Projected change in total production risk under two RCP scenarios (2.6 and 6.0, see Figure S7 for 8.5) and a set of hypothetical relationships between pollinator abundance and crop production (linear and varying degrees of convexity, see Figure S7 for a set of concave relationships). Results are shown both for an index of change in relative risk (A and B) and for the total production potentially at risk (in tonnes) (C and D). For each year into the future, the standardised temperature anomaly was projected globally for all cells with production of crops dependent on animal pollination, using a 3-year rolling average of temperature anomaly estimates in each cell, from an ensemble of different climate models. We used data on crop production from the year 2000 (the latest year when such data are available for all crops). For each annual projection of standardised temperature anomaly, insect pollinator abundance on cropland was predicted according to the model shown in Figure 1A, and then expressed as proportional abundance loss compared to cropland that has experienced no warming (i.e. standardised temperature anomaly of 0). In each cell, the total production of each crop⁴³ was multiplied by dependence on animal pollination,³⁴ and then adjusted for the predicted percentage reduction in insect pollinator abundance in that cell. These estimates of crop production at risk were summed across crops, and then across all terrestrial cells globally. Lines show different assumed relationships between insect pollinator abundance and crop production: dashed = convex relationships (of differing degrees, indicated by different colours: yellow = most convex; purple/blue = least convex); and solid = linear relationship.

138 Projected risk to crop production in 2050 from insect pollinator abundance losses, as a proportion of all
 139 production in a given location, is highest in the tropical regions of sub-Saharan Africa, South America, and
 140 southeast Asia (Figure 3A; see Figure S8 for underlying maps). In terms of total production potentially at
 141 risk, China, India, Indonesia, Brazil, and the Philippines emerge as being most at risk (Figure 3B). Among
 142 crops, cocoa is estimated to be at highest risk, by a large margin, especially in Africa, followed by mango
 143 (particularly in India) and watermelon (notably in China) (Figure 3B). The risk to cocoa production is

144 particularly significant in light of the social and economic context, as most cocoa is produced on small farms
145 (two to four hectares) that provide income to between 40 and 50 million people globally.⁵³ Coffee is also
146 expected to have a combination of relatively high production risk (Figure 3B) and high value, suggesting
147 that regions in which it is grown may experience economic difficulties, unless the pollination service can
148 be replaced cost effectively. Similarly to cocoa, coffee production provides income to millions of small-scale
149 farmers and their families in the tropics.⁵⁴ Therefore, the increased production risk due to loss of pollinators
150 could lead to increased income insecurity for some of the most vulnerable people globally. Projections of
151 local crop production risk are sensitive to the assumed abundance-production relationship (Figure S9), with
152 the exception of South East Asia, which is consistently projected to have high risk, and the temperate realm,
153 which is consistently projected to have low risk (Figure S10).

154 It is impossible to predict exactly how our estimates of production risk measure will translate into actual
155 crop production losses. There are multiple uncertainties associated with predicting pollinator biodiversity
156 changes and how this impacts crop production, some of which we explore here (e.g. the relationship between
157 pollinator abundance and crop production), but many of which we do not or cannot (e.g. the changing
158 distribution of crops, the economic viability of hand pollination, the buffering effects of managed pollinators,
159 the effects of climate change alone, the synergistic effects of abundance with other measures of pollinator
160 biodiversity, and other technological solutions such as the breeding or engineering of pollinator-independent
161 cultivars). Regardless, at the global scale, the changes to pollinator biodiversity that we estimate are likely
162 to be reflected in changes in the risk to production of crops dependent on animal pollination. For cocoa,
163 recent research has focused on the direct effects of climate change on crop production,^{55,56} often overlooking
164 those that might be pollinator-mediated, probably because the direct effects of climate change are easier to
165 capture. Importantly, solutions to direct and pollinator-mediated effects of climate change may differ. For
166 example, shade trees might protect crops from the detrimental effects of extreme temperatures,⁵⁵ but might
167 not for the ceratopogonid fly pollinators on which cocoa pollination depends. The focus in previous research
168 on direct climate impacts rather than pollinator-mediated effects is a key gap given that some cocoa varieties
169 are limited more by pollination availability than resource availability.⁵⁷




Figure 3: Projected change globally in crop production estimated to be at risk in 2050 under the RCP 6.0 climate scenario, assuming a linear relationship between insect pollinator abundance loss and production loss for crops dependent on animal pollination. All projections are based on mean projections of the standardized temperature anomaly based on temperature estimates from an ensemble of individual climate models. (A) The sum of crop production at risk across all crops with some dependence on animal pollination, as a proportion of the production of all crops grown in a location ('proportional production risk'). (B) The total crop production at risk for the 7 crops with the highest total pollination dependent production value globally (see Figure S11 for the top 20 crops by pollination dependent production alone, Figure S12 for country level proportional production risk, and Figure S13 for crop level proportional production risk), in million US\$ per annum, broken down into four main geographic regions. Each coloured bar represents a pollination dependent crop group: Grey, cocoa; black, tropical fruits (not recorded elsewhere); red, fruits (not recorded elsewhere); orange, mango; dark blue, coffee; yellow, watermelon; green, coconut; and other crops, light blue. Per tonne values of each crop are for the years 2015-2019 (US\$ in 2015-2019 values) taken from,⁵⁸ and total pollination dependent production according to⁴³ and.³⁴

Countries besides those with high production risk will feel the impact of losses of pollinator biodiversity and the crops that depend on them, through disruption to imports, especially as the most vulnerable crops tend to be valuable export products such as coffee and cocoa. In absolute terms, large countries such as China and the United States have the highest total import risks (Figure 4B). The Netherlands emerges as having surprisingly high risk given its size, the third largest overall import risk (Figure 4B), consistent with its status as the greatest importer of cocoa beans worldwide.⁵⁹ Import risk per capita (Figure 4A) highlights the challenges that could be faced by nations with limited agricultural production capacity, such as many island countries (e.g. Cayman Islands, Aruba, Singapore, the second, third and fourth highest import risk per capita, respectively) or countries with unfavourable environmental conditions for agriculture (e.g., Mongolia, with the 19th highest import risk per capita). Total import risk per capita tends to be high also in northern and high-income countries, particularly continental western Europe, which has large processing industries for crops such as coffee and cocoa. High income and unfavourable environment for agriculture could both account for high import risk per capita for some countries in the Middle East (United Arab Emirates, Kuwait, Saudi Arabia, which have the 5th, 13th, and 27th highest estimated pollination import risk per capita). Our predictions of import risk are based on trade patterns in the present day,³⁵ meaning we do not account for changes in trade flows that will likely occur in the future. We also assume that all production produced in a given country is exported, which is not true. Future work will explore removing the trade in re-exports from our projections.




Figure 4: Projected total import risk for 2050 under the RCP 6.0 scenario, assuming a linear relationship between insect pollinator abundance loss and production loss. Projections are based on changes in the mean standardized temperature anomaly based on projections from an ensemble of individual climate models. Import risk is a measure of how the effects of localised production risk might be distributed among other countries, calculated using trade flow data from.³⁵ For each yearly time step into the future, local production risk in each spatial cell is attributed to importers according to the quantity of pollination trade imported from each country. For example, if an importer is dependent on 3 countries for imports, at a proportion of 30%, 50%, and 20%, then any change in import risk should scale as a function of the local production risk aggregated at those same proportions. (A) The geographic distribution of total import risk adjusted for country population size and then converted to a percentile. Colours represent each percentile grouping: yellow, 0-25th percentile; orange, 25-50th; purple, 50-75th; dark blue, 75-100th. Inset plot represents the absolute import risk values on a log10 scale, with the same percentile breakpoints. (B) The total import crop production at risk in thousand tonnes per annum, for the 10 countries with the highest import production at risk.

188 Interacting effects of climate change and anthropogenic land use are rapidly restructuring biodiversity.^{26,60}
 189 Here we show that the combination of agricultural land use and recent climate change is associated with
 190 particularly large reductions in the abundance of insect pollinators. As a result, we predict that the tropics
 191 will likely experience the greatest risk of future crop pollination shortfalls, putting at risk the production of
 192 crops that depend on insect pollination. Future crop pollination risk is estimated to be highest in areas used
 193 to produce cocoa, mango, watermelon, and coffee. Given the many factors that determine crop production
 194 and crop price,⁶¹ the likely effects of insect pollinator losses on crop production are unclear, and even if
 195 they do occur, conclusive attribution is likely to be challenging. Such complications likely in part explain
 196 why identifying a strong effect of pollinator losses on global crop yield and price has thus far been so
 197 difficult.^{33,62,63} Regardless, there is sufficient evidence to suggest that declining insect pollinator abundance
 198 will influence crop production risk, particularly for those in the global south.⁶⁴ Such risk could manifest in
 199 the form of direct and immediate losses to crop production through pollinator shortfall, fluctuations in the
 200 stability of production, or through decreased resilience to changes that will happen in conjunction (e.g. the
 201 effects of extreme temperature and drought on crop growth). The health, well-being, and livelihoods of a
 202 high proportion of the global population, from small farmers to consumers, relies to some extent upon the
 203 availability and affordability of crops dependent on animal pollination, which is likely to be put at greater
 204 risk as a result of future pollinator losses as land-use and climate changes intensify.

205 References

- 206 1. Klink, R. van *et al.* Meta-analysis reveals declines in terrestrial but increases in freshwater insect
 207 abundances. *Science* **368**, 417–420 (2020).
- 208 2. Dornelas, M. *et al.* A balance of winners and losers in the Anthropocene. *Ecology Letters* **22**, 847–854
 209 (2019).
- 210 3. Lister, B. C. & Garcia, A. Climate-driven declines in arthropod abundance restructure a rainforest
 211 food web. *Proceedings of the National Academy of Sciences* **115**, E10397–E10406 (2018).
- 212 4. Biesmeijer, J. C. *et al.* Parallel declines in pollinators and insect-pollinated plants in Britain and the
 213 Netherlands. *Science* **313**, 351–354 (2006).
- 214 5. Potts, S. G. *et al.* Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology &*
 215 *Evolution* **25**, 345–353 (2010).
- 216 6. Oliver, T. H. *et al.* Declining resilience of ecosystem functions under biodiversity loss. *Nature Com-*
 217 *munications* **6**, 10122 (2015).
- 218 7. Powney, G. D. *et al.* Widespread losses of pollinating insects in Britain. *Nature Communications* **10**,
 219 1018 (2019).
- 220 8. Soroye, P., Newbold, T. & Kerr, J. Climate change contributes to widespread declines among bumble
 221 bees across continents. *Science* **367**, 685–688 (2020).
- 222 9. Basset, Y. & Lamarre, G. P. A. Toward a world that values insects. *Science* **364**, 1230–1231 (2019).
 223
- 224 10. Wagner, D. L. Insect declines in the Anthropocene. *Annual Review of Entomology* **65**, 457–480 (2020).
 225
- 226 11. Khan, S. A. *et al.* Declining abundance of pollinating insects drives falls in loquat (*Eriobotrya*
 227 *japonica*) fruit yields in the Pothwar region of Pakistan. *Agriculture, Ecosystems & Environment*
339, 108138 (2022).
- 228 12. Lewinsohn, T. M., Agostini, K., Lucci Freitas, A. V. & Melo, A. S. Insect decline in Brazil: an
 229 appraisal of current evidence. *Biology Letters* **18**, 20220219 (2022).

- 230 13. Janzen, D. H. & Hallwachs, W. Perspective: Where might be many tropical insects? *Biological*
231 *Conservation* **233**, 102–108 (2019).
- 232 14. Warren, M. S. *et al.* Rapid responses of British butterflies to opposing forces of climate and habitat
233 change. *Nature* **414**, 65–69 (2001).
- 234 15. De Palma, A. *et al.* Predicting bee community responses to land-use changes: Effects of geographic
235 and taxonomic biases. *Scientific Reports* **6**, 31153 (2016).
- 236 16. Oliver, I., Dorrough, J., Doherty, H. & Andrew, N. R. Additive and synergistic effects of land cover,
237 land use and climate on insect biodiversity. *Landscape Ecology* **31**, 2415–2431 (2016).
- 238 17. Millard, J. *et al.* Global effects of land-use intensity on local pollinator biodiversity. *Nature Commu-*
239 *nications* **12**, 2902 (2021).
- 240 18. Hickling, R., Roy, D. B., Hill, J. K., Fox, R. & Thomas, C. D. The distributions of a wide range of
241 taxonomic groups are expanding polewards. *Global Change Biology* **12**, 450–455 (2006).
- 242 19. Martay, B. *et al.* Impacts of climate change on national biodiversity population trends. *Ecography*
243 **40**, 1139–1151 (2017).
- 244 20. Lenoir, J. *et al.* Species better track climate warming in the oceans than on land. *Nature Ecology &*
245 *Evolution* **4**, 1044–1059 (2020).
- 246 21. Halsch, C. A. *et al.* Insects and recent climate change. *Proceedings of the National Academy of*
247 *Sciences* **118**, e2002543117 (2021).
- 248 22. Suggitt, A. J. *et al.* Extinction risk from climate change is reduced by microclimatic buffering. *Nature*
249 *Climate Change* **8**, 713–717 (2018).
- 250 23. Platts, P. J. *et al.* Habitat availability explains variation in climate-driven range shifts across multiple
251 taxonomic groups. *Scientific Reports* **9**, 15039 (2019).
- 252 24. Oliver, T. H. *et al.* Large extents of intensive land use limit community reorganization during climate
253 warming. *Global Change Biology* **23**, 2272–2283 (2017).
- 254 25. Peters, M. K. *et al.* Climate–land-use interactions shape tropical mountain biodiversity and ecosystem
255 functions. *Nature* **568**, 88–92 (2019).
- 256 26. Outhwaite, C. L., McCann, P. & Newbold, T. Agriculture and climate change are reshaping insect
257 biodiversity worldwide. *Nature* **605**, 97–102 (2022).
- 258 27. Deutsch, C. A. *et al.* Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings*
259 *of the National Academy of Sciences* **105**, 6668–6672 (2008).
- 260 28. Rodger, J. G. *et al.* Widespread vulnerability of flowering plant seed production to pollinator declines.
261 *Science Advances* **7**, eabd3524 (2021).
- 262 29. Klein, A., Steffan–Dewenter, I. & Tschardtke, T. Fruit set of highland coffee increases with the
diversity of pollinating bees. *Proceedings of the Royal Society of London. Series B: Biological Sciences*
263 **270**, 955–961 (2003).
- 264 30. Ricketts, T. H., Daily, G. C., Ehrlich, P. R. & Michener, C. D. Economic value of tropical forest to
265 coffee production. *Proceedings of the National Academy of Sciences* **101**, 12579–12582 (2004).
- 266 31. Holzschuh, A., Dormann, C. F., Tschardtke, T. & Steffan–Dewenter, I. Expansion of mass-flowering
crops leads to transient pollinator dilution and reduced wild plant pollination. *Proceedings of the*
267 *Royal Society B: Biological Sciences* **278**, 3444–3451 (2011).

- 268 32. Woodcock, B. A. *et al.* Meta-analysis reveals that pollinator functional diversity and abundance
269 enhance crop pollination and yield. *Nature Communications* **10**, 1481 (2019).
- 270 33. Aizen, M. A., Garibaldi, L. A., Cunningham, S. A. & Klein, A. M. Long-term global trends in crop
271 yield and production reveal no current pollination shortage but increasing pollinator dependency.
Current Biology **18**, 1572–1575 (2008).
- 272 34. Klein, A.-M. *et al.* Importance of pollinators in changing landscapes for world crops. *Proceedings of*
273 *the Royal Society B: Biological Sciences* **274**, 303–313 (2007).
- 274 35. Silva, F. D. S. *et al.* Virtual pollination trade uncovers global dependence on biodiversity of developing
275 countries. *Science Advances* **7**, eabe6636 (2021).
- 276 36. Albrecht, M., Schmid, B., Hautier, Y. & Müller, C. B. Diverse pollinator communities enhance plant
277 reproductive success. *Proceedings of the Royal Society B: Biological Sciences* **279**, 4845–4852 (2012).
- 278 37. Brittain, C., Kremen, C. & Klein, A.-M. Biodiversity buffers pollination from changes in environmen-
279 tal conditions. *Global Change Biology* **19**, 540–547 (2013).
- 280 38. Genung, M. A. *et al.* The relative importance of pollinator abundance and species richness for the
281 temporal variance of pollination services. *Ecology* **98**, 1807–1816 (2017).
- 282 39. Winfree, R. *et al.* Species turnover promotes the importance of bee diversity for crop pollination at
283 regional scales. *Science* **359**, 791–793 (2018).
- 284 40. Dainese, M. *et al.* A global synthesis reveals biodiversity-mediated benefits for crop production.
285 *Science Advances* **5**, eaax0121 (2019).
- 286 41. Lonsdorf, E. *et al.* Modelling pollination services across agricultural landscapes. *Annals of Botany*
287 **103**, 1589–1600 (2009).
- 288 42. Hudson, L. N. *et al.* The database of the PREDICTS (Projecting Responses of Ecological Diversity
289 In Changing Terrestrial Systems) project. *Ecology and Evolution* **7**, 145–188 (2017).
- 290 43. Monfreda, C., Ramankutty, N. & Foley, J. A. Farming the planet: 2. Geographic distribution of crop
291 areas, yields, physiological types, and net primary production in the year 2000. *Global Biogeochemical*
Cycles **22**, GB1022 (2008).
- 292 44. Rogelj, J., Meinshausen, M. & Knutti, R. Global warming under old and new scenarios using IPCC
293 climate sensitivity range estimates. *Nature Climate Change* **2**, 248–253 (2012).
- 294 45. Oliver, T. H. & Morecroft, M. D. Interactions between climate change and land use change on bio-
295 diversity: Attribution problems, risks, and opportunities. *Wiley Interdisciplinary Reviews: Climate*
Change **5**, 317–335 (2014).
- 296 46. Rech, A. R. *et al.* The macroecology of animal versus wind pollination: Ecological factors are more
297 important than historical climate stability. *Plant Ecology & Diversity* **9**, 253–262 (2016).
- 298 47. Peters, M. K., Peisker, J., Steffan-Dewenter, I. & Hoiss, B. Morphological traits are linked to the
299 cold performance and distribution of bees along elevational gradients. *Journal of Biogeography* **43**,
2040–2049 (2016).
- 300 48. Stavert, J. R. *et al.* Hairiness: The missing link between pollinators and pollination. *PeerJ* **4**, e2779
301 (2016).
- 302 49. Aizen, M. A., Garibaldi, L. A. & Harder, L. D. Myth and reality of a global crisis for agricultural
303 pollination. *Asociacion Argentina de Ecologia* (2022).

- 304 50. Ürge-Vorsatz, D., Herrero, S. T., Dubash, N. K. & Lecocq, F. Measuring the co-benefits of climate
305 change mitigation. *Annual Review of Environment and Resources* **39**, 549–582 (2014).
- 306 51. Timmermann, A. *et al.* El Niño–southern oscillation complexity. *Nature* **559**, 535–545 (2018).
307
- 308 52. Dee, L. E. *et al.* Clarifying the effect of biodiversity on productivity in natural ecosystems with
309 longitudinal data and methods for causal inference. *Nature Communications* **14**, 2607 (2023).
- 310 53. Beg, M. S., Ahmad, S., Jan, K. & Bashir, K. Status, supply chain and processing of cocoa—a review.
311 *Trends in food science & technology* **66**, 108–116 (2017).
- 312 54. Pham, Y., Reardon-Smith, K., Mushtaq, S. & Cockfield, G. The impact of climate change and
313 variability on coffee production: A systematic review. *Climatic Change* **156**, 609–630 (2019).
- 314 55. Schroth, G., Läderach, P., Martinez-Valle, A. I., Bunn, C. & Jassogne, L. Vulnerability to climate
change of cocoa in West Africa: Patterns, opportunities and limits to adaptation. *Science of the Total
315 Environment* **556**, 231–241 (2016).
- 316 56. Ofori-Boateng, K. & Insah, B. The impact of climate change on cocoa production in West Africa.
317 *International Journal of Climate Change Strategies and Management* (2014).
- 318 57. Groeneveld, J. H., Tschardtke, T., Moser, G. & Clough, Y. Experimental evidence for stronger cacao
yield limitation by pollination than by plant resources. *Perspectives in Plant Ecology, Evolution and
319 Systematics* **12**, 183–191 (2010).
- 320 58. Food and Agriculture Organization of the United Nations (FAO). Producer prices & crops and live-
321 stock products. (2022).
- 322 59. Simoes, A. J. G. & Hidalgo, C. A. The economic complexity observatory: An analytical tool for under-
standing the dynamics of economic development. in *Workshops at the twenty-fifth AAAI conference
323 on artificial intelligence* (2011).
- 324 60. Ganuza, C. *et al.* Interactive effects of climate and land use on pollinator diversity differ among taxa
325 and scales. *Science advances* **8**, eabm9359 (2022).
- 326 61. Gaetano, S. F., Emilia, L., Francesco, C., Gianluca, N. & Antonio, S. Drivers of grain price volatility:
327 A cursory critical review. *Agricultural Economics* **64**, 347–356 (2018).
- 328 62. Aizen, M. A., Garibaldi, L. A., Cunningham, S. A. & Klein, A. M. How much does agriculture depend
on pollinators? Lessons from long-term trends in crop production. *Annals of botany* **103**, 1579–1588
329 (2009).
- 330 63. Garibaldi, L. A., Aizen, M. A., Klein, A. M., Cunningham, S. A. & Harder, L. D. Global growth
and stability of agricultural yield decrease with pollinator dependence. *Proceedings of the National
331 Academy of Sciences* **108**, 5909–5914 (2011).
- 332 64. Dicks, L. V. *et al.* A global-scale expert assessment of drivers and risks associated with pollinator
333 decline. *Nature Ecology & Evolution* **5**, 1453–1461 (2021).

334 Acknowledgements

335 Thanks to Richard Gregory, Robin Freeman, and colleagues on the GLITRS project for useful discussions
336 and comments. Thanks to Andy Purvis and Melinda Mills, who both provided support and guidance.
337 Thanks also to Opeyemi Adedaja, Sabrina Gavini, Esther Kioko, Michael Kuhlmann, Zong-Xin Ren, and
338 Manu Saunders, who provided input on our set of likely pollinating species, and to all contributors of the
339 PREDICTS database.

340 Funding

341 JM received PhD funding from the London NERC DTP and the RSPB. JM, TN and CLO receive
342 funding from the UK Natural Environment Research Council (NERC; grant references: NE/R010811/1,
343 NE/V006533/1 and NE/V006800/1). TN and SC receive funding from the UK Economic and Social
344 Research Council (reference: ES/S008160/1). TN is additionally supported by a University Research
345 Fellowship from the Royal Society. LGC was funded by the brazilian Conselho Nacional de Desenvolvimento
346 Científico e Tecnológico [CNPq 307625/2021-4].

347 Contributions

348 JM and TN conceived the study. JM, CO, FDSS, JO, LD, LGC, SC, and TN designed the methodology of
349 crop pollination risk and import production risk. All authors edited and approved the final version of the
350 manuscript.

351 Competing interests

352 All authors declare that they have no competing interests.

353 Data availability

354 All code is currently hosted openly on GitHub here ([https://github.com/Joemillard/Worldwide-](https://github.com/Joemillard/Worldwide-vulnerability-of-local-pollinator-abundance-and-crop-pollination-to-land-use-and-climate)
355 [vulnerability-of-local-pollinator-abundance-and-crop-pollination-to-land-use-and-climate](https://github.com/Joemillard/Worldwide-vulnerability-of-local-pollinator-abundance-and-crop-pollination-to-land-use-and-climate)). On publi-
356 cation all code will be versioned on Zenodo.

357 Code availability

358 Links to all data are currently hosted openly on GitHub here ([https://github.com/Joemillard/Worldwide-](https://github.com/Joemillard/Worldwide-vulnerability-of-local-pollinator-abundance-and-crop-pollination-to-land-use-and-climate)
359 [vulnerability-of-local-pollinator-abundance-and-crop-pollination-to-land-use-and-climate](https://github.com/Joemillard/Worldwide-vulnerability-of-local-pollinator-abundance-and-crop-pollination-to-land-use-and-climate)). On publication
360 all primary data and links will be versioned on Zenodo.

361 Materials & Correspondence

362 Correspondence should be addressed to Joseph Millard (joseph.millard@nhm.ac.uk)