

## Heavy Metals in Estuarine Shellfish from Oregon

Richard S. Caldwell<sup>1,\*</sup> and Donald R. Buhler<sup>†</sup>

\*Department of Fisheries and Wildlife, Oregon State University, Marine Science Center, Newport, Oregon 97365 and <sup>†</sup>Department of Agricultural Chemistry and Environmental Health Sciences Center, Oregon State University, Corvallis, Oregon 97331

**Abstract.** The concentrations of Hg, Zn, Mn, Fe, Cu, and Cd were determined in marine shellfish from the Oregon coast. The levels of these metals in both crustaceans and molluscs were generally typical of those reported from other geographic areas. The primary exception was Cd, which in crustaceans was 5 to 40 times lower and in three of the six marine bivalve molluscs studied was 5 to 25 times lower than that found in most previous studies. The low concentrations of Cd may be due to the lower level and types of industrial activity in coastal Oregon estuaries compared to other regions of the world, and/or to a lower rate of input of this metal from natural weathering processes in this region. No significant variations due either to season or geographical location were apparent in the three principal shellfish species studied.

Concern for the contamination of natural environments with heavy metals has become widespread, derived both from the potential for deleterious effects on the biota (Bryan 1971; Eisler 1971; Thurberg *et al.* 1973) and from possible human health hazards where such species are consumed by man (Takeuchi 1972). Elevated metal concentrations in marine organisms are restricted to estuaries and the narrow coastal margin and are associated with surface runoff of coastal streams, drainage from industrial areas, the disposal of solid or liquid wastes, and various ship-related inputs (Eganhouse and Young 1976; Preston 1973; Taylor 1974; Young *et al.* 1973). Human populations, concentrated in coastal regions, can mobilize metals to the marine environment in amounts similar to or even exceed-

ing those introduced through continental weathering processes.

Certain heavy metals concentrate in marine food webs (Buhler *et al.* 1975), and filter feeding organisms such as bivalve molluscs often serve as important environmental sinks (Pringle *et al.* 1968). The principal objective of this study was to obtain information on the concentrations of six metals in selected shellfish from relatively unpolluted coastal areas in Oregon. The information resulting from examining several species including some commonly consumed by humans, could provide a baseline both in terms of mean metals concentrations and seasonal variability. In addition, the possibility that metal concentrations in shellfish from certain Oregon estuaries may be indicative of the relative degree of natural or anthropogenic contamination by heavy metals has been examined.

### Materials and Methods

Marine molluscs and crustaceans were collected from various Oregon estuaries and open coast locations (Figure 1). Characteristics of the sampling locations, such as size of drainage basin, level of freshwater input, human population levels, degree of industrialization and nature of industrial activities within the riverine systems are summarized in Table 1.

Recreationally important shellfish, including cockles (*Clinocardium nuttallii*), softshell clams (*Mya arenaria*) and the commercially important Dungeness crab (*Cancer magister*), were collected quarterly in 1972 from Oregon estuaries. Additional samples were obtained on an irregular basis in 1970, 1971, and 1974. Other molluscan and crustacean species collected irregularly during 1971 and 1972 were: the freshwater clams, *Anodonta* sp. and *Corbicula fluminea*; the ghost shrimp, *Callinassa californiensis*; the sand shrimp, *Crangon franciscorum*; the spotted shore crab, *Hemigrapsus nudus*; the mussels, *Mytilus californianus* and *Mytilus edulis*; the carnivorous gastropod, *Nucella lamellosa*; the herbivorous gastropod *Tegula funebris*; and the marine clams, *Protothaca staminea* and *Tre-*

<sup>1</sup> Address correspondence to R. S. Caldwell, Northwestern Aquatic Sciences, Inc., P.O. Box 1437, Newport, OR 97365




Fig. 1. Location of sampling stations along the Oregon coast

*sus capax*. Animals were rinsed, placed in plastic bags, and frozen for subsequent analysis.

Analyses were performed as described previously (Buhler *et al.* 1975; Phillips and Buhler 1980) on the rinsed whole soft tissues of clams and other molluscs and on hepatopancreas and muscle tissues dissected from crustaceans. Tissues were digested by boiling under reflux for three hr with concentrated nitric acid and 30% hydrogen peroxide. Mercury (Hg) was analyzed by flameless atomic absorption spectrophotometry using a Coleman Model 50 mercury analyzer. Zinc (Zn), manganese (Mn), iron (Fe), and copper (Cu) were measured in a Perkin-Elmer Model 403 atomic absorption spectrophotometer. Cadmium (Cd) was determined with the same instrument equipped with a Model HGA-2000 graphite furnace after ammonium pyrrolidine dithiocarbamate chelation and extraction of the digestion mixture with methyl isobutylketone (Koirtyohann and Wen 1973). Samples were analyzed by the method of additions and the accuracy of the analytical methods was established by the periodic analysis of a National Bureau of Standards bovine liver reference standard.

## Results and Discussion

The mean concentrations of six heavy metals in the muscle and hepatopancreas tissues of *C. magister*,

and in whole soft tissues of *C. nuttallii* and *M. arenaria*, from all sample sites along the Oregon coast are given in Table 2. Seasonal and geographic comparisons of metal concentrations are shown for *C. magister* in Table 3, for *M. arenaria* in Table 4 and for *C. nuttallii* in Table 5. The levels of Hg, Cd, and Zn in other shellfish species sampled on an irregular basis are presented in Table 6.

## Seasonal and Geographic Variations

No persistent differences in metal concentrations with sampling location were noted for the three major species and ten estuaries studied (Tables 3–5). The results are consistent with the relatively low degree of industrialization of the bays examined (Table 1) and the absence of any major differences in mining or weathering inputs into the streams that supply these estuaries. In addition, there were no consistent seasonal trends in the present study, with the possible exception of Hg (Tables 3–5). A tendency for higher concentrations of this metal to occur during the spring and summer months was observed in the three principle species of this study. These increases varied with sampling location but not in a manner similar for each species. Concentrations of nine trace metals in the Pacific oyster *Crassostrea gigas* from the Washington coast similarly failed to show a significant seasonal variation (Pringle *et al.* 1968).

Phillips (1976) observed seasonal variation in the concentrations of Zn, Cd, and Cu in *Mytilus edulis* from Australia, ascribed in part to variations in the wet weight. The source of seasonal fluctuation in wet weight was uncertain but may have been due to differences in nutritional or sexual status of the organisms at different times of the year. Bryan (1973) observed seasonal variations of metals in scallops and suggested that alterations in the metal content of phytoplankton may have been responsible.

## Crustaceans

The concentrations of Hg, Zn, and Fe found in *C. magister* (Table 2) and Hg and Zn in *C. californiensis*, *C. franciscorum* and *H. nudus* (Table 6) from Oregon were comparable to levels previously reported in other crustaceans and in other geographic locations (Fowler *et al.* 1975; Freeman *et al.* 1974; Reimer and Reimer 1975; Zook *et al.* 1976). Concentrations of Cu in *C. magister* muscle were also similar to those reported for four other decapods (Bryan 1964; Fowler *et al.* 1975; Topping 1973). In *C. magister* hepatopancreas, the concentrations of

**Table 1.** Characteristics of sampling locations shown in Figure 1

Sampling station	Location	Physical characteristics			Human population + pollutional characteristics			Comments
		Total estuarine area <sup>a</sup> (A)	Size of drainage basin (mi <sup>2</sup> )	Mean annual freshwater input <sup>b</sup> (ac-ft × 10 <sup>3</sup> )	Estimated human population <sup>c</sup>	Municipal sewage discharge <sup>b,d</sup>	Relative degree of industrialization <sup>b,e</sup>	
1	Columbia River	93,782	259,000 <sup>f</sup>	192,575	>1,000,000	+	+++++	Pulp mills, shipping, various industries
2	Nehalem Bay	2,309	855 <sup>b</sup>	2,700	1,188 <sup>b</sup>	—	+	Seafood processing
3	Tillamook Bay	8,289	540 <sup>b</sup>	>2,203		+	++	
4	Netarts Bay	2,325	14	42	930	—		
5	Nestucca Bay	1,000	322	1,600	695	+	+	Pulp mill, seafood processing
6	Salmon River	204	75	450	65	—		
7	Siletz Bay	1,187	373	1,800	1,203	+	+	
8	Yaquina Bay	3,910	253	780	8,381	+	+++	
9	Alsea Bay	2,146	474	1,500	900	+	+	Pulp mill, seafood processing
10	Umpqua River	6,830	4,560	6,700	>5,269	+	++	
11	Coos Bay	12,380	605	2,200	32,092	+	++++	
12	Coquille Bay	771	1,058	2,400	9,180	+	+	2 mi north of Pistol River Sampled near municipal discharge
13	Open coast	—	—	—	—	—		
14	Open coast—Brookings	—	—	—	—	+	+	

<sup>a</sup> Bourke *et al.* (1971)<sup>b</sup> Percy *et al.* (1974)<sup>c</sup> Human population of communities on estuary or riverine system<sup>d</sup> Municipal sewage discharge present (+) or absent (—)<sup>e</sup> Relative degree of industrialization indicated by the number of pluses<sup>f</sup> Anonymous (1973)**Table 2.** Concentrations of metals in three species of shellfish from all sites in Oregon sampled during 1970–1974 (expressed as µg/g wet weight)<sup>a</sup>

Metal	Dungeness crab ( <i>C. magister</i> )		Cockles ( <i>C. nuttallii</i> ) <sup>b</sup>	Softshell clam ( <i>M. arenaria</i> ) <sup>b</sup>	
	Muscle	Hepatopancreas			
Hg	0.108 ± 0.084 (180)	0.098 ± 0.062 (131)	0.019 ± 0.009 (71)	0.025 ± 0.018 (139)	
Cd	0.022 ± 0.020 (124)	0.149 ± 0.099 (98)	0.022 ± 0.031 (59)	0.028 ± 0.021 (111)	
Zn	41.9 ± 3.35 (166)	21.2 ± 8.10 (123)	10.0 ± 2.42 (62)	10.7 ± 3.8 (110)	
Mn	1.28 ± 0.754 (73)	6.67 ± 6.79 (59)	6.02 ± 4.45 (32)	24.4 ± 22.8 (31)	
Fe	5.71 ± 4.40 (73)	40.6 ± 34.0 (59)	460 ± 360 (47)	934 ± 711 (67)	
Cu	10.1 ± 4.41 (73)	27.5 ± 22.6 (59)	1.24 ± 0.445 (47)	2.54 ± 1.0 (67)	

<sup>a</sup> Mean ± S.D. Number of animals expressed in parentheses<sup>b</sup> Total soft tissues

**Table 3.** Concentration of metals in tissues of Dungeness crab (*Cancer magister*) from Oregon estuaries (expressed as  $\mu\text{g/g}$  wet weight)<sup>a</sup>

Collection site and date (Mo/Yr)	Concentration, $\mu\text{g/g}$ wet weight		
	Hg	Cd	Zn
<i>Muscle</i>			
Tillamook Bay			
2/71	0.085 $\pm$ 0.049 (13)	0.011 $\pm$ 0.007 (6)	37.5 $\pm$ 7.2 (10)
1/72	0.136 $\pm$ 0.152 (8)	0.019 $\pm$ 0.006 (6)	42.2 $\pm$ 7.8 (6)
4/72	0.092 $\pm$ 0.081 (3)	0.007 $\pm$ 0.003 (3)	40.9 $\pm$ 4.4 (3)
7/72	0.136 $\pm$ 0.056 (8)	0.022 $\pm$ 0.009 (6)	36.2 $\pm$ 8.8 (8)
10/72	0.059 $\pm$ 0.028 (8)	0.029 $\pm$ 0.032 (6)	43.1 $\pm$ 8.8 (8)
Netarts Bay			
2/71	0.124 $\pm$ 0.117 (10)	0.016 $\pm$ 0.027 (5)	36.8 $\pm$ 9.1 (8)
1/72	0.101 $\pm$ 0.030 (6)	0.014 $\pm$ 0.004 (6)	45.6 $\pm$ 2.6 (6)
4/72	0.097 $\pm$ 0.039 (7)	0.016 $\pm$ 0.014 (6)	41.0 $\pm$ 7.5 (6)
7/72	0.103 $\pm$ 0.032 (8)	0.012 $\pm$ 0.008 (6)	43.5 $\pm$ 7.6 (8)
Yaquina Bay			
12/71	0.093 $\pm$ 0.050 (6)	0.059 $\pm$ 0.024 (6)	40.5 $\pm$ 6.4 (6)
Umpqua River Estuary			
7/72	0.078 $\pm$ 0.029 (8)	0.031 $\pm$ 0.031 (6)	42.4 $\pm$ 10.6 (8)
10/72	0.088 $\pm$ 0.038 (8)	0.027 $\pm$ 0.015 (6)	50.4 $\pm$ 4.9 (8)
Coos Bay			
12/71	0.097 $\pm$ 0.070 (6)	0.019 $\pm$ 0.016 (6)	30.7 $\pm$ 3.1 (6)
4/72	0.153 $\pm$ 0.052 (16)	0.015 $\pm$ 0.009 (6)	50.6 $\pm$ 10.0 (16)
7/72	0.113 $\pm$ 0.061 (8)	0.018 $\pm$ 0.008 (6)	46.0 $\pm$ 7.5 (8)
9/72	0.051 $\pm$ 0.033 (8)	0.012 $\pm$ 0.003 (6)	44.8 $\pm$ 6.5 (8)
Coquille Bay			
3/71	0.085 $\pm$ 0.032 (13)	0.051 $\pm$ 0.039 (6)	31.6 $\pm$ 4.1 (6)
4/72	0.141 $\pm$ 0.110 (9)	0.033 $\pm$ 0.014 (6)	48.8 $\pm$ 9.6 (9)
7/72	0.126 $\pm$ 0.103 (8)	0.019 $\pm$ 0.011 (6)	45.4 $\pm$ 8.4 (8)
10/72	0.031 $\pm$ 0.019 (8)	0.021 $\pm$ 0.014 (6)	40.0 $\pm$ 7.2 (8)
<i>Hepatopancreas</i>			
Tillamook Bay			
1/72	0.083 $\pm$ 0.069 (8)	0.166 $\pm$ 0.044 (6)	13.1 $\pm$ 3.3 (6)
4/72	0.081 $\pm$ 0.055 (3)	0.199 $\pm$ 0.038 (3)	17.2 $\pm$ 4.4 (3)
7/72	0.165 $\pm$ 0.128 (8)	0.228 $\pm$ 0.214 (6)	18.5 $\pm$ 5.7 (8)
10/72	0.060 $\pm$ 0.029 (8)	0.150 $\pm$ 0.070 (6)	17.1 $\pm$ 8.0 (8)
Netarts Bay			
1/72	0.066 $\pm$ 0.025 (6)	0.064 $\pm$ 0.030 (6)	20.4 $\pm$ 5.2 (6)
4/72	0.073 $\pm$ 0.018 (7)	0.100 $\pm$ 0.046 (5)	16.9 $\pm$ 3.2 (5)
7/72	0.112 $\pm$ 0.045 (8)	0.163 $\pm$ 0.120 (6)	21.0 $\pm$ 10.6 (8)
Yaquina Bay			
12/71	0.058 $\pm$ 0.019 (6)	0.141 $\pm$ 0.051 (6)	20.6 $\pm$ 7.4 (6)
Umpqua River Estuary			
7/72	0.066 $\pm$ 0.032 (8)	0.169 $\pm$ 0.050 (6)	27.7 $\pm$ 6.7 (8)
10/72	0.103 $\pm$ 0.049 (8)	0.152 $\pm$ 0.075 (6)	22.0 $\pm$ 4.2 (8)
Coos Bay			
12/71	0.085 $\pm$ 0.026 (6)	0.091 $\pm$ 0.077 (6)	16.5 $\pm$ 6.2 (6)
4/72	0.148 $\pm$ 0.050 (16)	0.107 $\pm$ 0.044 (6)	19.2 $\pm$ 7.1 (12)
7/72	0.105 $\pm$ 0.052 (8)	0.109 $\pm$ 0.044 (6)	19.1 $\pm$ 4.5 (8)
9/72	0.062 $\pm$ 0.048 (8)	0.083 $\pm$ 0.028 (6)	21.7 $\pm$ 7.2 (8)
Coquille Bay			
4/72	0.158 $\pm$ 0.068 (9)	0.275 $\pm$ 0.170 (6)	35.7 $\pm$ 10.2 (9)
7/72	0.094 $\pm$ 0.048 (8)	0.122 $\pm$ 0.067 (6)	20.6 $\pm$ 6.1 (8)
10/72	0.058 $\pm$ 0.013 (8)	0.206 $\pm$ 0.089 (6)	23.6 $\pm$ 7.5 (6)

Cu were similar to those seen in *C. pagurus* (Topping 1973), but were 4 to 16 times lower than reported for the same tissues of *Homarus vulgaris* (Bryan 1964; Topping 1973). Variation in tissue Cu concentrations in crustaceans may be more related

to variation in hemocyanin content of the blood than to levels of environmental contamination. The Mn concentration of *C. magister* muscle, 1.28  $\mu\text{g/g}$  (Table 2), was three times that found by Fowler *et al.* 1975) in *Cancer anthonyi*.

Table 3. (cont'd)

Collection site and date (Mo/Yr)	Concentration, $\mu\text{g/g}$ wet weight		
	Mn	Fe	Cu
<i>Muscle</i>			
Tillamook Bay			
2/71	1.57 $\pm$ 1.04 (5)	11.8 $\pm$ 5.5 (5)	6.80 $\pm$ 1.08 (5)
1/72	1.78 $\pm$ 0.40 (3)	7.81 $\pm$ 2.45 (3)	8.79 $\pm$ 2.72 (3)
4/72	1.19 $\pm$ 0.14 (3)	8.20 $\pm$ 0.98 (3)	15.9 $\pm$ 11.4 (3)
7/72	1.09 $\pm$ 0.20 (6)	2.99 $\pm$ 1.07 (6)	11.0 $\pm$ 4.6 (6)
10/72	—	—	—
Netarts Bay			
2/71	—	—	—
1/72	1.49 $\pm$ 0.04 (6)	7.46 $\pm$ 3.65 (6)	14.9 $\pm$ 3.2 (6)
4/72	—	—	—
7/72	0.82 $\pm$ 0.26 (6)	1.80 $\pm$ 0.67 (6)	13.2 $\pm$ 5.5 (6)
Yaquina Bay			
12/71	1.61 $\pm$ 0.21 (6)	9.49 $\pm$ 5.49 (6)	10.2 $\pm$ 3.7 (6)
Umpqua River Estuary			
7/72	0.79 $\pm$ 0.25 (6)	1.68 $\pm$ 0.81 (6)	9.24 $\pm$ 2.48 (6)
10/72	—	—	—
Coos Bay			
12/71	1.72 $\pm$ 0.29 (6)	6.18 $\pm$ 0.28 (6)	9.74 $\pm$ 3.29 (6)
4/72	—	—	—
7/72	0.70 $\pm$ 0.34 (5)	1.56 $\pm$ 0.24 (5)	7.54 $\pm$ 2.72 (5)
9/72	—	—	—
Coquille Bay			
3/71	2.85 $\pm$ 1.34 (5)	11.7 $\pm$ 3.4 (5)	7.28 $\pm$ 2.08 (5)
4/72	1.42 $\pm$ 0.60 (5)	7.39 $\pm$ 2.37 (5)	9.11 $\pm$ 2.79 (5)
7/72	0.53 $\pm$ 0.02 (6)	1.77 $\pm$ 0.80 (6)	7.21 $\pm$ 2.37 (6)
10/72	0.71 $\pm$ 0.31 (5)	3.73 $\pm$ 1.52 (5)	11.7 $\pm$ 3.4 (5)
<i>Hepatopancreas</i>			
Tillamook Bay			
1/72	13.8 $\pm$ 12.3 (3)	41.7 $\pm$ 7.26 (3)	65.3 $\pm$ 41.8 (3)
4/72	3.73 $\pm$ 0.98 (3)	44.0 $\pm$ 27.8 (3)	48.9 $\pm$ 39.4 (3)
7/72	5.29 $\pm$ 1.96 (6)	23.8 $\pm$ 3.6 (6)	34.5 $\pm$ 33.0 (6)
10/72	—	—	—
Netarts Bay			
1/72	1.75 $\pm$ 0.14 (3)	49.6 $\pm$ 33.0 (3)	38.9 $\pm$ 22.3 (3)
4/72	—	—	—
7/72	6.06 $\pm$ 6.06 (6)	24.6 $\pm$ 14.5 (6)	30.6 $\pm$ 15.3 (6)
Yaquina Bay			
12/71	2.16 $\pm$ 0.81 (6)	99.8 $\pm$ 51.6 (6)	16.3 $\pm$ 6.5 (6)
Umpqua River Estuary			
7/72	5.20 $\pm$ 2.71 (6)	14.3 $\pm$ 5.1 (6)	16.0 $\pm$ 7.5 (6)
10/72	—	—	—
Coos Bay			
12/71	10.7 $\pm$ 9.6 (6)	32.6 $\pm$ 15.6 (6)	39.4 $\pm$ 21.4 (6)
4/72	—	—	—
7/72	5.14 $\pm$ 2.17 (4)	27.0 $\pm$ 5.0 (4)	11.9 $\pm$ 4.7 (4)
9/72	—	—	—
Coquille Bay			
4/72	18.9 $\pm$ 8.0 (5)	43.5 $\pm$ 25.7 (5)	21.3 $\pm$ 8.5 (5)
7/72	3.02 $\pm$ 1.14 (6)	52.9 $\pm$ 50.2 (6)	17.5 $\pm$ 10.3 (6)
10/72	5.22 $\pm$ 3.29 (5)	28.4 $\pm$ 22.1 (5)	16.2 $\pm$ 10.7 (5)

<sup>a</sup> Mean  $\pm$  S.D. Number of animals analyzed indicated in parentheses

The concentration of Cd in crustacean tissue was generally much lower than reported previously (Eisler *et al.* 1972; Fowler *et al.* 1975; Topping 1973; Zook *et al.* 1976). Mean Cd levels in *C. magister*

muscle and hepatopancreas were 0.022 and 1.49  $\mu\text{g/g}$ , respectively (Table 2), and 0.032  $\mu\text{g/g}$  in whole *H. nudus* (Table 6), values approximately 5 to 40 times lower than that found in the same tissues of

**Table 4.** Concentration of metals in softshell clams (*Mya arenaria*) from Oregon estuaries (expressed as  $\mu\text{g/g}$  wet weight)<sup>a</sup>

Collection site and date (Mo/Yr)	Hg	Cd	Zn	Mn	Fe	Cu
Nehalem Bay						
2/71	0.018 $\pm$ 0.004 (6)	0.068 $\pm$ 0.017 (6)	13.4 $\pm$ 2.8 (6)	—	—	—
Tillamook Bay						
1/72	0.014 $\pm$ 0.009 (10)	0.012 $\pm$ 0.009 (6)	6.36 $\pm$ 1.60 (6)	—	661 $\pm$ 317 (6)	1.63 $\pm$ 0.26 (6)
4/72	0.045 $\pm$ 0.017 (4)	0.037 $\pm$ 0.013 (4)	10.4 $\pm$ 2.2 (4)	—	1397 $\pm$ 1151 (4)	2.46 $\pm$ 0.97 (4)
7/72	0.028 $\pm$ 0.007 (4)	0.033 $\pm$ 0.008 (4)	10.1 $\pm$ 1.9 (4)	—	1446 $\pm$ 831 (4)	1.99 $\pm$ 0.06 (4)
10/72	0.017 $\pm$ 0.010 (4)	0.029 $\pm$ 0.006 (4)	11.0 $\pm$ 2.9 (4)	5.27 $\pm$ 1.70 (4)	363 $\pm$ 194 (4)	2.43 $\pm$ 1.22 (4)
Nestucca Bay						
4/71	0.020 $\pm$ 0.011 (6)	0.046 $\pm$ 0.015 (6)	11.8 $\pm$ 1.5 (6)	—	—	—
Salmon River Estuary						
4/71	0.018 $\pm$ 0.006 (5)	0.067 $\pm$ 0.006 (4)	7.64 $\pm$ 1.07 (4)	—	—	—
Siletz Bay						
4/71	0.019 $\pm$ 0.005 (6)	0.058 $\pm$ 0.025 (6)	9.22 $\pm$ 1.82 (6)	—	—	—
Yaquina Bay						
6/70	0.040 $\pm$ 0.010 (3)	—	—	—	—	—
11/70	0.016 $\pm$ 0.003 (6)	0.027 $\pm$ 0.005 (4)	6.99 $\pm$ 1.12 (4)	—	—	—
1/72	0.016 $\pm$ 0.009 (10)	0.018 $\pm$ 0.013 (6)	10.7 $\pm$ 0.8 (6)	23.2 $\pm$ 16.5 (5)	869 $\pm$ 330 (5)	1.93 $\pm$ 0.68 (5)
4/72	0.014 $\pm$ 0.006 (11)	0.001 $\pm$ 0.000 (3)	8.7 $\pm$ 1.1 (6)	—	1835 $\pm$ 845 (6)	2.69 $\pm$ 1.19 (6)
7/72	0.021 $\pm$ 0.007 (8)	0.001 $\pm$ 0.001 (6)	8.4 $\pm$ 3.4 (6)	—	705 $\pm$ 569 (6)	1.83 $\pm$ 0.42 (6)
9/72	0.014 $\pm$ 0.011 (4)	0.024 $\pm$ 0.008 (4)	18.3 $\pm$ 2.0 (4)	—	241 $\pm$ 110 (4)	3.46 $\pm$ 1.40 (4)
8/74	0.025 $\pm$ 0.010 (5)	0.007 $\pm$ 0.004 (6)	8.72 $\pm$ 1.43 (6)	27.4 $\pm$ 23.0 (6)	821 $\pm$ 319 (6)	4.24 $\pm$ 1.33 (6)
Alsea Bay						
6/70	0.050 $\pm$ 0.010 (3)	0.030 $\pm$ 0.009 (3)	—	—	—	—
Umpqua River Estuary						
1/72	0.017 $\pm$ 0.004 (9)	0.016 $\pm$ 0.006 (6)	9.04 $\pm$ 0.89 (6)	—	1378 $\pm$ 441 (6)	2.97 $\pm$ 0.49 (6)
4/72	0.019 $\pm$ 0.004 (4)	0.033 $\pm$ 0.006 (4)	8.6 $\pm$ 1.4 (4)	15.8 $\pm$ 11.8 (4)	714 $\pm$ 193 (4)	2.02 $\pm$ 0.21 (4)
7/72	0.012 $\pm$ 0.004 (4)	0.043 $\pm$ 0.014 (4)	11.8 $\pm$ 1.6 (4)	—	1473 $\pm$ 1111 (4)	3.21 $\pm$ 1.41 (4)
10/72	0.031 $\pm$ 0.014 (4)	0.034 $\pm$ 0.007 (4)	17.3 $\pm$ 3.5 (4)	—	—	—
Coos Bay						
3/71	0.074 $\pm$ 0.022 (6)	0.019 $\pm$ 0.004 (3)	10.6 $\pm$ 3.1 (3)	—	—	—
4/72	0.020 $\pm$ 0.004 (4)	0.007 $\pm$ 0.003 (4)	11.3 $\pm$ 1.6 (4)	18.9 $\pm$ 15.6 (4)	524 $\pm$ 118 (4)	2.34 $\pm$ 0.77 (4)
7/72	0.032 $\pm$ 0.012 (4)	0.016 $\pm$ 0.003 (4)	11.9 $\pm$ 1.4 (4)	34.7 $\pm$ 26.5 (4)	300 $\pm$ 144 (4)	2.23 $\pm$ 0.92 (4)
9/72	0.028 $\pm$ 0.006 (4)	0.018 $\pm$ 0.011 (4)	20.9 $\pm$ 2.3 (4)	—	—	—
Coquille Bay						
3/71	0.063 $\pm$ 0.012 (5)	0.027 $\pm$ 0.007 (5)	7.82 $\pm$ 0.42 (5)	—	—	—

<sup>a</sup> Total soft tissues. Mean  $\pm$  S.D. Number of animals analyzed indicated in parentheses

other decapods. Since there is little loss of metals from estuarine systems (Turekian 1977), the results suggest that Oregon estuaries are relatively free of man-made Cd enrichments.

### Molluscs

The metal profiles for *C. nuttallii* and *M. arenaria* are similar to one another (Table 2) although the concentrations of Mn, Fe, and Cu, were generally higher in *M. arenaria* than in *C. nuttallii*. Of the other molluscs examined, *Anodonta* sp., *C. fluminea* and *M. californianus* tended to have higher tissue Zn concentrations, and *M. edulis*, *P. staminea* and *T. capax* had similar Zn concentrations as

*M. arenaria* and *C. nuttallii* (Tables 2 and 6). The concentrations of Hg were relatively alike in all bivalve molluscs examined and ranged from a low of 0.009  $\mu\text{g/g}$  in *T. capax* to a high of 0.041  $\mu\text{g/g}$  in *P. staminea* (Table 6). In *C. nuttallii* and *M. arenaria* Hg concentrations were 0.019 and 0.025  $\mu\text{g/g}$ , respectively (Table 2).

Molluscan Cd concentrations were substantially more variable and ranged from a low of 0.001  $\mu\text{g/g}$  in two collections of *M. arenaria* and one of *C. nuttallii* to a high of 1.99  $\mu\text{g/g}$  in the turban snail *T. funebris* (Table 6). The mean Cd concentrations from all collections of *C. nuttallii* and *M. arenaria* were near the lower end of the range of concentrations found in molluscs in this study, 0.022 and 0.028  $\mu\text{g/g}$ , respectively (Table 2).

**Table 5.** Concentration of metals in cockles (*Clinocardium nuttallii*) from Oregon estuaries (expressed as  $\mu\text{g/g}$  wet weight)<sup>a</sup>

Collection site and date (Mo/Yr)	Hg	Cd	Zn	Mn	Fe	Cu
Tillamook Bay						
1/72	0.019 $\pm$ 0.006 (8)	0.011 $\pm$ 0.007 (6)	8.67 $\pm$ 1.61 (6)	—	—	—
4/72	0.018 $\pm$ 0.005 (4)	0.023 $\pm$ 0.004 (4)	9.07 $\pm$ 0.58 (4)	13.2 $\pm$ 1.2 (4)	1288 $\pm$ 115 (4)	1.94 $\pm$ 0.14 (4)
7/72	0.029 $\pm$ 0.019 (4)	0.026 $\pm$ 0.012 (4)	11.1 $\pm$ 0.2 (4)	11.9 $\pm$ 3.7 (3)	949 $\pm$ 323 (3)	2.09 $\pm$ 0.03 (3)
10/72	0.020 $\pm$ 0.004 (4)	0.018 $\pm$ 0.007 (4)	13.0 $\pm$ 2.7 (4)	7.63 $\pm$ 3.02 (4)	738 $\pm$ 226 (4)	1.53 $\pm$ 0.61 (4)
Netarts Bay						
2/71	0.019 (1)	0.20 (1)	7.83 (1)	—	—	—
7/72	0.014 $\pm$ 0.004 (4)	0.045 $\pm$ 0.005 (4)	11.0 $\pm$ 1.4 (4)	6.35 $\pm$ 2.59 (4)	161 $\pm$ 37 (4)	0.98 $\pm$ 0.02 (4)
Yaquina Bay						
12/71	0.011 $\pm$ 0.005 (6)	0.006 $\pm$ 0.006 (6)	6.91 $\pm$ 1.13 (6)	—	225 $\pm$ 62 (4)	1.32 $\pm$ 0.24 (4)
4/72	0.024 $\pm$ 0.006 (10)	0.006 $\pm$ 0.006 (6)	10.1 $\pm$ 1.0 (6)	—	613 $\pm$ 76 (5)	1.44 $\pm$ 0.10 (5)
7/72	0.024 $\pm$ 0.007 (8)	0.005 $\pm$ 0.004 (6)	8.9 $\pm$ 1.0 (6)	—	254 $\pm$ 69 (6)	1.30 $\pm$ 0.45 (6)
9/72	0.012 $\pm$ 0.005 (4)	0.013 $\pm$ 0.003 (4)	12.6 $\pm$ 4.1 (4)	3.63 $\pm$ 2.51 (4)	395 $\pm$ 71 (4)	0.99 $\pm$ 0.05 (4)
8/74	0.012 $\pm$ 0.007 (2)	0.001 $\pm$ 0.001 (2)	9.41 $\pm$ 0.81 (2)	3.32 $\pm$ 0.44 (2)	363 $\pm$ 7 (2)	1.17 $\pm$ 0.40 (2)
Coos Bay						
1/72	0.014 $\pm$ 0.005 (6)	0.015 $\pm$ 0.012 (5)	9.10 $\pm$ 0.87 (5)	1.93 $\pm$ 0.23 (5)	271 $\pm$ 61 (5)	0.75 $\pm$ 0.16 (5)
4/72	0.033 $\pm$ 0.004 (4)	0.040 $\pm$ 0.010 (4)	11.6 $\pm$ 2.2 (4)	—	—	—
7/72	0.011 $\pm$ 0.006 (4)	0.080 $\pm$ 0.022 (4)	8.88 $\pm$ 1.32 (4)	3.67 $\pm$ 0.79 (4)	200 $\pm$ 116 (4)	0.97 $\pm$ 0.05 (4)
9/72	0.012 $\pm$ 0.006 (4)	0.021 $\pm$ 0.002 (4)	13.3 $\pm$ 1.8 (4)	2.33 $\pm$ 1.65 (4)	215 $\pm$ 70 (4)	0.99 $\pm$ 0.03 (4)

<sup>a</sup> Total soft tissues. Mean  $\pm$  S.D. Number of animals analyzed indicated in parentheses

Compared with molluscs from other geographic areas, Oregon molluscs were typical with respect to the concentrations of Zn, Mn, Cu, and Fe. For example, Pringle *et al.* (1968) found mean Zn concentrations of 17 and 20.6  $\mu\text{g/g}$  in *M. arenaria* and *Mercenaria mercenaria*, respectively, from the East coast of the U.S. compared to mean Zn concentrations of 8 to 59  $\mu\text{g/g}$  for Oregon coast bivalves (Tables 2 and 6). Similar comparisons between the present findings and those of Pringle *et al.* (1968) can be made for Cu and Mn. In addition, Cu and Zn concentrations in *C. nuttallii* and *M. arenaria* are comparable to those found in Atlantic coast surf clams, *Spisula solidissima*, and ocean quahogs, *Arctica islandica* (Wenzloff *et al.* 1979). Pringle *et al.* (1968) reported a mean Fe concentration of 405  $\mu\text{g/g}$  in East coast *M. arenaria*, which compares favorably to values of 460  $\mu\text{g/g}$  and 934  $\mu\text{g/g}$  in *C. nuttallii* and *M. arenaria* (Table 2) from Oregon. However, East coast quahogs (*M. mercenaria*) contained only 30  $\mu\text{g/g}$  of Fe (Pringle *et al.* 1968).

The concentrations of Hg in the eight species of bivalve molluscs analyzed in this study, averaging about 0.02  $\mu\text{g/g}$  (Tables 2 and 6), were very comparable to those reported by other authors; for example, *Mya arenaria* and *Mercenaria mercenaria* (Freeman *et al.* 1974; Zook *et al.* 1976), *Crassostrea virginica* (Freeman *et al.* 1974; Reimer and Reimer 1975), *Mytilus californianus* (Eganhouse and Young 1976), and *Anodonta grandis* (Smith and Green 1975). In contrast, Schell and Nevisi (1977)

reported Hg concentrations in *Mytilus edulis* and clams from Puget Sound, Washington of approximately 0.002 to 0.004  $\mu\text{g/g}$ , values 5 to 10-fold lower than reported here. The reason for this difference is unknown.

Compared to the concentrations of Cd seen in bivalve molluscs from other geographic areas, Oregon bivalves tended to have lower Cd levels. The mean concentrations of Cd in *C. nuttallii* and *M. arenaria* from Oregon were only 0.022 and 0.028  $\mu\text{g/g}$  (Table 2) compared with mean values from 0.184 to 0.27  $\mu\text{g/g}$  for East coast *Mya arenaria* (Pringle *et al.* 1968; Zook *et al.* 1976) and from 0.122 to approximately 0.25  $\mu\text{g/g}$  for East coast *Mercenaria mercenaria* (Pringle *et al.* 1968; Zook *et al.* 1976). Other Oregon bivalves generally had higher Cd levels than *C. nuttallii* and *M. arenaria*, concentration means up to 0.283  $\mu\text{g/g}$  (Table 6), but were still appreciably lower than values reported elsewhere (Eisler *et al.* 1972; Eustace, 1974; Pringle *et al.* 1968; Wenzloff *et al.* 1979; Zook *et al.* 1976). The unusually low Cd levels found in bivalves from Oregon bays presumably reflects the general lack of industrial pollution and/or geological inputs in these estuaries.

## Conclusions

Concentrations of several toxic metals in Oregon were well below the maximum concentrations gen-

**Table 6.** Concentration of metals in miscellaneous Oregon coastal species (expressed as  $\mu\text{g/g}$  wet weight)<sup>a</sup>

Collection site and date (Mo/Yr)	Hg	Cd	Zn
<i>Anodonta</i> sp.			
Columbia River			
2/72	0.023 (1)	0.042 (1)	25.5 (1)
4/72	0.025 $\pm$ 0.012 (4)	0.148 $\pm$ 0.065 (4)	51.7 $\pm$ 25.4 (4)
7/72	0.033 $\pm$ 0.023 (4)	0.132 $\pm$ 0.029 (4)	51.1 $\pm$ 12.5 (4)
10/72	0.033 $\pm$ 0.023 (4)	—	—
<i>Callianassa californiensis</i> <sup>b</sup>			
Netarts Bay			
2/71	0.014 <sup>d</sup>	—	—
<i>Corbicula fluminea</i>			
Columbia River			
2/72	0.019 $\pm$ 0.023 (8)	0.009 $\pm$ 0.010 (6)	12.0 $\pm$ 3.5 (6)
4/72	0.016 $\pm$ 0.005 (4)	0.069 $\pm$ 0.005 (4)	13.4 $\pm$ 1.7 (4)
7/72	0.018 $\pm$ 0.008 (4)	0.103 $\pm$ 0.006 (4)	26.3 $\pm$ 3.9 (4)
<i>Crangon franciscorum</i> <sup>b</sup>			
Tillamook Bay			
2/71	0.059 <sup>e</sup>	—	—
<i>Hemigrapsus nudus</i> <sup>c</sup>			
Oregon Coast, Brookings			
3/71	0.056 $\pm$ 0.023 (5)	0.032 $\pm$ 0.007 (3)	37.9 $\pm$ 0.7 (3)
<i>Mytilus californianus</i>			
Oregon Coast, Pistol River			
3/71	0.020 $\pm$ 0.005 (6)	0.147 $\pm$ 0.065 (4)	17.8 $\pm$ 8.1 (4)
<i>Mytilus edulis</i>			
Umpqua River Estuary			
1/72	0.016 $\pm$ 0.005 (16)	0.283 $\pm$ 0.147 (6)	9.70 $\pm$ 2.12 (6)
7/72	0.011 $\pm$ 0.002 (4)	0.262 $\pm$ 0.101 (4)	8.0 $\pm$ 0.7 (4)
10/72	0.018 $\pm$ 0.005 (4)	0.240 $\pm$ 0.081 (4)	17.8 $\pm$ 3.5 (4)
<i>Nucella lamellosa</i>			
Tillamook Bay			
2/71	0.035 $\pm$ 0.033 (18)	—	—
<i>Protothaca staminea</i>			
Netarts Bay			
2/71	0.015 $\pm$ 0.004 (5)	0.082 $\pm$ 0.014 (5)	8.62 $\pm$ 0.96 (5)
Oregon Coast, Brookings			
3/71	0.041 $\pm$ 0.012 (5)	0.202 $\pm$ 0.027 (5)	—
<i>Tegula funebris</i>			
Oregon Coast, Brookings			
3/71	0.039 $\pm$ 0.012 (16)	1.99 $\pm$ 2.14 (4)	—
<i>Tresus capax</i>			
Netarts Bay			
2/71	0.009 (1)	0.025 (1)	7.82 (1)

<sup>a</sup> Total soft tissues unless indicated otherwise. Mean  $\pm$  S.D. Number of animals analyzed indicated in parentheses<sup>b</sup> Abdominal muscle<sup>c</sup> Claw muscle<sup>d</sup> Pooled sample from 20 animals<sup>e</sup> Pooled sample from 39 animals

erally regarded as safe for human consumption. These limits, set by the United States Food and Drug Administration for Hg and by the National Health and Medical Research Council of Australia (MacKay *et al.* 1975) for Cd, Cu, and Zn, are 1.0, 2.0, 30 and 1,000  $\mu\text{g/g}$  wet weight, respectively. The levels of these metals found in shellfish from Oregon bays, in this study, are among the lowest levels reported in the literature. The low values and the absence of any significant correlations between loca-

tions within Oregon estuaries suggest that the levels represent normal background concentrations of metals in the organisms examined and do not reflect a measurable degree of contamination from anthropogenic sources.

**Acknowledgments.** This research was supported in part by NIH research grants ES-00040 and ES-00210 and in part by the Oregon State University Sea Grant College Program Grant 04-3-



158-4, NOAA, U.S. Department of Commerce. Issued as Oregon Agricultural Experiment Station Technical Paper No. 5563. The authors are grateful to Mr. L. H. Heaton and Mr. R. O. Sinnhuber for some of the samples and to Ms. M. C. Henderson, Ms. B. A. Hobe, Ms. Rebecca Roberts, Ms. R. Wasserman, and Ms. M. E. Rasmuson for analytical assistance.

## References

- Anonymous*: Oregon Estuaries. Oregon Division of State Lands, Salem, Oregon, (1973).
- Bourke, R. H., B. Glenne, and B. W. Adams: The nearshore physical oceanographic environment of the Pacific Northwest coast. Department of Oceanography, Oregon State University, Corvallis, OR (1971).
- Bryan, G. W.: Zinc regulation in the lobster, *Homarus vulgaris*. I. Tissue zinc and copper concentrations. J. Mar. Biol. Assoc. U.K. **44**, 549 (1964).
- : The effects of heavy metals (other than mercury) on marine and estuarine organisms. Proc. Roy. Soc. (Ser. B) **177**, 389 (1971).
- : The occurrence and seasonal variation of trace metals in the scallops *Pecten maximus* (L.) and *Chlamys opercularis* (L.). J. Mar. Biol. Assoc. U.K. **53**, 145 (1973).
- Buhler, D. R., R. R. Claeys, and B. R. Mate: Heavy metal and chlorinated hydrocarbon residues in California sea lions (*Zalophus californianus californianus*). J. Fish. Res. Bd. Can. **32**, 239 (1975).
- Eganhouse, R. P., and D. R. Young: Mercury in tissues of mussels off southern California. Mar. Pollut. Bull. **7**, 145 (1976).
- Eisler, R.: Cadmium poisoning in *Fundulus heteroclitus* (Pisces: Cyprinodontidae) and other marine organisms. J. Fish. Res. Bd. Can. **28**, 1225 (1971).
- Eisler, R., G. E. Zarogian, and R. J. Hennekey: Cadmium uptake by marine organisms. J. Fish. Res. Bd. Can. **29**, 1367 (1972).
- Eustace, I. J.: Zinc, cadmium, copper, and manganese in species of finfish and shellfish caught in the Derwent Estuary, Tasmania. Aust. J. Mar. Freshwat. Res. **25**, 209 (1974).
- Fowler, B. A., R. C. Fay, R. L. Walter, R. D. Willis, and W. F. Gutknecht: Levels of toxic metals in marine organisms collected from southern California coastal waters. Environ. Health Perspec. **12**, 71 (1975).
- Freeman, H. C., D. A. Horne, B. McTague, and M. McMenemy: Mercury in some Canadian Atlantic coast fish and shellfish. J. Fish. Res. Bd. Can. **31**, 369 (1974).
- Koirtiyohann, S. R., and J. W. Wen: Critical study of the APDC-MIBK extraction system for atomic absorption. Anal. Chem. **45**, 1986 (1973).
- MacKay, N. J., R. J. Williams, J. L. Kacprzac, M. N. Kazacos, A. J. Collins, and E. H. Auty: Heavy metals in cultivated oysters (*Crassostrea commercialis* = *Saccostrea cucullata*) from the estuaries of New South Wales. Aust. J. Mar. Freshwater Res. **26**, 31 (1975).
- Percy, K. L., C. Sutterlin, D. A. Bella, and P. C. Klingeman: Descriptions and information sources for Oregon estuaries. Sea Grant College Program, Oregon State University, Corvallis, OR (1974).
- Phillips, D. J. H.: The common mussel *Mytilus edulis* as an indicator of pollution by zinc, cadmium, lead, and copper. I. Effect of environmental variables on uptake of metals. Marine Biol. **38**, 59 (1976).
- Phillips, G. R., and D. R. Buhler: Mercury accumulation in and growth rate of rainbow trout, *Salmo gairdneri*, stocked in an eastern Oregon reservoir. Arch. Environ. Contam. Toxicol. **9**, 99 (1980).
- Preston, A.: Heavy metals in British waters. Nature **242**, 95 (1973).
- Pringle, G. H., D. E. Hissong, E. L. Katz, and S. T. Mulawka: Trace metal accumulation by estuarine mollusks. Am. Soc. Civil Eng. Proc. J. Sanit. Eng. Div. **94**(SA3), Paper 5970, 455 (1968).
- Reimer, A. A., and R. D. Reimer: Total mercury in some fish and shellfish along the Mexican coast. Bull. Environ. Contam. Toxicol. **14**, 105 (1975).
- Schell, W. R., and A. Nevissi: Heavy metals from waste disposal in Central Puget Sound. Environ. Sci. Technol. **11**, 887 (1977).
- Smith, A. L., and R. H. Green: Uptake of mercury by freshwater clams (Family Unionidae). J. Fish. Res. Bd. Can. **32**, 1297 (1975).
- Takeuchi, T.: Biological reactions and pathological changes in human beings and animals caused by organic mercury contamination. In R. Hartung and B. D. Dinman (eds.): Environmental mercury contamination. p. 247. Ann Arbor, MI. Ann Arbor Science Publishers (1972).
- Taylor, D.: Natural distribution of trace metals in sediments from a coastal environment, Tor Bay, England. Estuar. Coastal Mar. Sci. **2**, 417 (1974).
- Thurberg, F. P., M. A. Dawson, and R. S. Collier: Effects of copper and cadmium on osmoregulation and oxygen consumption in two species of estuarine crabs. Mar. Biol. **23**, 171 (1973).
- Topping, G.: Heavy metals in shellfish from Scottish waters. Aquacult. **1**, 379 (1973).
- Turekian, K. K.: The fate of metals in the oceans. Geochim. Cosmochim. Acta **41**, 1139 (1977).
- Wenzloff, D. R., R. A. Greig, A. S. Merrill, and J. W. Ropes: A survey of heavy metals in the surf clam, *Spisula solidissima*, and the ocean quahog, *Artica islandica*, of the Mid-Atlantic coast of the United States. Fish. Bull. **77**, 280 (1979).
- Young, D. R., C. S. Young, and G. E. Hlavka: Sources of trace metals from highly-urbanized Southern California to the adjacent marine ecosystems. In Cycling and Control of Metals. p. 21. U.S. Environmental Protection Agency, Cincinnati, OH (1973).
- Zook, E. G., J. J. Powell, B. M. Hackley, J. A. Emerson, J. R. Brooker, and G. M. Knobl, Jr.: National Marine Fisheries Service preliminary survey of selected sea foods for mercury, lead, cadmium, chromium, and arsenic content. J. Agr. Food Chem. **24**, 47 (1976).

Manuscript received July 25, 1981; accepted April 22, 1982.