References

- Barfield, C. S., Mitchell, E., and Poeb, S. (1978). A temperature-dependent model for fall armyworm development. Annals of the Entomological Society of America, 71(1):70–74.
- Butler Jr, G., Bryan, D., and Jackson, C. (1968). Development of the salt-marsh caterpillar parasite, exorista mella at controlled constant and variable temperatures in the laboratory. *Journal of economic entomology*, 61(1):161–162.
- Butler Jr, G. and Lopez, J. (1980). Trichogramma pretiosum: development in two hosts in relation to constant and fluctuating temperatures. *Annals of the Entomological Society of America*, 73(6):671–673.
- Butler Jr, G. D. (1966). Development of the beet armyworm and its parasite chelonus texanus in relation to temperature. *Journal of economic entomology*, 59(6):1324–1327.
- Davis, J., Radcliffe, E., and Ragsdale, D. (2006). Effects of high and fluctuating temperatures on myzus persicae (hemiptera: Aphididae). *Environmental Entomology*, 35(6):1461–1468.
- Eubank, W., Atmar, J., and Ellington, J. (1973). The significance and thermodynamics of fluctuating versus static thermal environments on heliothis zea egg development rates. *Environmental Entomology*, 2(4):491–496.
- Fielding, D. J. and Ruesink, W. G. (1988). Prediction of egg and nymphal developmental times of the squash bug (hemiptera: Coreidae) in the field. *Journal of economic entomology*, 81(5):1377–1382.
- Hagstrum, D. W. and Milliken, G. A. (1991). Modeling differences in insect developmental times between constant and fluctuating temperatures. *Annals of the Entomological Society of America*, 84(4):369–379.
- Harries, F. and Douglass, J. (1948). Bionomic studies on the beet leafhopper. *Ecological Monographs*, 18(1):45–79.
- Joshi, D. (1996). Effect of fluctuating and constant temperatures on development, adult longevity and fecundity in the mosquito aedes krombeini. *Journal of Thermal Biology*, 21(3):151–154.
- Lin, S., Hodson, A. C., and Richards, A. G. (1954). An analysis of threshold temperatures for the development of oncopeltus and tribolium eggs. *Physiological Zoology*, 27(4):287–311.
- Liu, S.-S., Zhang, G.-M., and Zhu, J. (1995). Influence of temperature variations on rate of development in insects: analysis of case studies from entomological literature. *Annals of the Entomological Society of America*, 88(2):107–119.
- Ludwig, D. (1928). The effects of temperature on the development of an insect (popillia japonica newman). *Physiological Zoology*, 1(3):358–389.
- Ludwig, D. and Cable, R. M. (1933). The effect of alternating temperatures on the pupal development of drosophila melanogaster meigen. *Physiological Zoology*, 6(4):493–508.
- Messenger, P. (1969). Bioclimatic studies of the aphid parasite praon exsoletum. 2. thermal limits to development and effects of temperature on rate of development and occurrence of diapause. *Annals of the Entomological Society of America*, 62(5):1026–1031.
- Messenger, P. and Flitters, N. (1958). Effect of constant temperature environments on the egg stage of three species of hawaiian fruit flies. *Annals of the Entomological Society of America*, 51(2):109–119.
- Niehaus, A. C., Angilletta, M. J., Sears, M. W., Franklin, C. E., and Wilson, R. S. (2012). Predicting the physiological performance of ectotherms in fluctuating thermal environments. *Journal of Experimental Biology*, 215(4):694–701.
- Siddiqui, W. and Barlow, C. (1972). Population growth of drosophila melanogaster (diptera: Drosophilidae) at constant and alternating temperatures. *Annals of the Entomological Society of America*, 65(5):993–1001.
- Simonet, D. and Davenport, B. (1981). Temperature requirements for development and oviposition of the carrot weevil. *Annals of the Entomological Society of America*, 74(3):312–315.