Currying and the Lambda Calculus

CS 350

Dr. Joseph Eremondi

Last updated: July 19, 2024

Overview

• To learn how multi-argument functions can be desugared into single-argument functions

- To learn how multi-argument functions can be desugared into single-argument functions
 - Curly-Curry

- To learn how multi-argument functions can be desugared into single-argument functions
 - o Curly-Curry
- To see that everything can be desugared into single-argument functions

- To learn how multi-argument functions can be desugared into single-argument functions
 - Curly-Curry
- To see that everything can be desugared into single-argument functions
 - by learning about the Lambda Calculus

Details

 Say we allow {lambda $\{x\ y\ z\}\ \{+\ x\ \{*\ y\ z\}\}\}$ in Curly

- Say we allow {lambda $\{x\ y\ z\}\ \{+\ x\ \{*\ y\ z\}\}\}$ in Curly
- How can we interpret a call to this function?

- Say we allow {lambda $\{x \ y \ z\} \ \{+ \ x \ \{* \ y \ z\}\}\}$ in Curly
- How can we interpret a call to this function?
 - o Evaluate the body with either

- Say we allow {lambda $\{x \ y \ z\} \ \{+ \ x \ \{* \ y \ z\}\}\}$ in Curly
- How can we interpret a call to this function?
 - Evaluate the body with either
 - x, y, z replaced by concrete argument values (substitution)

- Say we allow {lambda $\{x \ y \ z\} \ \{+ \ x \ \{* \ y \ z\}\}\}$ in Curly
- How can we interpret a call to this function?
 - Evaluate the body with either
 - x, y, z replaced by concrete argument values (substitution)
 - x,y,z bound to concrete values in an environment

• We can achieve this with nested lambda expressions

• We can achieve this with nested lambda expressions

• We can achieve this with nested lambda expressions

```
{let f {fun {x} {fun {y} {fun {z} {+ x {* y z}}}}} ....}
```

• We can achieve this with nested lambda expressions

```
{let f {fun {x} {fun {y} {fun {z} {+ x {* y z}}}}} ....}
```

We can achieve this with nested lambda expressions

```
{let f {fun {x} {fun {y} {fun {z} {+ x {* y z}}}}} ....}
```

• To call, we do nested calls

```
{{{f 1} 2} 3}
```

Step by step:

We can achieve this with nested lambda expressions

```
{let f {fun {x} {fun {y} {fun {z} {+ x {* y z}}}}} ....}
```

```
{{{f 1} 2} 3}
```

- Step by step:
 - o f 1 produces {fun {y} {fun {z} {+ 1 {* y z}}}}

We can achieve this with nested lambda expressions

```
{let f {fun {x} {fun {y} {fun {z} {+ x {* y z}}}}} ....}
```

```
{{{f 1} 2} 3}
```

- Step by step:
 - o f 1 produces {fun {y} {fun {z} {+ 1 {* y z}}}}
 - \circ Calling that on 2 produces {fun {z} {+ 1 {* 2 z}}}

We can achieve this with nested lambda expressions

```
{let f {fun {x} {fun {y} {fun {z} {+ x {* y z}}}}} ....}
```

```
{{{f 1} 2} 3}
```

- Step by step:
 - o f 1 produces {fun {y} {fun {z} {+ 1 {* y z}}}}
 - \circ Calling that on 2 produces $\{fun \{z\} \{+ 1 \{* 2 z\}\}\}\$
 - \circ Calling that on 3 produces $\{+1 \ \{*2 \ 3\}\}\}$

We can achieve this with nested lambda expressions

```
{let f {fun {x} {fun {y} {fun {z} {+ x {* y z}}}}} ....}
```

```
{{{f 1} 2} 3}
```

- Step by step:
 - o f 1 produces {fun {y} {fun {z} {+ 1 {* y z}}}}
 - \circ Calling that on 2 produces $\{fun \{z\} \{+ 1 \{* 2 z\}\}\}\$
 - Calling that on 3 produces {+ 1 {* 2 3}}}
 - Exactly what we want for {f 1 2 3}