Template SFD E(B-V), |b| > 25° (Uchuu full sky, NSIDE=64): the same systematic matters less for the broader, more biased samples

Contamination (t: template, mean 0, std 1)

additive: $N_c = N + \varepsilon \bar{N} t$

multiplicative: $N_c = N(1 + \varepsilon t)$

Fit and correction with strength k

$\hat{\varepsilon} = \sum (N_c/\bar{N} - 1)\, t / \sum t^2$

add.: $N_k = N_c - k\hat{\varepsilon}\bar{N}t$; mult.: $N_k = N_c/(1 + k\hat{\varepsilon}t)$

k = fraction of the fitted systematic removed: 0 none, 0.5 half (under), 1 all (nominal), 1.5 50% too much (over: an inverted copy is imprinted)

Clustering width, shot noise removed (Poisson-log-normal moments)

$\sigma^2 = \ln\left[1 + (\mathrm{Var}\, N_k - \bar{N})/\bar{N}^2\right]$

Width to bias to halo mass

$\sigma^2 = \ln(1 + b^2 \sigma^2_{m,\mathrm{pix}}) \;\Rightarrow\; b = \sqrt{e^{\sigma^2} - 1}\,/\,\sigma_{m,\mathrm{pix}}$

$b(M_{\min})$: Comparat17 averaged over $n(>M_{\min})$























