Summary of symmetry calculations

November 10, 2021

Contents

1	DBH_model	5
2	Lotka_Volterra	7
3	linear_model	9
4	hydons_model	13

4 CONTENTS

DBH_{model}

Run $04_15PM_10_November-2021$

Degree in tangential ansätze: 2. The system of ODEs is given by:

$$\begin{aligned} \frac{\mathrm{d}w_1}{\mathrm{d}t} &= -w_1w_2 - w_1w_3 + w_2w_3, \\ \frac{\mathrm{d}w_2}{\mathrm{d}t} &= -w_1w_2 + w_1w_3 - w_2w_3, \\ \frac{\mathrm{d}w_3}{\mathrm{d}t} &= w_1w_2 - w_1w_3 - w_2w_3. \end{aligned}$$

The calculated generators are:

$$X_1 = (t^2) \partial t + (1 - 2tw_1) \partial w_1 + (1 - 2tw_2) \partial w_2 + (1 + 2tw_3) \partial w_3$$

$$X_2 = (-t) \partial t + (w_1) \partial w_1 + (w_2) \partial w_2 + (w_3) \partial w_3,$$

$$X_3 = (1) \partial t$$
,

$$X_4 = (1) \partial t$$
,

$$X_5 = (\mathbf{f}_1\left(t\right))\,\partial t + (w_2w_3\,\mathbf{f}_1\left(t\right) - w_1w_2\,\mathbf{f}_1\left(t\right) - w_1w_3\,\mathbf{f}_1\left(t\right))\,\partial w_1 + (w_1w_3\,\mathbf{f}_1\left(t\right) - w_1w_2\,\mathbf{f}_1\left(t\right) \\ + -w_2w_3\,\mathbf{f}_1\left(t\right))\,\partial w_2 + (w_1w_2\,\mathbf{f}_1\left(t\right) - w_1w_3\,\mathbf{f}_1\left(t\right) - w_2w_3\,\mathbf{f}_1\left(t\right))\,\partial w_3.$$

Some of the generators might contain the following arbitrary functions:

 f_1

The execution time of the script was:

 $0~\mathrm{hours}~5~\mathrm{minutes}~7~\mathrm{seconds}.$

Lotka_Volterra

Run $04_16PM_10_November-2021$

Degree in tangential ansätze: 2. The system of ODEs is given by:

$$\frac{\mathrm{d}N}{\mathrm{d}t} = N\left(-Pb + a\right),$$

$$\frac{\mathrm{d}P}{\mathrm{d}t} = P\left(Nc - d\right).$$

The calculated generators are:

$$X_1 = (1) \partial t$$
,

$$X_2 = \left(-\frac{1}{a}\right)\partial t,$$

$$X_{3} = \left(\frac{\mathbf{f}_{1}\left(t\right)}{c}\right)\partial t + \left(\frac{Na\,\mathbf{f}_{1}\left(t\right)}{c} - \frac{NPb\,\mathbf{f}_{1}\left(t\right)}{c}\right)\partial N + \left(NP\,\mathbf{f}_{1}\left(t\right) - \frac{Pd\,\mathbf{f}_{1}\left(t\right)}{c}\right)\partial P.$$

Some of the generators might contain the following arbitrary functions:

 f_1

The execution time of the script was:

0 hours 1 minutes 8 seconds.

linear_model

Run $04_09PM_10_November-2021$

Degree in tangential ansätze: 2. The system of ODEs is given by:

$$\frac{\mathrm{d}u}{\mathrm{d}t} = u + v,$$
$$\frac{\mathrm{d}v}{\mathrm{d}t} = u + v.$$

The calculated generators are:

$$X_{1} = \left(\frac{e^{2t}}{4} - \frac{e^{-2t}}{4}\right) \partial t + \left(\frac{u}{2} + \frac{ve^{2t}}{2} - \frac{ue^{-2t}}{2} - \frac{ve^{-2t}}{2}\right) \partial u + \left(\frac{v}{2} + \frac{ve^{2t}}{2}\right) \partial v$$

$$\begin{split} X_2 &= \left(\frac{u}{8} + \frac{v}{8} - \frac{3ve^{-2t}}{8} - \frac{ue^{-4t}}{8} - \frac{ue^{2t}}{8} - \frac{ve^{-4t}}{8} \right. \\ &+ \frac{ue^{-2t}}{8} + \frac{3ve^{2t}}{8} \right) \partial t + \left(-\frac{u^2}{4} + \frac{v^2}{4} + \frac{uv}{2} + \frac{u^2e^{-2t}}{2} \right. \\ &+ \frac{v^2e^{2t}}{2} - \frac{v^2e^{-2t}}{2} - \frac{u^2e^{-4t}}{4} - \frac{v^2e^{-4t}}{4} - \frac{uve^{-4t}}{2} \right) \partial u + \left(\frac{v^2}{2} + \frac{v^2e^{2t}}{2} \right) \partial v \end{split}$$

$$X_3 = \left(\frac{v}{2} - \frac{u}{2} + \frac{ue^{-2t}}{2} + \frac{ve^{-2t}}{2}\right)\partial t$$

$$\begin{split} X_4 &= \left(-\frac{3u}{8} + \frac{5v}{8} - \frac{3ue^{-2t}}{8} - \frac{ue^{-4t}}{8} - \frac{ue^{2t}}{8} - \frac{ve^{-4t}}{8} \right. \\ &\quad + \frac{ve^{-2t}}{8} + \frac{3ve^{2t}}{8} \right) \partial t + \left(-\frac{u^2}{4} + \frac{v^2}{4} + \frac{uv}{2} + \frac{v^2e^{-2t}}{2} \right. \\ &\quad + \frac{v^2e^{2t}}{2} - \frac{u^2e^{-2t}}{2} - \frac{u^2e^{-4t}}{4} - \frac{v^2e^{-4t}}{4} - \frac{uve^{-4t}}{2} \right) \partial u + \left(\frac{v^2}{2} + \frac{v^2e^{2t}}{2} \right) \partial v \end{split}$$

$$X_5 = \left(-\frac{e^{2t}}{4} - \frac{1}{2} - \frac{e^{-2t}}{4}\right)\partial t + \left(-\frac{u}{2} - \frac{ue^{-2t}}{2} - \frac{ve^{2t}}{2} - \frac{ve^{-2t}}{2}\right)\partial u + \left(-\frac{ve^{2t}}{2} - \frac{v}{2}\right)\partial v,$$

$$\begin{split} X_6 &= \left(-\frac{u}{8} - \frac{v}{8} - \frac{3ve^{-2t}}{8} - \frac{ue^{2t}}{8} + \frac{ue^{-4t}}{8} + \frac{ue^{-2t}}{8} \right. \\ &+ \left. \frac{ve^{-4t}}{8} + \frac{3ve^{2t}}{8} \right) \partial t + \left(-\frac{v^2}{4} + \frac{u^2}{4} + \frac{u^2e^{-2t}}{2} + \frac{v^2e^{2t}}{2} \right. \\ &+ \left. -\frac{uv}{2} - \frac{v^2e^{-2t}}{2} + \frac{u^2e^{-4t}}{4} + \frac{v^2e^{-4t}}{4} + \frac{uve^{-4t}}{2} \right) \partial u + \left(\frac{v^2e^{2t}}{2} - \frac{v^2}{2} \right) \partial v, \end{split}$$

$$X_7 = \left(\frac{1}{2} + \frac{e^{2t}}{2}\right)\partial u + \left(\frac{e^{2t}}{2} - \frac{1}{2}\right)\partial v,$$

$$X_8 = (1) \partial t$$

$$\begin{split} X_9 &= \left(-\frac{5v}{8} + \frac{3u}{8} - \frac{3ue^{-2t}}{8} - \frac{ue^{2t}}{8} + \frac{ue^{-4t}}{8} + \frac{ve^{-4t}}{8} \right. \\ &+ \left. \frac{ve^{-2t}}{8} + \frac{3ve^{2t}}{8} \right) \partial t + \left(-\frac{v^2}{4} + \frac{u^2}{4} + \frac{v^2e^{-2t}}{2} + \frac{v^2e^{2t}}{2} \right. \\ &+ \left. -\frac{uv}{2} - \frac{u^2e^{-2t}}{2} + \frac{u^2e^{-4t}}{4} + \frac{v^2e^{-4t}}{4} + \frac{uve^{-4t}}{2} \right) \partial u + \left(\frac{v^2e^{2t}}{2} - \frac{v^2}{2} \right) \partial v, \end{split}$$

$$\begin{split} X_{10} = & \left(-\frac{3v}{8} + \frac{u}{8} - \frac{3ve^{2t}}{8} - \frac{ue^{-4t}}{8} - \frac{ue^{-2t}}{8} - \frac{ve^{-4t}}{8} \right. \\ & + \left. -\frac{ve^{-2t}}{8} + \frac{ue^{2t}}{8} \right) \partial t + \left(\frac{u^2}{4} - \frac{u^2e^{-4t}}{4} - \frac{uv}{2} - \frac{uve^{-4t}}{2} - \frac{v^2e^{2t}}{2} - \frac{v^2}{4} - \frac{v^2e^{-4t}}{4} \right) \partial u + \left(-\frac{v^2e^{2t}}{2} - \frac{v^2}{2} \right) \partial v, \end{split}$$

$$X_{11} = \left(-\frac{u^2}{4} - \frac{v^2}{4} + \frac{uv}{2} + \frac{u^2e^{-4t}}{4} + \frac{v^2e^{-4t}}{4} + \frac{uve^{-4t}}{2}\right)\partial t$$

$$X_{12} = \left(\frac{u^2}{4} + \frac{v^2}{4} + \frac{u^2 e^{-2t}}{2} - \frac{uv}{2} - \frac{v^2 e^{-2t}}{2} + \frac{u^2 e^{-4t}}{4} + \frac{v^2 e^{-4t}}{4} + \frac{uv e^{-4t}}{2}\right) \partial t$$

$$\begin{split} X_{13} = & \left(\frac{u^2}{4} + \frac{v^2}{4} + \frac{v^2 e^{-2t}}{2} - \frac{uv}{2} - \frac{u^2 e^{-2t}}{2} + \frac{u^2 e^{-4t}}{4} + \frac{v^2 e^{-4t}}{4} + \frac{uv e^{-4t}}{2} \right) \partial t \end{split}$$

$$X_{14} = \left(\frac{u}{2} - \frac{v}{2} + \frac{ue^{-2t}}{2} + \frac{ve^{-2t}}{2}\right)\partial t$$

$$X_{15} = \left(\frac{e^{2t}}{2} - \frac{1}{2}\right)\partial u + \left(\frac{1}{2} + \frac{e^{2t}}{2}\right)\partial v$$

$$X_{16} = \left(-\frac{e^{2t}}{4} + \frac{e^{-2t}}{4}\right) \partial t + \left(\frac{u}{2} + \frac{ue^{-2t}}{2} + \frac{ve^{-2t}}{2} - \frac{ve^{2t}}{2}\right) \partial u + \left(\frac{v}{2} - \frac{ve^{2t}}{2}\right) \partial v$$

$$\begin{split} X_{17} = \left(-\frac{1}{2} + \frac{e^{-2t}}{4} + \frac{e^{2t}}{4} \right) \partial t + \left(-\frac{u}{2} + \frac{ue^{-2t}}{2} + \frac{ve^{-2t}}{2} \right) \\ + \frac{ve^{2t}}{2} \right) \partial u + \left(\frac{ve^{2t}}{2} - \frac{v}{2} \right) \partial v, \end{split}$$

$$\begin{split} X_{18} &= \left(-\frac{u}{8} + \frac{3v}{8} - \frac{3ve^{2t}}{8} - \frac{ue^{-2t}}{8} - \frac{ve^{-2t}}{8} + \frac{ue^{-4t}}{8} \right. \\ &+ \left. \frac{ue^{2t}}{8} + \frac{ve^{-4t}}{8} \right) \partial t + \left(-\frac{u^2}{4} + \frac{v^2}{4} + \frac{uv}{2} - \frac{v^2e^{2t}}{2} \right. \\ &+ \left. \frac{u^2e^{-4t}}{4} + \frac{v^2e^{-4t}}{4} + \frac{uve^{-4t}}{2} \right) \partial u + \left(\frac{v^2}{2} - \frac{v^2e^{2t}}{2} \right) \partial v \end{split}$$

$$X_{19} = (u f_2(t) + v f_1(t) - v f_2(t) + f_3(t)) \partial t + (u f_3(t) + v f_3(t) + u^2 f_2(t) + v^2 f_1(t) - v^2 f_2(t) + uv f_1(t)) \partial u + (u f_3(t) + v f_3(t) + u^2 f_2(t) + v^2 f_1(t) - v^2 f_2(t) + uv f_1(t)) \partial v$$

Some of the generators might contain the following arbitrary functions:

 f_1

 f_2

 f_3

The execution time of the script was:

0 hours 1 minutes 9 seconds.

$hydons_model$

$Run\ 04_08PM_10_November-2021$

Degree in tangential ansätze: 2. The system of ODEs is given by:

$$\begin{split} \frac{\mathrm{d}y_1}{\mathrm{d}t} &= \frac{ty_1 + y_2^2}{-t^2 + y_1 y_2}, \\ \frac{\mathrm{d}y_2}{\mathrm{d}t} &= \frac{ty_2 + y_1^2}{-t^2 + y_1 y_2}. \end{split}$$

The calculated generators are:

$$X_1 = (t) \partial t + (y_1) \partial y_1 + (y_2) \partial y_2,$$

$$X_{2}=\left(-t^{2} \operatorname{f}_{1}\left(t\right)+y_{1} y_{2} \operatorname{f}_{1}\left(t\right)\right) \partial t+\left(y_{2}^{2} \operatorname{f}_{1}\left(t\right)+t y_{1} \operatorname{f}_{1}\left(t\right)\right) \partial y_{1}+\left(y_{1}^{2} \operatorname{f}_{1}\left(t\right)+t y_{2} \operatorname{f}_{1}\left(t\right)\right) \partial y_{2}$$

Some of the generators might contain the following arbitrary functions:

 f_1

The execution time of the script was:

0 hours 0 minutes 26 seconds.