

Math Methods Assignment #4

Johannes Byle

September 28, 2021

1. It would move in a parabola, which can be seen from the fact that if we rotate our coordinate system such that the rockets total acceleration is pointing down this becomes a simple projectile motion problem, which we know is parabolic.
2. If we take a rhombus with corners A, B, C, D we can represent the sides as $\vec{AB}, \vec{BC}, \vec{CD}, \vec{DA}$. We know that $|\vec{AB}| = |\vec{BC}| = |\vec{CD}| = |\vec{DA}|$, and that $\vec{AC} = \vec{AB} + \vec{BC}$ and $\vec{BD} = \vec{BC} + \vec{CD}$. Since the sides of a rhombus are parallel we can for the sake of this problem consider $\vec{AC} = \vec{AB} + \vec{AD}$ and $\vec{BD} = \vec{AD} - \vec{AB}$. To show that the diagonals are orthogonal we need to show that the dot products are equal to zero: $\vec{AC} \cdot \vec{BD} = (\vec{AB} + \vec{AD}) \cdot (\vec{AD} - \vec{AB}) = \vec{AD}^2 - \vec{AB}^2 = 0$ since the sides all have equal length.
3. This is equivalent to $\sum_i \sum_j \sum_k \epsilon_{ijk} \epsilon_{ijk}$. The number of permutations of n numbers is $n!$, which means there are $3! = 6$ permutations in our case. Since whether the permutation is even or odd the term $\epsilon_{ijk} \epsilon_{ijk} = 1$ and otherwise 0 the sum is equal to 6.
4. Using the definition of the cross product $\mathbf{a} \times \mathbf{b} = \epsilon_{ijk} \hat{\mathbf{e}}_i a_j b_k$:

$$\begin{aligned}\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) &= \mathbf{a} \times (\epsilon_{ijk} \hat{\mathbf{e}}_i b_j c_k) = \epsilon_{ijk} \epsilon_{ijl} \hat{\mathbf{e}}_l a_j b_j c_k \\ (\mathbf{a} \times \mathbf{b}) \times \mathbf{c} &= (\epsilon_{ijk} \hat{\mathbf{e}}_i a_j b_k) \times \mathbf{c} = \epsilon_{ijk} \epsilon_{ijl} \hat{\mathbf{e}}_l a_j b_j c_k\end{aligned}$$

5. The following matrix converts between x and x' :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \quad (1)$$

Multiplying the gradient of the scalar function we can see that they are both equal and thus covariant:

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \hat{e}_1 \\ \hat{e}_2 \end{bmatrix} = \frac{1}{\sqrt{2}} \hat{e}_1 + \left(\frac{1}{\sqrt{2}} + 1 \right) \hat{e}_2 = \frac{1}{\sqrt{2}} (\hat{e}_1 + \hat{e}_2) + \hat{e}_2 \quad (2)$$

6. An orthogonal transformation follows the following rule: $e'_i =$