

Math Methods Assignment #4

Johannes Byle

September 27, 2021

1. It would depend on whether the rocket rotated. Assuming that it's acceleration was constant in both the horizontal and vertical directions it would move in a parabola, unless v_0 is 0 in which case it would move in a straight line. If the rocket were allowed to rotate it would move in a parabolic path, as it would start asymptotically in horizontal direction, and then as it tilted down it would move asymptotically close to the vertical axis.
2. If we take a rhombus with corners A, B, C, D we can represent the sides as $\vec{AB}, \vec{BC}, \vec{CD}, \vec{DA}$. We know that $|\vec{AB}| = |\vec{BC}| = |\vec{CD}| = |\vec{DA}|$, and that $\vec{AC} = \vec{AB} + \vec{BC}$ and $\vec{BD} = \vec{BC} + \vec{CD}$. Since the sides of a rhombus are parallel we can for the sake of this problem consider $\vec{AC} = \vec{AB} + \vec{AD}$ and $\vec{BD} = \vec{AD} - \vec{AB}$. To show that the diagonals are orthogonal we need to show that the dot products are equal to zero: $\vec{AC} \cdot \vec{BD} = (\vec{AB} + \vec{AD}) \cdot (\vec{AD} - \vec{AB}) = \vec{AD}^2 - \vec{AB}^2 = 0$ since the sides all have equal length.
3. This is equivalent to $\sum_i \sum_j \sum_k \epsilon_{ijk} \epsilon_{ijk}$. The number of permutations of n numbers is $n!$, which means there are $3! = 6$ permutations in our case. Since whether the permutation is even or odd the term $\epsilon_{ijk} \epsilon_{ijk} = 1$ and otherwise 0 the sum is equal to 6.
4. Using the definition of the cross product $\mathbf{a} \times \mathbf{b} = \epsilon_{ijk} \hat{\mathbf{e}}_i a_j b_k$:

$$\begin{aligned}\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) &= \mathbf{a} \times (\epsilon_{ijk} \hat{\mathbf{e}}_i b_j c_k) = \epsilon_{ijk} \hat{\mathbf{e}}_i a_j b_j c_k \\ (\mathbf{a} \times \mathbf{b}) \times \mathbf{c} &= (\epsilon_{ijk} \hat{\mathbf{e}}_i a_j b_k) \times \mathbf{c} = \epsilon_{ijk} \hat{\mathbf{e}}_i a_j b_j c_k\end{aligned}$$