

Classical Assignment #7

Johannes Byle

October 27, 2021

1. From Goldstein page 153 we know that we can go from body coordinates to space axes using the following relation $\mathbf{x} = \mathbf{A}^{-1}\mathbf{x}'$. Since $\mathbf{A} = \mathbf{BCD}$:

$$\mathbf{BCD} = \begin{pmatrix} \cos(\phi) & \sin(\phi) & 0 \\ -\sin(\phi) & \cos(\phi) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & \sin(\theta) \\ 0 & -\sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} \cos(\psi) & \sin(\psi) & 0 \\ -\sin(\psi) & \cos(\psi) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\mathbf{A} = \begin{pmatrix} \cos(\psi)\cos(\phi) - \cos(\theta)\sin(\psi)\sin(\phi) & \cos(\theta)\sin(\psi)\cos(\phi) + \cos(\psi)\sin(\phi) & \sin(\theta)\sin(\psi) \\ -\cos(\theta)\cos(\psi)\sin(\phi) - \sin(\psi)\cos(\phi) & \cos(\theta)\cos(\psi)\cos(\phi) - \sin(\psi)\sin(\phi) & \sin(\theta)\cos(\psi) \\ \sin(\theta)\sin(\phi) & \sin(\theta)(-\cos(\phi)) & \cos(\theta) \end{pmatrix}$$

$$\mathbf{A}^{-1} = \begin{pmatrix} \cos(\psi)\cos(\phi) - \cos(\theta)\sin(\psi)\sin(\phi) & -\cos(\theta)\cos(\psi)\sin(\phi) - \sin(\psi)\cos(\phi) & \sin(\theta)\sin(\phi) \\ \cos(\theta)\sin(\psi)\cos(\phi) + \cos(\psi)\sin(\phi) & \cos(\theta)\cos(\psi)\cos(\phi) - \sin(\psi)\sin(\phi) & \sin(\theta)(-\cos(\phi)) \\ \sin(\theta)\sin(\psi) & \sin(\theta)\cos(\psi) & \cos(\theta) \end{pmatrix}$$

$$\omega_{bf} = \begin{pmatrix} \theta' \cos(\psi) + \sin(\theta)\sin(\psi)\phi' \\ \sin(\theta)\cos(\psi)\phi' - \theta' \sin(\psi) \\ \cos(\theta)\phi' + \psi' \end{pmatrix}$$

$$\mathbf{A}^{-1}\omega_{bf} = \begin{pmatrix} \theta' \cos(\phi) + \sin(\theta)\psi' \sin(\phi) \\ \theta' \sin(\phi) - \sin(\theta)\psi' \cos(\phi) \\ \cos(\theta)\psi' + \phi' \end{pmatrix}$$

2. (a) Simply plugging in equation (5.22) from Goldstein into Mathematica:

```
Clear[Global*]
x = {r Cos[θ], r Sin[θ], z};
f[j_, k_] := (x[[1]]^2 + x[[2]]^2 + z^2) KroneckerDelta[j, k] - x[[j]] x[[k]];
i[j_, k_, hi_, hf_] := M Integrate[Integrate[Integrate[f[j, k] r, {z, ((hf - hi)/R) r + hi, hf}], {r, 0, R}], {θ, 0, 2Pi}]
ITensor[hi_, hf_] := Table[FullSimplify[i[j, k, hi, hf]], {j, 1, 3}, {k, 1, 3}];
com = ITensor[-(3/4)h, (1/4)h];
origin = ITensor[0, h];
MatrixForm[com]
MatrixForm[origin]
RVec = {0, 0, -3/4};
steiner[j_, k_] := (origin[[j, k]] - M (Norm[RVec]^2 KroneckerDelta[j, k] - RVec[[j]] RVec[[k]]))
MatrixForm[Table[TrueQ[com[[j, k]] == steiner[j, k]], {j, 1, 3}, {k, 1, 3}]]
```

$$\text{i.} \quad \begin{pmatrix} \frac{1}{80} h M \pi R^2 (h^2 + 4R^2) & 0 & 0 \\ 0 & \frac{1}{80} h M \pi R^2 (h^2 + 4R^2) & 0 \\ 0 & 0 & \frac{1}{10} h M \pi R^4 \end{pmatrix}$$

$$\text{ii.} \quad \begin{pmatrix} \frac{1}{20} h M \pi R^2 (4h^2 + R^2) & 0 & 0 \\ 0 & \frac{1}{20} h M \pi R^2 (4h^2 + R^2) & 0 \\ 0 & 0 & \frac{1}{10} h M \pi R^4 \end{pmatrix}$$

(b) This array should be all True, but I think I've made a mistake in my definition of the Steiner equation.

$$\begin{pmatrix} \text{False} & \text{True} & \text{True} \\ \text{True} & \text{False} & \text{True} \\ \text{True} & \text{True} & \text{True} \end{pmatrix}$$

3.

4. The kinetic energy will have a term from the translation, which will be a rotation around the apex of the cone. Defining θ as the angle between the cone and the x-axis (the cone is resting on the x-y plane) we get the following translational kinetic energy:

$$V = \dot{\theta} \frac{3}{4} h \cos \alpha$$

$$T = \frac{9}{32} M \dot{\theta}^2 h^2 \cos^2 \alpha$$

For the rotational kinetic energy we need to find the angular velocity which we can define from $\dot{\theta}$:

$$\omega = \frac{4V}{3h \sin \alpha} = \frac{\dot{\theta} \cos \alpha}{\sin \alpha} = \dot{\theta} \cot \alpha$$

Using the moment of inertia tensor derived in the previous problem:

$$I = \begin{bmatrix} \frac{1}{20} h M \pi R^2 (4h^2 + R^2) & 0 & 0 \\ 0 & \frac{1}{20} h M \pi R^2 (4h^2 + R^2) & 0 \\ 0 & 0 & \frac{1}{10} h M \pi R^4 \end{bmatrix}$$

Substituting $h \sin \alpha = R$:

$$I = \begin{bmatrix} \frac{1}{20} h M \pi R^2 (4h^2 + h \sin^2 \alpha) & 0 & 0 \\ 0 & \frac{1}{20} h M \pi h \sin^2 \alpha (4h^2 + h \sin^2 \alpha) & 0 \\ 0 & 0 & \frac{1}{10} h M \pi h \sin^4 \alpha \end{bmatrix}$$

Finding the rotational kinetic energy with $T = \omega \cdot I \cdot \omega$:

$$\omega \cdot I \cdot \omega = \begin{bmatrix} \dot{\theta} \\ 0 \\ \dot{\theta} \cot \alpha \end{bmatrix} \begin{bmatrix} \frac{1}{20} h M \pi R^2 (4h^2 + h \sin^2 \alpha) & 0 & 0 \\ 0 & \frac{1}{20} h M \pi h \sin^2 \alpha (4h^2 + h \sin^2 \alpha) & 0 \\ 0 & 0 & \frac{1}{10} h M \pi h \sin^4 \alpha \end{bmatrix} \begin{bmatrix} \dot{\theta} \\ 0 \\ \dot{\theta} \cot \alpha \end{bmatrix}$$

$$\omega \cdot I \cdot \omega = \dot{\theta}^2 \frac{1}{20} h M \pi R^2 (4h^2 + h \sin^2 \alpha) + \dot{\theta}^2 \cot^2 \alpha \frac{1}{10} h M \pi h \sin^4 \alpha$$

The final kinetic energy is then:

$$T = \frac{9}{32} M \dot{\theta}^2 h^2 \cos^2 \alpha + \frac{1}{2} \left[\dot{\theta}^2 \frac{1}{20} h M \pi R^2 (4h^2 + h \sin^2 \alpha) + \dot{\theta}^2 \cot^2 \alpha \frac{1}{10} h M \pi h \sin^4 \alpha \right]$$