

Survival analysis in an experimental microbial aquatic community: Visualizations

Samuel Pawel

29 Juli 2021

```
## some libraries
library(ggplot2)
library(dplyr)
library(survival)

## import data set
dd <- readRDS("Dataset/extinction.week.Rdata")

## glimpse at data set
str(dd)



## 'data.frame': 3400 obs. of 5 variables:
## $ jar : int 1 2 3 4 5 6 7 8 9 10 ...
## $ temp : int 15 20 20 20 15 20 20 20 15 20 ...
## $ energy : num 0.275 0.275 0.275 0.825 0.825 0.825 0.275 0.275 0.275 0.825 ...
## $ species : chr "blepharisma" "blepharisma" "blepharisma" "blepharisma" ...
## $ week.persist: int 8 8 4 3 6 1 6 6 8 2 ...

summary(dd)

## jar temp energy species
## Min. : 1.00 Min. :15.0 Min. :0.275 Length:3400
## 1st Qu.: 50.75 1st Qu.:15.0 1st Qu.:0.275 Class :character
## Median :100.50 Median :17.5 Median :0.550 Mode :character
## Mean :100.50 Mean :17.5 Mean :0.550
## 3rd Qu.:150.25 3rd Qu.:20.0 3rd Qu.:0.825
## Max. :200.00 Max. :20.0 Max. :0.825
## week.persist
## Min. :0.000
## 1st Qu.:0.000
## Median :1.000
## Mean :3.302
## 3rd Qu.:8.000
## Max. :8.000

## visualize how long species persisted stratified by experimental conditions
dd$condition <- paste0("Temperature: ", dd$temp,
                      "°C x Energy: ", dd$energy)
ggplot(data = dd, aes(x = week.persist, fill = condition)) +
  geom_bar() +
  facet_wrap(~ species, ncol = 4) +
  scale_x_continuous(minor_breaks = NULL) +
```

```
labs(x = "Weeks persisted", fill = "") +
guides(fill = guide_legend(ncol = 2)) +
theme_bw() +
theme(legend.position = "top")
```



```


## create survival outcome that takes into account interval censoring
dd$timeSurv <- Surv(time = dd$week.persist,
                    time2 = ifelse(dd$week.persist == 8, Inf, dd$week.persist + 1),
                    type = "interval2")

## compute Kaplan Meier survival curves for each species and for all treatments
applyGrid <- expand.grid(species = unique(dd$species),
                        energy = unique(dd$energy),
                        temp = unique(dd$temp), stringsAsFactors = FALSE)
plotList <- lapply(X = seq(1, nrow(applyGrid)), FUN = function(i) {
  spDat <- filter(dd,
                  temp == applyGrid$temp[i],
                  species == applyGrid$species[i],
                  energy == applyGrid$energy[i])
  kmFit <- survfit(formula = timeSurv ~ 1, data = spDat)
  kmPredictions <- summary(kmFit, times = seq(0, 8, 1), extend = TRUE)
  data.frame(surv = kmPredictions$surv,
             lower = kmPredictions$lower,
             upper = kmPredictions$upper,
             time = kmPredictions$time,
             applyGrid[i,])
})

plotDF <- do.call("rbind", plotList)
plotDF$condition <- paste0("Temperature: ", plotDF$temp,
                          "°C x Energy: ", plotDF$energy)

## plot survival curves
ggplot(data = plotDF, aes(x = time , y = surv, color = condition)) +
  geom_step(alpha = 0.8) +
  facet_wrap(~ species, ncol = 4) +
  scale_x_continuous(minor_breaks = NULL) +
  labs(x = "Time (weeks)", y = "Survival probability", color = "") +
  guides(color = guide_legend(ncol = 2)) +
  theme_bw() +
  theme(legend.position = "top")

```



- There are species which went extinct already in the first week or which survived all 8 weeks. The estimated survivor curves look odd in both cases.
- Some species show either increased or decreased survival depending on the experimental conditions.
- The survival of some species seems unaffected by the experimental conditions.