

Contents

1	On Manifolds with Boundary	3
1.1	Regular values and transversality	3
1.2	Morse Theory	5
1.3	Integral curves and flows	7
2	Cartan Calculus	11
2.1	Lie derivative and Cartan's magic formula	11
2.2	Computation rules for time-dependent forms	15
2.3	Formulas for the Hodge star operator and codifferential	19
3	Riemannian Geometry	23
3.1	Product Manifolds	23
3.2	Induced connections	26
3.3	Miscellaneous	28
4	Chern Weil Theory	30
4.1	Construction of the Chern Weil Homomorphism	30
4.2	Proof of the Chern-Weil Theorem	33
4.3	Chern and Pontryagin Classes	35
5	Algebraic Topology	38
5.1	General Statements about Homology	38
5.2	Seifert-Van Kampen for Pushouts	39
5.3	CW-Complexes	39
5.4	Remarks about Urs' lecture on Algebraic Topology	41
5.5	Local Computation of the Degree	43
6	Point Set Topology	45
7	Fiber Bundles	46
7.1	Construction of Fiber Bundles	46
7.2	Changing the fiber	53
7.3	Principal Bundles	55
7.4	Vector Bundles	58
7.5	Universal Bundles	60
8	Construction of Diffeomorphisms	65
8.1	Existence of Smooth Isotopies	65
9	Some Linear Algebra	70
9.1	General Statements	70
9.2	Multilinear Algebra	71
9.3	Symplectic Vector Spaces	72
9.4	The Exterior Algebra	73
10	Functional Analysis	76
10.1	L^p and Sobolev Spaces	76
10.2	Complemented Subspaces	79
10.3	The Arzelà-Ascoli Theorem	80
11	Whitney Topologies	82
11.1	Approximation Results	82
11.2	An equivalent definition of the Whitney Topologies	96
11.3	C^k -convergence	98
12	Miscellaneous	103
12.1	Smoothness on arbitrary subsets	103
12.2	Integrating vector valued functions	104

12.3 Miscellaneous Statements about Manifolds	105
13 Solutions to selected exercises	106
14 Alternative Proofs	108
References	109

1 On Manifolds with Boundary

In this section we discuss how known theorems/lemmas fail for manifolds with non-empty boundary and how they can be generalized.

1.1 Regular values and transversality

It is known that for the regular value theorem, and more generally statements about transversality, to still hold for manifolds with boundary one needs to impose additional conditions on the boundary behaviour of a smooth function $f: M \rightarrow N$. Here is another lemma, that fits into the same category. We call $q \in N$ a **boundary value** of f if $q \in f(\partial M)$.

Lemma 1.1. *Let M be a manifold with boundary and $f: M \rightarrow \mathbb{R}$ a smooth function. Furthermore, let $a \in \mathbb{R}$ be a regular value of f , that is not a boundary value. Then $M_a := f^{-1}(-\infty, a]$ is a submanifold with boundary*

$$\partial M_a = f^{-1}(a) \cup (M_a \cap \partial M).$$

Proof sketch. For $p \in f^{-1}(a)$ we have $p \in \text{int}(M)$ by assumption. Since $\text{int}(M)$ is a boundaryless manifold (and an open subset of M) the local submersion theorem holds, i.e. f looks like the orthogonal projection $\mathbb{R}^n \rightarrow \mathbb{R}$ around p . Then the proof goes essentially the same as in the boundaryless case. $\square$

Counterexample 1.2. To see that Lemma 1.1 fails without the assumption that a is not a boundary value consider $M := S^1 \times [0, 1]$ and $f: M \rightarrow \mathbb{R}, (z, t) \mapsto t$. Then f is everywhere regular but $f^{-1}(-\infty, 0]$ does not satisfy the statements in Lemma 1.1.

Let M, N be manifolds with boundary and $f: M \rightarrow N$ smooth. We define the **boundary map** ∂f of f by

$$\partial f := f|_{\partial M}.$$

We now generalize two theorems from [GP74] about transversal maps by allowing the codomain to have boundary as well.

Theorem 1.3 (Transversality Theorem). *Let M, N be manifolds with boundary, $f: M \rightarrow N$ smooth and $X \subseteq N$ a submanifold without boundary. If*

$$f \pitchfork X \text{ and } \partial f \pitchfork X$$

then $X' := f^{-1}(X)$ is a submanifold with boundary

$$\partial X' = X' \cap \partial M$$

and

$$\text{codim}(X') = \text{codim}(X).$$

Proof. The case $\partial N = \emptyset$ is shown in [GP74] (p. 60). If $\partial N \neq \emptyset$ consider the inclusion

$$\iota: N \rightarrow D(N),$$

where $D(N)$ denotes the double of N and define $F := \iota \circ f$. Then $Y := \iota(X)$ is a submanifold of $D(N)$ and $X' = F^{-1}(Y)$. Furthermore, since ι is an embedding (and $\dim(N) = \dim(D(N))$) it follows that

$$F \pitchfork Y \text{ and } \partial F \pitchfork Y,$$

such that the statement follows from the boundaryless case. $\square$

We can also make a statement about the tangent space of X' in the above theorem. To this end let $f: M \rightarrow N$ and $X \subseteq N, X' \subseteq M$ be as above. Then we can also characterize transversality $f \pitchfork X$ as follows: Choose a metric on N and for $x \in X', y = f(x)$ define the **vertical differential**

$$D_x^V f: T_x M \rightarrow N_y X$$

as the composition

$$T_x M \xrightarrow{d_x f} T_y N = T_y X \oplus N_y X \xrightarrow{\text{proj.}} N_y X.$$

Then $f \pitchfork X$ if and only if $D_x^V f$ is surjective for all $x \in X'$. If this is the case we moreover have (at least if $\partial M = \emptyset$, otherwise we also require $\partial f \pitchfork X$)

$$\text{codim}(\ker D_x^V f) = \dim(\text{im}(D_x^V f)) = \text{codim}(X) = \text{codim}(X')$$

and thus (since $T_x X' \subseteq \ker D_x^V f$) we get

$$T_x X' = \ker D_x^V f.$$

This generalizes the formula for the tangent space of the preimage of a regular value. It should also be noted, that the Riemannian metric on N was only chosen, to define a complement of $T_y X$ in a consistent manner, such that the vertical differential $D_x^V f$ depends smoothly on x .

Theorem 1.4 (Parametric Transversality Theorem). *Let M, N be manifolds with boundary and $X \subseteq N$ a submanifold without boundary. Furthermore, let S be a manifold without boundary and $F: M \times S \rightarrow N$ a smooth map with*

$$F \pitchfork X \text{ and } \partial F \pitchfork X.$$

Then for almost every $s \in S$ the map $f_s: M \rightarrow N$ defined by $f_s(p) := F(p, s)$ satisfies

$$f_s \pitchfork X \text{ and } \partial f_s \pitchfork X.$$

Proof. By Theorem 1.3 we know that

$$X' := F^{-1}(X)$$

is a submanifold (with boundary) of $M \times S$. Consider the projection $\pi: M \times S \rightarrow S$. We can then show, that for a regular value $s \in S$ of the map $\pi|_{X'}$ the maps f_s and ∂f_s are transversal to X , which is just some linear algebra. The proof is exactly the same as in the case $\partial N = \emptyset$, see [GP74] (p. 68). Then the statement follows from Sard's Theorem. $\square$

In the above theorems the submanifold X which we pulled back by f was always assumed to be without boundary. Here is a version, which allows for X to have boundary as well.

Lemma 1.5. *Let M, N be smooth manifolds where M has no boundary and let $f: M \rightarrow N$ be a submersion. Given a submanifold $X \subseteq N$ with boundary the preimage $f^{-1}(X) \subseteq M$ is a submanifold with boundary*

$$\partial f^{-1}(X) = f^{-1}(\partial X)$$

and $\text{codim}(f^{-1}(X)) = \text{codim}(X)$.

Proof. First we may also assume $\partial N = \emptyset$, otherwise embed N into its double. Let $p \in f^{-1}(\partial X)$, $q = f(p) \in \partial X$ and choose a chart $\psi: V \rightarrow N$ around q , such that

$$\psi^{-1}(X) = V \cap \{(x_1, \dots, x_n) \mid x_1 \leq 0, x_2 = \dots = x_{s+1} = 0\},$$

where $n = \dim(N)$ and $s = \text{codim}(X)$. Since f is a submersion we can find a chart $\phi: U \rightarrow M$ around p , such that

$$\psi^{-1} \circ f \circ \phi: U \rightarrow V$$

is given by projection onto the first n coordinates, i.e. $(x_1, \dots, x_m) \mapsto (x_1, \dots, x_n)$ for $m = \dim(M) \geq n$. Thus in the coordinates of ϕ , $f^{-1}(X)$ is given by

$$\phi^{-1}(f^{-1}(X)) = U \cap \{(x_1, \dots, x_m) \mid x_1 \leq 0, x_2 = \dots = x_{s+1} = 0\},$$

such that ϕ is a submanifold chart for $f^{-1}(X)$ around p and $\partial(f^{-1}(X) \cap \phi(U)) = f^{-1}(X) \cap \phi(U)$. Similarly we construct submanifold charts around points $p \in f^{-1}(\text{int}(X))$ from which the statement follows. $\square$

Remark 1.6. The above Lemma becomes false if $\partial M \neq \emptyset$. Consider for example the inclusion $f: [0, 1] \hookrightarrow \mathbb{R}$ and $S := [1, 2] \subseteq \mathbb{R}$. Then

$$\partial f^{-1}(S) = \emptyset \neq f^{-1}(\partial S).$$

Or even worse, consider the 2-dimensional half space $\mathbb{H}^2 := \{(x, y) \in \mathbb{R}^2 \mid x \leq 0\}$ and the submersion $f: \mathbb{H}^2 \rightarrow \mathbb{R}$, $(x, y) \mapsto y$. Then $f^{-1}([0, 1])$ is not even a manifold anymore.

Moreover we cannot replace the submersion condition by only a transversality condition. Consider for example the map

$$f: \mathbb{R} \rightarrow \mathbb{R}^2, x \mapsto (0, x),$$

which is transversal to $S := [0, 1] \times \{0\} \subseteq \mathbb{R}^2$. Then again $\partial f^{-1}(S) = \emptyset \neq f^{-1}(\partial S)$.

1.2 Morse Theory

For a smooth function $f: M \rightarrow \mathbb{R}$ and a regular value $a \in \mathbb{R}$ that is not a boundary value we define the **sublevel set** of a to be

$$M_a := f^{-1}(-\infty, a].$$

Lemma 1.1 guarantees, that M_a is a (sub-)manifold with boundary.

Lemma 1.7. *Let M be a compact manifold with boundary and $f: M \rightarrow \mathbb{R}$ a Morse function. Furthermore let $a, b \in \mathbb{R}$ be regular values of f , such that $[a, b]$ does not contain any critical values nor any boundary values. Then*

$$M_a \cong M_b.$$

Proof. Let $K := f^{-1}[a, b]$ and let $U := \{p \in \text{int}(M) \mid p \text{ is a regular point of } f\}$. Then $U \subseteq M$ is open and $K \subseteq U$. Choose a cutoff function $\phi: M \rightarrow \mathbb{R}$ with

- $\phi|_K = 1$
- $\text{supp}(\phi) \subseteq U$
- $0 \leq \phi \leq 1$

Now choose a pseudo gradient¹ X of f and define the smooth vector field

$$Y := \phi \frac{-1}{X \cdot f} X,$$

which we define to be 0 outside of U . Since Y vanishes in a neighbourhood around ∂M (or more generally since Y is tangent to ∂M , see [Lee11] Theorem 9.34) we can consider the flow of Y :

$$\theta: M \times \mathbb{R} \rightarrow M.$$

Since M is compact (and Y tangent to ∂M) each integral curve is defined for all time. Let $\theta_t := \theta(-, t): M \rightarrow M$. We claim that

$$\theta_{b-a}(M_b) \subseteq M_a, \tag{1}$$

$$\theta_{a-b}(M_a) \subseteq M_b. \tag{2}$$

Let $p \in f^{-1}[a, b]$ and let $\gamma_p: \mathbb{R} \rightarrow M$ be the integral curve of Y starting at p . Consider

$$h: \mathbb{R} \rightarrow \mathbb{R}, \quad t \mapsto f(\gamma_p(t)).$$

Then for $t \in \mathbb{R}$ with $h(t) \in [a, b]$ we have $\phi(\gamma_p(t)) = 1$ and thus

$$h'(t) = \dot{\gamma}_p(t) \cdot f = Y_{\gamma_p(t)} \cdot f = -1.$$

So h is a smooth function with $c := h(0) = f(p) \in [a, b]$ and $h'(t) = -1$ as long as $h(t) \in [a, b]$. We claim that $h(c - a) = a$. To see this let

$$s := \sup\{t \in [0, c - a] \mid h([0, t]) \subseteq [a, b]\}.$$

Assume $s < c - a$. For continuity reasons we get that $h([0, s]) \subseteq [a, b]$ and thus $h'(t) = -1$ for $t \in [0, s]$. Then $h(t) = c - t$ for $t \in [0, s]$. But then $h(s) \in (a, b]$ and h is strictly decreasing around $t = s$ such that there exists $\varepsilon > 0$ with $h([0, s + \varepsilon]) \subseteq [a, b]$, contradicting the definition of s . So $s = c - a$ and we get that $h(t) = c - t$ for $t \in [0, c - a]$. Thus $h(c - a) = a$.

As $h'(t) \leq 0$ for all $t \in \mathbb{R}$ we conclude

$$f(\theta_{b-a}(p)) = h(b - a) \leq h(c - a) = a,$$

thus $\theta_{b-a}(p) \in M_a$. For $p \in M_b \setminus f^{-1}[a, b] = \text{int} M_a$ we also have

$$\theta_{b-a}(p) \in M_a$$

¹Here we agree to the convention that $X \cdot f < 0$ outside of critical points.

as f decreases along gradient flow lines of Y . This shows (1). We argue similarly for (2). Let $p \in M_a$ and let $\gamma_p: \mathbb{R} \rightarrow M$ be the integral curve of Y starting at p . Consider

$$h: \mathbb{R} \rightarrow \mathbb{R}, t \mapsto f(\gamma_p(-t)).$$

Then $h'(t) = \phi(\gamma_p(-t)) \leq 1$. Assume that $h(b-a) = f(\theta_{a-b}(p)) > b$. Then by the mean value theorem there exists some $\xi \in [0, b-a]$ with

$$h'(\xi) = \frac{h(b-a) - h(0)}{b-a} \geq \frac{h(b-a) - a}{b-a} > 1,$$

which is a contradiction. So $f(\theta_{a-b}(p)) \leq b$, showing (2). Thus θ_{b-a} restricts to a diffeomorphism $M_b \xrightarrow{\cong} M_a$ with inverse θ_{a-b} . $\square$

Counterexample 1.8. Example 1.2 also shows, that we need to impose conditions on the boundary values in Lemma 1.7. To see that we also need that condition for Lemma 1.10 we may similarly define f as the height function on the disjoint union of two cylinders which are slightly shifted in height. Similar examples show, that Theorem 1.11 fails without the assumption on boundary values.

Remark 1.9. We can weaken the assumptions of Lemma 1.7. First of all f does not need to be a Morse function. We can then still construct a vector field X , such that $X \cdot f < 0$ outside of critical points. Secondly, we may replace the assumption that M is compact by instead requiring $f^{-1}[a, b]$ to be compact, which is for example the case if f is proper. We then only have to modify the proof in the definition of U . Instead we define U to be a *compact* neighbourhood of $K := f^{-1}[a, b]$, i.e. $K \subseteq \text{int}U$, which does not contain any critical points nor boundary points. For example we may define U as a union of finitely many (here we need compactness of $f^{-1}[a, b]$) compact sets K_i not containing neither critical points nor boundary points such that $\text{int}K_i$ still cover K .

Similarly we may proof the following, for which we can also weaken the assumptions as described in the remark above.

Lemma 1.10. *Let M be a compact manifold with boundary and $f: M \rightarrow \mathbb{R}$ a Morse function. Furthermore let $a, b \in \mathbb{R}$ be regular values of f , such that $[a, b]$ does not contain any critical values nor any boundary values. Then M_a is a (not necessarily smooth) deformation retract of M_b .*

Proof. We consider the flow θ as in Lemma 1.7 and define a retraction

$$r: M_b \rightarrow M_a, p \mapsto \begin{cases} \theta(f(p) - a, p), & f(p) \geq a \\ p, & f(p) \leq a \end{cases}$$

As we have shown above $\theta(f(p) - a, p) \in f^{-1}(a)$. It follows that r is continuous and well-defined. To see that r is a deformation retract consider

$$H: M_b \times [0, 1] \rightarrow M_b, (p, t) \mapsto \begin{cases} \theta(t(f(p) - a), p), & f(p) \geq a \\ p, & f(p) \leq a \end{cases}$$

Then $H(-, 0) = \text{id}_{M_b}$ and $H(-, 1) = r$. $\square$

We would now also like to generalize the result, that the topology of M changes by attaching a λ -cell when crossing a critical point c_0 of index λ to compact manifolds with boundary. This statement can now be shown using the two lemmas above by requiring $f^{-1}[a, b]$ to only contain the critical point c_0 and no boundary points. The proof now works exactly the same as for the boundaryless case, see [Mil63]. Let us state the theorem for manifolds with boundary:

Theorem 1.11. *Let M be a compact manifold with boundary and $f: M \rightarrow \mathbb{R}$ a Morse function. Let $c_0 \in M$ be a critical point of f of index λ and $c := f(c_0)$ as well as $\varepsilon > 0$, such that $f^{-1}[c - \varepsilon, c + \varepsilon]$ contains no boundary points and no critical points besides c . Then for ε sufficiently small $M_{c+\varepsilon}$ has the homotopy type of $M_{c-\varepsilon}$ with a λ -cell attached. More precisely, there exists an attaching map*

$$\phi: S^{\lambda-1} = \partial D^\lambda \rightarrow \partial M_{c-\varepsilon}$$

and a homotopy equivalence

$$r: M_{c+\varepsilon} \rightarrow M_{c-\varepsilon} \cup_\phi D^\lambda$$

with homotopy inverse

$$\iota: M_{c-\varepsilon} \cup_\phi D^\lambda \rightarrow M_{c+\varepsilon},$$

such that $r \circ \iota = id_{M_{c-\varepsilon} \cup_\phi D^\lambda}$.

Proof Sketch. Following [Mil63] we first construct $F: M \rightarrow \mathbb{R}$, such that

$$F^{-1}(-\infty, c + \varepsilon] = f^{-1}(-\infty, c + \varepsilon] = M_{c+\varepsilon}.$$

Furthermore $F^{-1}[c - \varepsilon, c + \varepsilon]$ does not contain any critical points or boundary points, such that $F^{-1}(-\infty, c - \varepsilon]$ is a deformation retract of $F^{-1}(-\infty, c + \varepsilon]$ by Lemma 1.10. We then show that $M_{c-\varepsilon} \cup_\phi D^\lambda$ is a deformation retract of $F^{-1}(-\infty, c - \varepsilon]$ (or more precisely the image of $M_{c-\varepsilon} \cup_\phi D^\lambda$ under an embedding is a deformation retract). This then shows that $M_{c-\varepsilon} \cup_\phi D^\lambda$ is a deformation retract of $M_{c+\varepsilon}$. $\square$

However this does not yield a cell decomposition of M as we do not know how the topology changes when crossing boundary points. Instead we can use the fact that any compact manifold without boundary admits a cell-decomposition directly, to proof the case when M has non-empty boundary.

Theorem 1.12 (Cell-decompositions of Manifolds). *Let M be compact manifold with or without boundary. Then M has the homotopy type of a CW-complex.*

Proof. The case when M has no boundary is shown in [Mil63]. If $\partial M \neq \emptyset$ we can argue as follows: consider the double $D(M)$ of M . Then $D(M)$ is a compact manifold without boundary. Now we construct a Morse function on $D(M)$. Choose a Morse function $f: M \rightarrow \mathbb{R}$ with $f \leq 0$ and $f(p) = 0$ if and only if $p \in \partial M$, such that f has no critical points on the boundary. Such Morse functions always exist, see [Vin24]. We can then glue f and $-f$ together to get a Morse function F on $D(M)$, again see [Vin24]. Then $M \cong F^{-1}(-\infty, 0]$ and 0 is a regular value of F , allowing us to refer back to the boundaryless case, which tells us that every sublevel set M_a for a regular value $a \in \mathbb{R}$ has the homotopy type of a CW-complex. $\square$

Remark 1.13. We may also formulate Theorem 1.11 and 1.12 for handle bodies instead of CW-complexes. The proof is essentially the same but instead of constructing a deformation retract $F^{-1}(-\infty, c - \varepsilon] \rightarrow M_{c-\varepsilon} \cup_\phi D^\lambda$ one constructs a homeomorphism $F^{-1}(-\infty, c - \varepsilon] \rightarrow M_{c-\varepsilon} \cup_\phi D^\lambda \times D^{n-\lambda}$.

1.3 Integral curves and flows

For manifolds with boundary we essentially define integral curves in the same way as for boundaryless manifolds, but allow the domain to be any non-trivial interval (i.e. not a single point) containing 0. In particular this interval may only be half-open or closed. In general, the existence of integral curves on manifolds with boundary fails as illustrated by the following example:

Counterexample 1.14. Consider the lower halfspace $M := \mathbb{R}_{y \leq 0}^2$ and the vector field $X := \frac{\partial}{\partial x} + 2x \frac{\partial}{\partial y}$. Then for $p = (0, 0)$ the integral curve is only defined for $t = 0$, since on $\mathbb{R}^2$ it would be given by $\gamma(t) = (t, t^2)$ but $t^2 > 0$ for $t \neq 0$.




Figure 1: The vector field X . The integral curves are parabolas.

We do however still get a uniqueness result *if* the integral curve exists. Furthermore, if the vector field is sufficiently well-behaved we can guarantee the existence.

Theorem 1.15 (Uniqueness and existence of Integral curves). *Let M be a manifold with boundary, X a smooth vector field on M and $p \in M$. If there exists an integral curve starting at p , then there exists a unique maximal integral curve $\gamma: I \rightarrow M$ starting at p , where one of the four following cases holds for $a \leq 0 \leq b$, $a < b$:*

- (i) $I = (a, b)$
- (ii) $I = [a, b)$ and $\gamma(a) \in \partial M$
- (iii) $I = (a, b]$ and $\gamma(b) \in \partial M$
- (iv) $I = [a, b]$ and $\gamma(a), \gamma(b) \in \partial M$

Furthermore, if X is either transversal to the boundary or everywhere tangent, the unique maximal integral curve starting at p always exists.

Proof. Let's begin by showing uniqueness. The proof is essentially the same as in the boundaryless case (see for example [Lee11]): We show that two integral curves $\gamma: I \rightarrow M$ and $\gamma': I' \rightarrow M$ starting at p agree on their common domain. This then allows us to define the maximal integral curve on some interval I and every interval containing 0 is of the form (i) – (iv) ². The only difference is that I and I' may not be open intervals. So we have to show that uniqueness still applies. We claim: Let $U \subseteq \mathbb{R}^n$ be open, $F: U \rightarrow \mathbb{R}^n$ a smooth vector field, I an interval of the form (i) – (iv) and $\gamma, \gamma': I \rightarrow U$ integral curves both starting at $p \in U$. Then $\gamma = \gamma'$. If 0 is an interior point of I we can apply the usual uniqueness theorem for γ, γ' restricted to $\text{int}(I)$. By continuity they must then also agree at the endpoint. So it remains to show uniqueness if 0 is a boundary point of I (say for example $I = [0, b)$). But by the existence theorem for integral curves we can find an integral curve $\eta: (-\varepsilon, \varepsilon) \rightarrow U$ starting at p and define

$$\beta: (-\varepsilon, b) \rightarrow U, t \mapsto \begin{cases} \eta(t), t \leq 0 \\ \gamma(t), t \geq 0 \end{cases}$$

Since $\eta(0) = \gamma(0)$ and $\dot{\eta}(0) = F(p) = \dot{\gamma}(0)$ we get that β is at least C^1 . Similarly we define

$$\beta': (-\varepsilon, b) \rightarrow U, t \mapsto \begin{cases} \eta(t), t \leq 0 \\ \gamma'(t), t \geq 0 \end{cases}$$

By the uniqueness theorem we have

$$\beta = \beta'$$

and our claim follows. The case $I = (a, 0]$ goes analogous and for $I = [0, b]$ or $I = [a, 0]$ it is either trivial (if $a = b = 0$) or we consider $I \setminus \{b\}$ resp. $I \setminus \{a\}$ and apply the above argument. Equality at $t = a, t = b$ respectively then again follows by continuity.

To see that $\gamma(t) \in \partial M$ for $t \in \{a, b\}$ a boundary point of I we note that for $p = \gamma(t) \in \text{int}(M)$ we can define the integral curve starting at p on some open interval $(-\varepsilon, \varepsilon)$ by the existence theorem for integral curves (Picard-Lindelöf). But then t is not a boundary point of I since I is maximal.

Next we show the existence if X is transversal to ∂M . For $p \in \text{int}(M)$ this follows directly from the Picard-Lindelöf Theorem as in the boundaryless case. If $p \in \partial M$, we may argue similarly: Consider a chart $h: \mathbb{R}_{x_n \geq 0}^n \supseteq U \rightarrow M$ around $p = h(0)$ and let Y be the vector field X in this chart, i.e. $D_q h(Y_q) = X_{h(q)}$. Then by definition of smoothness we can extend Y to a smooth vector field Y' on some open neighbourhood $U' \subseteq \mathbb{R}^n$ around $0 = h^{-1}(p)$. Applying Picard-Lindelöf we get the existence of an integral curve $\gamma: (-\varepsilon, \varepsilon) \rightarrow U'$ of Y' starting at 0. Write

$$Y' = \sum_i F_i \frac{\partial}{\partial x_i}$$

and

$$\gamma = (\gamma_1, \dots, \gamma_n).$$

Assume that X is inwards pointing along the boundary. After restricting to a smaller domain we may assume $F_n \geq 0$. Then

$$\gamma'_n(t) = F_n(\gamma(t)) \geq 0,$$

such that γ_n is increasing. In particular $\gamma_n(t) \geq 0$ for $t \in [0, \varepsilon)$, hence $\gamma(t) \in U$ for $t \in [0, \varepsilon)$. Then

$$\eta := h \circ \gamma: [0, \varepsilon) \rightarrow M$$

is the desired integral curve.

For the case that X is everywhere tangent to ∂M see [Lee11] (Theorem 9.34). □

On compact manifolds we can say more about the maximal Interval I :

Lemma 1.16 (Existence of Integral curves on compact manifolds). *If M is compact in Lemma 1.15 then the maximal interval I has one of the following forms:*

- (i) $I = (-\infty, \infty)$
- (ii) $I = [a, \infty)$ and $\gamma(a) \in \partial M$
- (iii) $I = (-\infty, b]$ and $\gamma(b) \in \partial M$
- (iv) $I = [a, b]$ and $\gamma(a), \gamma(b) \in \partial M$

Proof. Let $\gamma: I \rightarrow M$ be the unique maximal integral curve guaranteed by Lemma 1.15. We will show that any open boundary point of I must be $\pm\infty$. For simplicity let us only consider $I = [a, b)$. The other cases are completely analogous. By compactness we can find a sequence $0 < t_n \nearrow b$ such that $\gamma(t_n)$ converges to a point $q \in M$.

Case 1: $q \in \text{int}(M)$. By Picard-Lindelöf we can find an open neighbourhood $U \subseteq M$ around q and $\varepsilon > 0$, such that for every point $x \in U$ the integral curve starting at x is defined for $t \in (-\varepsilon, \varepsilon)$. For some fixed m sufficiently large we have $q_0 := \gamma(t_m) \in U$. Assume that $b - t_m < \frac{\varepsilon}{2}$. But then we can extend γ to $[a, b + \frac{\varepsilon}{2})$ as follows. Consider

$$\gamma': [a - t_m, b - t_m) \rightarrow M, \quad t \mapsto \gamma(t + t_m).$$

Then γ' is a integral curve starting at q_0 . Let $\alpha: (-\varepsilon, \varepsilon) \rightarrow M$ be another integral curve starting at q_0 . Then α and γ' agree on their common domain by the arguments in Lemma 1.15 and thus we can put them together to define an integral curve

$$\beta: J \rightarrow M$$

for $J := [a - t_m, b - t_m) \cup (-\varepsilon, \varepsilon) \supseteq [a - t_m, \varepsilon)$. But then we can define

$$\gamma'': [a, \varepsilon + t_m) \rightarrow M, \quad t \mapsto \beta(t - t_m),$$

which is an integral curve starting at $\gamma(0) = p$. However

$$b + \frac{\varepsilon}{2} < t_m + \varepsilon$$

and so $[a, b + \frac{\varepsilon}{2}) \subseteq [a, \varepsilon + t_m)$, contradicting maximality of γ .

Case 2: $q \in \partial M$. We argue similarly as above. In a local chart around q we can again apply Picard-Lindelöf (smoothness of X guarantees by definition the existence of a smooth extension of X to an open neighbourhood in $\mathbb{R}^n$). We then again take $q_0 := \gamma(t_m)$ sufficiently close to q , choose an integral curve α starting at q , which we can use to define an integral curve α' starting at q_0 defined on some open interval $(-\varepsilon, \varepsilon)$ by translation in time as above. Also we define γ' as above. The same argument as above then shows that we can extend γ' (in the local chart) to an integral curve β on $J := [a - t_m, b - t_m) \cup (-\varepsilon, \varepsilon) \supseteq [a - t_m, b - t_m]$. In particular we have $\beta([a - t_m, b - t_m]) \subseteq M$. Translating in time again yields an integral curve $\gamma'': [a, b] \rightarrow M$ contradicting maximality. □

As an application we can show:

²If we put together integral curves defined on open intervals that agree on their common domain, it is clear that this results in a smooth curve, since around each t this curve is given by one of the smooth curves which we put together. If we put together integral curves γ_1, γ_2 defined on something like $I_1 = (a, b]$ and $I_2 = [b, c)$ we have to be more careful around the boundary point b . But the same argument as we applied to β in the proof of Lemma 1.15 shows that putting γ_1 and γ_2 together results in a C^1 -curve. We can then apply the usual uniqueness theorem to see that this curve is already given by a C^∞ -curve.

Lemma 1.17. *Let I be some proper interval containing 0 and $\gamma: I \rightarrow M$ any smooth regular curve, i.e. $\dot{\gamma}(t) \neq 0$ for all $t \in I$. Then there exists a unique unit speed reparametrization of γ , that is an orientation preserving diffeomorphism $\phi: J \rightarrow I$ with $\phi(0) = 0$, such that $\gamma \circ \phi$ is a unit speed curve.*

Proof. We have

$$(\gamma \circ \phi)'(t) = \phi'(t)\dot{\gamma}(\phi(t)),$$

so we need to find ϕ , such that (note that we also need $\phi' > 0$ to preserve orientation)

$$\phi'(t) = \frac{1}{|\dot{\gamma}(\phi(t))|}$$

This is a flow equation for the smooth vector field

$$X(p) := \frac{1}{|\dot{\gamma}(p)|}$$

on I . Since X is transversal to the boundary of I (if I even has boundary), Theorem 1.15 yields the existence of a unique maximal integral curve $\phi: J \rightarrow I$ of X with $\phi(0) = 0$. We now only need to show that ϕ is surjective, since $\phi' > 0$ already shows that it is an embedding.

To this end, consider any compact subinterval $I' := [a, b] \subseteq I$ ($a < b$) containing 0 and let $\psi: J' \rightarrow I'$ be the unique maximal integral curve of $X|_{I'}$ starting at 0. By Lemma 1.16 ψ has to be defined on a compact interval $J' = [c, d]$, since $X|_{I'}$ is bounded below by a positive constant, so if J' would be infinite in the positive/negative direction, ψ would also tend to positive/negative infinity, but by definition it is contained in the compact interval I' . Moreover Lemma 1.16 then yields

$$\psi(c) = a, \quad \psi(d) = b.$$

Since ϕ is the unique maximal integral curve of X we get that $J' \subseteq J$ and $\phi|_{J'} = \psi$. In particular $I' \subseteq \text{im}(\phi)$. This completes the proof. $\square$

Remark 1.18 (A note on flows for manifolds with boundary). In general the concept of a flow is not well behaved if M has boundary, as integral curves may not even exist (see Counterexample 1.14). In [Lee11] the author shows, that if X is tangent to ∂M we still get a well-behaved flow, i.e. it is defined on some open subset $\mathcal{O} \subseteq M \times \mathbb{R}$. I think that if we instead assume, that X is transversal to ∂M , we still get a "good" flow domain in the sense that we can define the flow on some subset $\mathcal{D} \subseteq M \times \mathbb{R}$, which is a codimension 0 submanifold with boundary and $I_p := \{t \in \mathbb{R} \mid (p, t) \in \mathcal{D}\}$ is an interval around 0 for each $p \in M$.

The statement in the above remark should follow from the following lemma, which allows us to define submanifold charts for $\mathcal{D}$.

Lemma 1.19. *Let M be a manifold with boundary and X a vector field of M , which is inward pointing along ∂M . Let*

$$M_{\partial} := \{p \in M \mid \text{im} \gamma_p \cap \partial M \neq \emptyset\}$$

be all points in M whose trajectory contains boundary points (this is at most one for each $p \in M$). For $p \in M_{\partial}$ the integral curve γ_p starting at p is defined on some interval $[-\tau(p), a)$ for some $\tau(p) \geq 0$ and $a > -\tau(p)$. Then the following statements hold true

- (i) M_{∂} is an open subset of M
- (ii) $\tau: M_{\partial} \rightarrow \mathbb{R}_{\geq 0}, p \mapsto \tau(p)$ is smooth

Proof sketch. The idea is to consider the maximal flow

$$\theta: \mathcal{D} \rightarrow M, (p, t) \mapsto \gamma_p(t)$$

on some open subset $\mathcal{D} \subseteq \partial M \times [0, \infty)$ and show that this is an embedding. Then

$$M_{\partial} = \text{im} \theta$$

is open and

$$\tau = \pi \circ \theta^{-1}: M_{\partial} \rightarrow [0, \infty)$$

where $\pi: \partial M \times [0, \infty) \rightarrow [0, \infty)$ is the canonical projection map. $\square$

Lemma 1.20. *Injective Flowouts are already embeddings*

2 Cartan Calculus

2.1 Lie derivative and Cartan's magic formula

In this section we discuss two differential operators on the graded algebra $\Omega^\bullet(M)$ of differential forms on a manifold M , namely the Lie derivative $\mathcal{L}_X$ and the interior product ι_X , where X is a smooth vector field on M . The main goal of this section is to show well-definedness of $\mathcal{L}_X$ and proof the following theorem, describing the relationships between $\mathcal{L}_X$, ι_X and the exterior derivative d .

Theorem 2.1. *Let M be a manifold and $X, Y \in \mathcal{X}(M)$. Then the maps*

$$\begin{aligned} d: \Omega^r(M) &\rightarrow \Omega^{r+1}(M), \\ \mathcal{L}_X: \Omega^r(M) &\rightarrow \Omega^r(M), \\ \iota_X: \Omega^r(M) &\rightarrow \Omega^{r-1}(M) \end{aligned}$$

satisfy the following Leibniz rules for $\alpha, \beta \in \Omega^\bullet(M)$:

$$\begin{aligned} d(\alpha \wedge \beta) &= d\alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge d\beta, \\ \mathcal{L}_X(\alpha \wedge \beta) &= \mathcal{L}_X\alpha \wedge \beta + \alpha \wedge \mathcal{L}_X\beta, \\ \iota_X(\alpha \wedge \beta) &= \iota_X\alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge \iota_X\beta, \end{aligned}$$

as well as the following identities:

- (i) $d^2 = 0$,
- (ii) $d\mathcal{L}_X = \mathcal{L}_Xd$,
- (iii) $\mathcal{L}_X = d\iota_X + \iota_Xd$,
- (iv) $\mathcal{L}_X\mathcal{L}_Y - \mathcal{L}_Y\mathcal{L}_X = \mathcal{L}_{[X,Y]}$,
- (v) $\mathcal{L}_X\iota_Y - \iota_Y\mathcal{L}_X = \iota_{[X,Y]}$,
- (vi) $\iota_X\iota_Y + \iota_Y\iota_X = 0$.

Firstly let us restate the definition of the Lie derivative. For a smooth manifold M without boundary, $\alpha \in \Omega^r(M)$ and $X \in \mathcal{X}(M)$ we define

$$(\mathcal{L}_X\alpha)_p := \left. \frac{d}{dt} \right|_{t=0} (\phi_t^*\alpha)_p,$$

where $\phi_t = \phi(t, -)$ denotes the flow of X .

Lemma 2.2. *$(\mathcal{L}_X\alpha)_p$ exists for all $p \in M$ and defines a smooth r -form on M .*

Before showing the lemma, let us introduce the notion of a smooth family of r -forms.

Definition 2.3 (Smooth family of r -forms). Let β_t be a family of r -forms on M indexed over some open Interval I . We call β_t smooth if in some (then all) local coordinates $x^1, \dots, x^n$ we can write

$$(\beta_t)_p = \sum \beta^{i_1, \dots, i_r}(t, p) d_px_{i_1} \wedge \dots \wedge d_px_{i_r}$$

for smooth functions $\beta^{i_1, \dots, i_r}$. If I is closed we call β_t smooth if it can be extended to a smooth family β'_t indexed over some open interval $I' \supseteq I$.

Then $t \mapsto (\beta_t)_p$ is a smooth curve in the vector space $\Lambda^r(T_p^*M)$, such that

$$\left(\left. \frac{d}{dt} \right|_{t=t_0} \beta_t \right)_p := \left. \frac{d}{dt} \right|_{t=t_0} (\beta_t)_p$$

is a well-defined r -form. Furthermore, in local coordinates we have

$$\left. \frac{d}{dt} \right|_{t=t_0} \beta_t = \sum \frac{\partial \beta^{i_1, \dots, i_r}}{\partial t}(t_0, -) dx_{i_1} \wedge \dots \wedge dx_{i_r},$$

such that this r -form is also smooth and moreover depends smoothly on t_0 . So

$$\dot{\beta}_t := \frac{d}{dt}\beta_t$$

defines a smooth family of r -forms by

$$\left(\frac{d}{dt}\beta_t\right)_{t_0} := \frac{d}{dt}\Big|_{t=t_0} \beta_t.$$

Note that if $\partial M = \emptyset$ we can also directly allow for closed intervals I in the definition above. However we may then always extend the family β_t to an open interval (in fact to all of $\mathbb{R}$), as β_t can equivalently be considered a smooth r -form β on $M \times I \subseteq M \times \mathbb{R}$. Under this correspondence we have the following interesting relationship (without proof):

$$\dot{\beta}_t = \mathcal{L}_{\frac{\partial}{\partial t}} \beta.$$

Proof of Lemma 2.2. Let $x_1, \dots, x_n$ be local coordinates on some open neighbourhood $U \subseteq M$ of p . Choose a smaller neighbourhood $U' \subseteq U$ of p and $\varepsilon > 0$, such that the flow

$$\phi: (-\varepsilon, \varepsilon) \times U' \rightarrow U$$

of X is defined. Write

$$\alpha = \sum \alpha^{i_1, \dots, i_r} dx_{i_1} \wedge \dots \wedge dx_{i_r}.$$

Then:

$$(\phi_t^* \alpha)_p = \sum \alpha^{i_1, \dots, i_r}(\phi_t(p)) d_p(x_{i_1} \circ \phi_t) \wedge \dots \wedge d_p(x_{i_r} \circ \phi_t)$$

Expanding the differentials

$$d_p(x_{i_k} \circ \phi_t) = \sum_{j=1}^n \frac{\partial(x_{i_k} \circ \phi_t)}{\partial x_j}(p) d_p x_j$$

we see that $\beta_t := \phi_t^* \alpha$ is a smooth family of r -forms on U' indexed over $t \in (-\varepsilon, \varepsilon)$. Thus

$$(\mathcal{L}_X \alpha)_p = \left(\frac{d}{dt}\Big|_{t=0} \beta_t\right)_p$$

exists and depends smoothly on p . □

Proposition 2.4 (Basic properties of the Lie derivative). *Let $\alpha \in \Omega^r(M)$, $\beta \in \Omega^s(M)$, $\lambda \in \mathbb{R}$ and $X \in \mathcal{X}(M)$. Then the Lie derivative satisfies the following properties:*

- (i) *Locality:* $(\mathcal{L}_X \alpha)|_U = \mathcal{L}_{X|_U} \alpha|_U$ for $U \subseteq M$ open
- (ii) *Linearity in α :* $\mathcal{L}_X(\lambda\alpha + \beta) = \lambda\mathcal{L}_X \alpha + \mathcal{L}_X \beta$
- (iii) *Leibniz rule:* $\mathcal{L}_X(\alpha \wedge \beta) = \mathcal{L}_X \alpha \wedge \beta + \alpha \wedge \mathcal{L}_X \beta$
- (iv) *Compatibility with d :* $\mathcal{L}_X(d\alpha) = d(\mathcal{L}_X \alpha)$

Proof. (i) follows from the fact that locally both sides are given by

$$\frac{d}{dt}\Big|_{t=0} \beta_t$$

for the smooth family β_t defined in the proof of Lemma 2.2. Moreover, (ii) follows immediately from the definition. To see (iii) write

$$\mathcal{L}_X(\alpha \wedge \beta)_p = \frac{d}{dt}\Big|_{t=0} (\phi_t^* \alpha)_p \wedge (\phi_t^* \beta)_p.$$

We claim: For a vector space V and smooth curves $\alpha_t \in \Lambda^r(V^*)$, $\beta_t \in \Lambda^s(V^*)$ the following holds:

$$\left. \frac{d}{dt} \right|_{t=0} (\alpha_t \wedge \beta_t) = \left(\left. \frac{d}{dt} \right|_{t=0} \alpha_t \right) \wedge \beta_0 + \alpha_0 \wedge \left(\left. \frac{d}{dt} \right|_{t=0} \beta_t \right)$$

W.l.o.g. let

$$\alpha_t = a(t)\alpha', \quad \beta_t = b(t)\beta'$$

for some $\alpha' \in \Lambda^r(V^*)$, $\beta' \in \Lambda^s(V^*)$ and smooth functions a, b . Then the general case follows from linearity. We compute:

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=0} (\alpha_t \wedge \beta_t) &= \left. \frac{d}{dt} \right|_{t=0} a(t)b(t)\alpha' \wedge \beta' \\ &= \left(\left. \frac{da}{dt} \right|_{t=0} b(0) + a(0) \left. \frac{db}{dt} \right|_{t=0} \right) \alpha' \wedge \beta' \\ &= \left. \frac{da}{dt} \right|_{t=0} \alpha' \wedge b(0)\beta' + a(0)\alpha' \wedge \left. \frac{db}{dt} \right|_{t=0} \beta' \\ &= \left(\left. \frac{d}{dt} \right|_{t=0} \alpha_t \right) \wedge \beta_0 + \alpha_0 \wedge \left(\left. \frac{d}{dt} \right|_{t=0} \beta_t \right). \end{aligned}$$

This shows the claim and thus (iii). To see (iv) let β_t be again the smooth family of r -forms defined in the proof of Lemma 2.2 and write in local coordinates $x_1, \dots, x_n$:

$$\beta_t = \sum \beta^{i_1, \dots, i_r}(t, -) dx_{i_1} \wedge \dots \wedge dx_{i_r}.$$

By linearity we may assume that all functions $\beta^{i_1, \dots, i_r}$ except for one vanish, i.e.

$$\beta_t = f(t, -) dx_{j_1} \wedge \dots \wedge dx_{j_r}$$

for some smooth function f and $1 \leq j_1 < \dots < j_r \leq n$. Then

$$d\beta_t = \sum_{k=1}^n \frac{\partial f(t, -)}{\partial x_k} dx^k \wedge dx_{j_1} \wedge \dots \wedge dx_{j_r}.$$

Thus

$$\begin{aligned} \mathcal{L}_X(d\alpha) &= \left. \frac{d}{dt} \right|_{t=0} \phi_t^* d\alpha = \left. \frac{d}{dt} \right|_{t=0} d\beta_t \\ &= \sum_{k=1}^n \frac{\partial^2 f}{\partial t \partial x_k}(0, -) dx_k \wedge dx_{j_1} \wedge \dots \wedge dx_{j_r} \\ &= d \left(\left. \frac{\partial f}{\partial t} \right|_{t=0}(0, -) dx_{j_1} \wedge \dots \wedge dx_{j_r} \right) \\ &= d \left(\left. \frac{d}{dt} \right|_{t=0} \beta_t \right) = d(\mathcal{L}_X \alpha) \end{aligned}$$

□

For convenience, let us also restate the definition of ι_X

Definition 2.5. Let $X \in \mathcal{X}(M)$ and $\alpha \in \Omega^r(M)$. For $r > 0$ define

$$\iota_X \alpha := \alpha(X, -) \in \Omega^{r-1}(M)$$

For $r = 0$ let $\iota_X \alpha := 0$.

Clearly ι_X is linear and only depends on the local behaviour of X and α . Furthermore, it is easily seen that

$$\iota_X(\alpha \wedge \beta) = \iota_X \alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge \iota_X \beta.$$

Thus far the Lie derivative has only been defined on manifolds without boundary. To extend the definition of $\mathcal{L}_X \alpha$ to manifolds with boundary, first choose extensions X' and α' of X and α , respectively, to the double of M and then define

$$\mathcal{L}_X \alpha := (\mathcal{L}_{X'} \alpha')|_M$$

Of course a priori it is not clear that this definition does not depend on the choice of X' and α' . However, the following proposition, which is one of the formulas of Theorem 2.1, shows that it is in fact independent.

Proposition 2.6 (Cartan's magic formula). *Let $X \in \mathcal{X}(M)$. Then the following identity holds:*

$$\mathcal{L}_X = d\iota_X + \iota_X d$$

Proof. Let $\mathcal{D}_X := d\iota_X + \iota_X d$. From the discussion above it is easy to verify:

$$\mathcal{D}_X(\alpha \wedge \beta) = \mathcal{D}_X \alpha \wedge \beta + \alpha \wedge \mathcal{D}_X \beta.$$

Moreover, $\mathcal{D}_X$ clearly satisfies locality, linearity and commutes with d . Thus we can establish

$$\mathcal{D}_X = \mathcal{L}_X$$

by showing equality on $\Omega^0(M)$. So let $f \in \Omega^0(M)$, then

$$\mathcal{D}_X f = \iota_X df = X \cdot f$$

and

$$\mathcal{L}_X f = \left. \frac{d}{dt} \right|_{t=0} (f \circ \phi_t) = df(X) = X \cdot f$$

This shows the proposition. $\square$

Now let us proof the remaining formulas of Theorem 2.1, which are shown similarly.

Proof of Theorem 2.1. Equation (i) is only mentioned for the sake of completeness and will not be shown here. See for example [Ste23a] for a proof. Equations (ii) and (iii) have already been shown above.

To see (iv) note that again both sides of the equation are differential operators satisfying the same Leibniz rule and commute with d , such that it suffices to show equality on $\Omega^0(M)$. But this follows immediately from the definition of the Lie bracket.

We argue similarly for (v), but in this equation the differential operators on each side do not commute with d . However they still satisfy the same Leibniz rule, so we only need to show equality on $\Omega^0(M)$ as well as on the exact one-forms. Clearly both sides vanish on $\Omega^0(M)$ (by definition of the interior product) and for $\alpha = df \in \Omega^1(M)$ we compute:

$$\begin{aligned} (\mathcal{L}_X \iota_Y - \iota_Y \mathcal{L}_X)(df) &= \mathcal{L}_X(Y \cdot f) - (\iota_Y d\mathcal{L}_X)(f) \\ &= XYf - YXf \\ &= [X, Y] \cdot f = \iota_{[X, Y]}(df). \end{aligned}$$

Finally, (vi) is just a way to rewrite

$$\alpha(X, Y, -) = -\alpha(Y, X, -).$$

$\square$

As an application of Theorem 2.1 we can show the following identity:

Corollary 2.7. *For $\alpha \in \Omega^1(M)$ and $X, Y \in \mathcal{X}(M)$ the following identity holds:*

$$d\alpha(X, Y) = X \cdot \alpha(Y) - Y \cdot \alpha(X) - \alpha([X, Y])$$

Proof. By Theorem 2.1 (iii) we have:

$$(\iota_Y \mathcal{L}_X)(\alpha) = Y \cdot \alpha(X) + d\alpha(X, Y).$$

Moreover, applying (v) yields:

$$(\iota_Y \mathcal{L}_X)(\alpha) = X \cdot \alpha(Y) - \alpha([X, Y]).$$

The corollary follows. $\square$

Integrating time dependent forms

Let β_t be a smooth time dependent family of differential forms on a manifold M defined on some interval I . At the beginning of the section we described what it means to take the derivative of β_t with respect to time. Often one is also interested in the *integral* with respect to time, which is defined similarly, i.e. pointwise. For $a, b \in I$ and $p \in M$ we set

$$\left(\int_a^b \beta_t dt \right)_p := \int_a^b (\beta_t)_p dt,$$

where the right side is just the integral of a smooth curve $[a, b] \rightarrow \Lambda^r(T_p^*M)$ (see section 12.2 for a definition of such an integral). Furthermore, in local coordinates we have

$$\int_a^b \beta_t dt = \sum \int_a^b \beta^{i_1, \dots, i_r}(t, -) dt dx_{i_1} \wedge \dots \wedge dx_{i_r},$$

so $\int_a^b \beta_t dt$ is a smooth r -form on M .

2.2 Computation rules for time-dependent forms

The goal of this section is to establish a few coordinate free rules for computing expressions involving the derivative or integral of time dependent differential forms in practice. We begin by discussing some elementary common cases and explain how to deal with more complex ones later on.

The Lie derivative $\mathcal{L}_X \alpha$ was defined as

$$\left. \frac{d}{dt} \right|_{t=0} \phi_t^* \alpha,$$

where ϕ_t is the flow of X . We might also want to differentiate this expression at some other time $t = t_0$. The following proposition shows how this can be computed in terms of $\mathcal{L}_X \alpha$. In fact, we show a more general statement, where ϕ_t is the flow of a time dependent vector field.

Proposition 2.8. *Let M be a manifold without boundary, X_t a time dependent vector field on M and $\alpha \in \Omega^r(M)$. Denote by ϕ_t the flow of X_t and consider the open set $M_{t_0} = \{p \in M \mid \phi_t(p) \text{ exists for } t = t_0\}$. Then*

$$\left. \frac{d}{dt} \right|_{t=t_0} (\phi_t^* \alpha)_p = \phi_{t_0}^* (\mathcal{L}_{X_{t_0}} \alpha)_p$$

for all points $p \in M_{t_0}$.

Proof. We begin by introducing some notation. Say X_t is defined for time $t = t_1$. We then write $\phi_{t_1, t}$ for the solution of the following nonautonomous ODE

$$\frac{d}{dt} \phi_{t_1, t} = X_{t+t_1}(\phi_{t_1, t}),$$

i.e. the flow of the reparametrised time-dependent vector field X_{t+t_1} . Then we have $\phi_{0, t} = \phi_t$ and

$$\phi_{t+t_0} = \phi_{t_0, t} \circ \phi_{0, t_0}$$

wherever this equation makes sense. Then we can compute:

$$\begin{aligned} \frac{d}{dt} \Big|_{t=t_0} (\phi_t^* \alpha)_p &= \frac{d}{dt} \Big|_{t=0} (\phi_{t+t_0}^* \alpha)_p \\ &= \frac{d}{dt} \Big|_{t=0} (\phi_{0,t_0}^* \phi_{t_0,t}^* \alpha)_p \\ &= \phi_{0,t_0}^* \frac{d}{dt} \Big|_{t=0} (\phi_{t_0,t}^* \alpha)_p \end{aligned}$$

We claim that

$$\frac{d}{dt} \Big|_{t=0} (\phi_{t_0,t}^* \alpha)_p = (\mathcal{L}_{X_{t_0}} \alpha)_p$$

To see this, we can explicitly express both sides in local coordinates and then use (ii) from Lemma 2.9 and (i) from Lemma 2.10 to reduce this to the case, where α is a 0-form, which follows immediately. Essentially, this is again using the trick that two differential operators are equal if they satisfy the same Leibniz rule, commute with d and agree on 0-forms. $\square$

Lemma 2.9. *Let α_t, β_t be smooth time-dependent families of differential forms on a manifold M and $f: M \rightarrow N$ a smooth map. Then the following holds:*

- (i) $\frac{d}{dt} f^* \alpha_t = f^* \frac{d}{dt} \alpha_t$
- (ii) $\frac{d}{dt} (\alpha_t \wedge \beta_t) = \frac{d}{dt} \alpha_t \wedge \beta_t + \alpha_t \wedge \frac{d}{dt} \beta_t$

Proof. For (i) note that for a given point $p \in M$ the pullback along f is given by a linear map

$$f^*: \Lambda^r(T_{f(p)}^* N) \rightarrow \Lambda^r(T_p^* M),$$

so the statement follows directly from the chain rule. (ii) follows from the ordinary Leibniz rule for smooth functions and can be seen by computing the derivative in local coordinates. We have in fact already seen this computation in the proof of Proposition 2.4. $\square$

This next Lemma shows, that the time derivative of time-dependent differential forms commutes with the differential operators from Theorem 2.1.

Lemma 2.10. *Let α_t be a time-dependent family of differential forms and X a vector field on a manifold M . Then the following holds:*

- (i) $\frac{d}{dt} d\alpha_t = d \frac{d}{dt} \alpha_t$
- (ii) $\frac{d}{dt} \mathcal{L}_X \alpha_t = \mathcal{L}_X \frac{d}{dt} \alpha_t$
- (iii) $\frac{d}{dt} \iota_X \alpha_t = \iota_X \frac{d}{dt} \alpha_t$

Proof. (i) follows from Schwarz's Theorem by computing the expression in local coordinates. This computation has essentially already been carried out in the proof of Proposition 2.4. (ii) follows from (i) and (iii) using Cartan's magic formula. For (iii) note that for $p \in M$ the map

$$\iota_{X_p}: \Lambda^r(T_p^* M) \rightarrow \Lambda^{r-1}(T_p^* M)$$

is linear so the statement follows again from the chain rule. $\square$

Moreover, the integral with respect to time also commutes with the differential operators, as the following Lemma shows.

Lemma 2.11. *Let α_t be a time-dependent family of differential forms defined on some interval I and X a vector field on a manifold M . Given $a, b \in I$ the following holds:*

- (i) $d \int_a^b \alpha_t dt = \int_a^b d\alpha_t dt$
- (ii) $\mathcal{L}_X \int_a^b \alpha_t dt = \int_a^b \mathcal{L}_X \alpha_t dt$
- (iii) $\iota_X \int_a^b \alpha_t dt = \int_a^b \iota_X \alpha_t dt$

Proof. For (i) choose local coordinates $x_1, \dots, x_n$ and let $\alpha^{i_1, \dots, i_r}(t, x)$ be a component function of α_t with respect to these coordinates. Then

$$\frac{\partial}{\partial x_i} \int_a^b \alpha^{i_1, \dots, i_r}(t, x) dt = \int_a^b \frac{\partial}{\partial x_i} \alpha^{i_1, \dots, i_r}(t, x) dt$$

since $[a, b]$ is compact. The statement follows. Again (ii) follows from (i) and (iii) using Cartan's magic formula. (iii) follows from linearity of the integral (see Lemma 12.8) by again considering the linear map ι_{X_p} for $p \in M$. $\square$

Finally, we explain how to deal with time dependent vector fields in the differential operators.

Lemma 2.12. *Let X_t be a time-dependent vector field on a manifold M . Then the following holds:*

$$(i) \quad \frac{d}{dt} \mathcal{L}_{X_t} = \mathcal{L}_{\frac{d}{dt} X_t}$$

$$(ii) \quad \frac{d}{dt} \iota_{X_t} = \iota_{\frac{d}{dt} X_t}$$

Proof. (i) follows from (ii) as well as Lemma 2.10 (i) using Cartan's magic formula. For (ii) let $\alpha \in \Omega^r(M)$ and consider the linear map

$$A: T_p M \rightarrow \Lambda^{r-1}(T_p^* M), \quad v \mapsto \alpha_p(v, -)$$

for $p \in M$. Then

$$\left(\frac{d}{dt} \iota_{X_t} \alpha \right)_p = \frac{d}{dt} A(X_t(p))$$

and so the statement follows again from the chain rule. $\square$

Remark 2.13. Of course analogous statements hold for $\mathbb{R}^n$ -valued time parameters $t \in \mathbb{R}^n$.

Moser's trick

In many cases one has to differentiate a time-dependent family of differential forms, which is given by an expression where the time variable appears at multiple instances. To understand how a formula for this is obtained, we begin by discussing an important example, namely Moser's trick, which is used to prove the Darboux Theorem about symplectic/contact manifolds. Following that, we present a straightforward cooking recipe on how to compute the derivative of such an expression in general, illustrated by another example.

Let M be a manifold (without boundary). Given a smooth family α_t of differential forms on M indexed over $t \in \mathbb{R}$ we want to find a family of diffeomorphisms $\varphi_t: M \rightarrow M$ (i.e. an isotopy), such that $\varphi_0 = id_M$ and

$$\varphi_t^* \alpha_t = \alpha_0 \quad \text{for all } t \in [0, 1],$$

or equivalently:

$$\frac{d}{dt} \varphi_t^* \alpha_t = 0. \tag{3}$$

Directly constructing such an isotopy is difficult but we can make use of the following trick: The isotopy φ_t corresponds to the level preserving diffeomorphism

$$\Phi: M \times [0, 1] \rightarrow M \times [0, 1], (p, t) \mapsto (\varphi_t(p), t).$$

This diffeomorphism then defines a vector field $\Phi_* \frac{\partial}{\partial t}$ on $M \times [0, 1]$, whose integral curves uniquely determine φ_t . This suggests that in order to construct an isotopy φ_t we can construct a vector field Y on $M \times [0, 1]$, of the form

$$Y_{(p,t)} = ((X_t)_p, \frac{\partial}{\partial t})$$

for a time dependent vector field X_t on M (this trick is also explained in [Mil65] Thm. 5.6).

Then (3) can be reformulated as a condition on X_t , which is shown in the following Lemma.

Lemma 2.14. *In the situation above the following holds:*

$$\frac{d}{dt} \varphi_t^* \alpha_t = \varphi_t^* (L_{X_t} \alpha_t + \dot{\alpha}_t)$$

Proof. The idea is to "decouple" the time variables and consider the two parameter family

$$g: [0, 1] \times \mathbb{R} \rightarrow T_p^*M, (t, s) \mapsto (\varphi_t^* \alpha_s)_p$$

The partial derivatives are now much easier to compute. By Proposition 2.8 we have:

$$\left. \frac{\partial}{\partial t} \right|_{t=t_0} \varphi_t^* \alpha_s = \varphi_{t_0}^* (\mathcal{L}_{X_{t_0}} \alpha_s)$$

Moreover, by Lemma 2.9

$$\left. \frac{\partial}{\partial s} \right|_{s=t_0} \varphi_t^* \alpha_s = \varphi_t^* \left(\left. \frac{\partial}{\partial s} \right|_{s=t_0} \alpha_s \right) = \varphi_t^* \dot{\alpha}_{t_0}.$$

In order to compute the derivative of

$$f: [0, 1] \rightarrow T_p^*M, t \mapsto (\varphi_t^* \alpha_t)_p$$

consider the diagonal map

$$\Delta: [0, 1] \rightarrow [0, 1] \times \mathbb{R}, t \mapsto (t, t).$$

Then

$$f = g \circ \Delta,$$

hence

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=t_0} f(t) &= D_{t_0} f(1) = D_{(t_0, t_0)} g \circ D_{t_0} \Delta(1) \\ &= D_{(t_0, t_0)}(1, 1) = D_{(t_0, t_0)}(1, 0) + D_{(t_0, t_0)}(0, 1) \\ &= \frac{\partial g}{\partial t}(t_0, t_0) + \frac{\partial g}{\partial s}(t_0, t_0) \end{aligned}$$

The Lemma follows. □

So (3) can be reformulated to the condition:

$$0 = L_{X_t} \alpha_t + \dot{\alpha}_t = d\iota_{X_t} \alpha_t + \iota_{X_t} d\alpha_t + \dot{\alpha}_t,$$

where we used Theorem 2.1 (iii) in the second equation. Hence it suffices to construct a time dependent vector field X_t satisfying the above condition and whose integral curves exist for time $t = 1$.

A cooking recipe

The proof of Lemma 2.14 now gives the announced cooking recipe for computing derivatives of smooth families of differential forms indexed over some time t , where t appears at multiple instances. Namely, fix t for all but one instance and differentiate. Then take the sum of all of the results. We illustrate this by an example.

Example 2.15. Let X_t be some time dependent vector field on a manifold M with flow ϕ_t . Furthermore, let α_t be a smooth family of differential forms and f_t a smooth family of real valued functions on M . We want to compute:

$$\frac{d}{dt} \phi_t^* (e^{f_t} \alpha_t).$$

Begin by fixing all but the first occurrence of t and differentiate, i.e. compute

$$\begin{aligned} \frac{\partial}{\partial t} \phi_t^* (e^{f_s} \alpha_s) &= \phi_t^* (\mathcal{L}_{X_t} (e^{f_s} \alpha_s)) = \phi_t^* ((X_t \cdot e^{f_s}) \alpha_s + e^{f_s} \mathcal{L}_{X_t} \alpha_s) \\ &= \phi_t^* ((X_t \cdot f_s) e^{f_s} \alpha_s + e^{f_s} \mathcal{L}_{X_t} \alpha_s) \end{aligned}$$

Here we used Proposition 2.8 and the Leibniz rule for the Lie derivative. Next fix all but the second occurrence of t and using Lemma 2.9 compute:

$$\frac{\partial}{\partial t} \phi_s^*(e^{f_t} \alpha_s) = \phi_s^* \left(\frac{\partial}{\partial t} e^{f_t} \alpha_s \right) = \phi_s^*(\dot{f}_t e^{f_t} \alpha_s)$$

And lastly

$$\frac{\partial}{\partial t} \phi_s^*(e^{f_s} \alpha_t) = \phi_s^* \left(e^{f_s} \frac{\partial}{\partial t} \alpha_t \right) = \phi_s^*(e^{f_s} \dot{\alpha}_t)$$

Finally plug in $s = t$ and sum the results:

$$\begin{aligned} \frac{d}{dt} \phi_t^*(e^{f_t} \alpha_t) &= \phi_t^*((X_t \cdot f_t) e^{f_t} \alpha_t + e^{f_t} \mathcal{L}_{X_t} \alpha_t) + \phi_t^*(\dot{f}_t e^{f_t} \alpha_t) + \phi_t^*(e^{f_t} \dot{\alpha}_t) \\ &= \phi_t^*(e^{f_t} ((X_t \cdot f_t) \alpha_t + \mathcal{L}_{X_t} \alpha_t + \dot{f}_t \alpha_t + \dot{\alpha}_t)) \end{aligned}$$

2.3 Formulas for the Hodge star operator and codifferential

In this section we briefly discuss the definition of the Hodge star operator and codifferential as well as their basic properties. This is supposed to serve as a canonical reference for the correct signs in the formulas involved. After the basic properties we proof a few more formulas that might be interesting.

Let $(V, \langle \cdot, \cdot \rangle)$ be an oriented n -dimensional euclidian vector space. Then the Hodge star operator is the unique linear map

$$\star: \Lambda^k V \rightarrow \Lambda^{n-k} V$$

satisfying

$$\alpha \wedge \star \beta = \langle \alpha, \beta \rangle \text{vol},$$

for all $\alpha, \beta \in \Lambda^k V$, where $\text{vol} \in \Lambda^n V$ is the canonical volume form on V . If $e_1, \dots, e_n \in V$ is an oriented orthonormal basis we see that for any ordered tuple $(i_1, \dots, i_k)$

$$\star(e_{i_1} \wedge \dots \wedge e_{i_k}) = e_{j_1} \wedge \dots \wedge e_{j_{n-k}},$$

where $(j_1, \dots, j_{n-k})$ is the complementary ordered tuple. Since tuples of this form define an orthonormal basis it follows that $\star$ is an isometry. Using this fact we compute

$$\begin{aligned} \langle \star \star \beta, \alpha \rangle \text{vol} &= \star \star \beta \wedge \star \alpha = (-1)^{k(n-k)} \star \alpha \wedge \star \star \beta \\ &= (-1)^{k(n-k)} \langle \star \alpha, \star \beta \rangle \text{vol} = (-1)^{k(n-k)} \langle \alpha, \beta \rangle \text{vol} \end{aligned}$$

from which we conclude that

$$\star \star = (-1)^{k(n-k)} \text{id}_{\Lambda^k V}.$$

The Hodge star operator can also be used to relate the interior and exterior product. For $v \in V$ and $\omega \in \Lambda^k V^*$ we denote the interior product by $v \lrcorner \omega \in \Lambda^{k-1} V^*$, that is

$$(v \lrcorner \omega)(v_1, \dots, v_{k-1}) := \omega(v, v_1, \dots, v_{k-1}).$$

(Some authors define this with the opposite sign, i.e. $v \lrcorner \omega := -\omega(v, -)$). Moreover the notation $v \wedge \omega \in \Lambda^{k+1} V^*$ is to be understood as first dualizing v and then taking the usual wedge product, that is

$$v \wedge \omega := \langle v, - \rangle \wedge \omega.$$

We then have the following Lemma.

Lemma 2.16. *For $v \in V$, $\omega \in \Lambda^k V$ and $\eta \in \Lambda^{k-1} V$ the following holds*

- (i) $\langle \eta, v \lrcorner \omega \rangle = \langle v \wedge \eta, \omega \rangle$
- (ii) $v \lrcorner \omega = (-1)^{n(k+1)} \star(v \wedge \star \omega)$

Proof. For (i) note that the expression on both sides is linear in v, ω and η , such that it suffices to show equality on basis vectors. So let $e_1, \dots, e_n \in V$ be an oriented orthonormal basis and $\varepsilon^1, \dots, \varepsilon^n \in V^*$ its dual basis. We have to compute (i) for

$$v = e_m, \quad \omega = \varepsilon^{i_1} \wedge \dots \wedge \varepsilon^{i_k}, \quad \eta = \varepsilon^{j_1} \wedge \dots \wedge \varepsilon^{j_{k-1}}.$$

First note that we may assume $m = i_p$ and

$$(i_1, \dots, \widehat{i_p}, \dots, i_k) = (j_1, \dots, j_{k-1}).$$

for some $1 \leq p \leq k$ since otherwise both sides vanish. But then

$$v \lrcorner \omega = (-1)^{p-1} \eta$$

and

$$v \wedge \eta = (-1)^{p-1} \omega$$

from which (i) follows. Using this we can show (ii):

$$\begin{aligned} \langle \eta, v \lrcorner \omega \rangle \text{vol} &= \langle v \wedge \eta, \omega \rangle \text{vol} \\ &= v \wedge \eta \wedge \star \omega \\ &= (-1)^{k-1} \eta \wedge v \wedge \star \omega \\ &= (-1)^{k-1+(n-k+1)(k-1)} \eta \wedge \star \star (v \wedge \star \omega) \\ &= (-1)^{(n-k+2)(k-1)} \langle \eta, \star(v \wedge \star \omega) \rangle \text{vol} \end{aligned}$$

Moreover we compute modulo 2:

$$(n-k+2)(k-1) = (n-k)(k-1) = n(k-1) - k(k-1) = n(k+1)$$

Since η was arbitrary the statement follows. $\square$

Remark 2.17. Part (i) of Lemma 2.16 states that the interior product is the adjoint of the exterior product.

Next we show a useful formula for the exterior derivative and codifferential in terms of the Levi-Civita connection. Let (M, g) be an oriented n -dimensional Riemannian manifold. First let's fix a sign convention for the codifferential δ . On k -forms we define δ by

$$\begin{array}{ccc} \Omega^k(M) & \xrightarrow{(-1)^k \delta} & \Omega^{k-1}(M) \\ \downarrow \star & & \downarrow \star \\ \Omega^{n-k}(M) & \xrightarrow{d} & \Omega^{n-k+1}(M) \end{array}$$

or in formulas

$$\delta := (-1)^k \star^{-1} d \star = (-1)^{n(k-1)+1} \star d \star: \Omega^k(M) \rightarrow \Omega^{k-1}(M).$$

With this sign convention we have for $\omega \in \Omega^k(M)$ and $\eta \in \Omega^{k+1}(M)$

$$d(\omega \wedge \star \eta) = d\omega \wedge \star \eta - \omega \wedge \star d\eta. \quad (4)$$

If $\alpha, \beta \in \Omega^k(M)$, such that the intersection of their supports is compact, we denote by

$$\langle \langle \alpha, \beta \rangle \rangle := \int_M \langle \alpha, \beta \rangle \text{vol}$$

their L^2 -scalar product. Using Stoke's Theorem it then follows from (4):

Lemma 2.18. *Let $\omega \in \Omega^k(M)$ and $\eta \in \Omega^{k+1}(M)$, such that the intersection of their supports is compact and contained in $M \setminus \partial M$. Then*

$$\langle \langle d\omega, \eta \rangle \rangle = \langle \langle \omega, \delta\eta \rangle \rangle.$$

With these sign conventions in mind we can now show:

Lemma 2.19. *Let M be a Riemannian manifold and $\omega \in \Omega^*(M)$. Let $e_1, \dots, e_n$ be an orthonormal local frame of TM , then*

$$d\omega = \sum_{i=1}^n e_i \wedge \nabla_i \omega,$$

where $\nabla_i \omega := \nabla_{e_i} \omega$ denotes the connection on the bundle $\Lambda^* T^* M$ induced by the Levi-Civita connection on M . Moreover, if M is oriented we have

$$\delta\omega = - \sum_{i=1}^n e_i \lrcorner \nabla_i \omega.$$

Proof. We first show the formula for the exterior derivative for 1-forms $\alpha \in \Omega^1(M)$. Using Corollary 2.7 and the symmetry of the Levi-Civita connection we get

$$\begin{aligned} d\alpha(X, Y) &= X \cdot \alpha(Y) - Y \cdot \alpha(X) - \alpha([X, Y]) \\ &= \nabla_X(\alpha(Y)) - \nabla_Y(\alpha(X)) - \alpha(\nabla_X Y - \nabla_Y X) \\ &= (\nabla_X \alpha)(Y) - (\nabla_Y \alpha)(X) \end{aligned}$$

for any smooth vector fields $X, Y \in \mathcal{X}(M)$. Let $\varepsilon^1, \dots, \varepsilon^n$ be the dual frame of $e_1, \dots, e_n$. It then follows that

$$\begin{aligned} d\alpha &= \sum_{i,j=1}^n d\alpha(e_i, e_j) \varepsilon^i \otimes \varepsilon^j \\ &= \sum_{i=1}^n \varepsilon^i \otimes \left(\sum_{j=1}^n (\nabla_i \alpha)(e_j) \varepsilon^j \right) - \sum_{j=1}^n \left(\sum_{i=1}^n (\nabla_j \alpha)(e_i) \varepsilon^i \right) \otimes \varepsilon^j \\ &= \sum_{i=1}^n \varepsilon^i \otimes \nabla_i \alpha - \sum_{j=1}^n \nabla_j \alpha \otimes \varepsilon^j \\ &= \sum_{i=1}^n \varepsilon^i \wedge \nabla_i \alpha \\ &= \sum_{i=1}^n e_i \wedge \nabla_i \alpha \end{aligned}$$

where we have used $\varepsilon^i = \langle e_i, - \rangle$ in the last equality (since $e_1, \dots, e_n$ is orthonormal). Now define a differential operator D by

$$D\omega := \sum_{i=1}^n e_i \wedge \nabla_i \omega.$$

Clearly D agrees with the exterior derivative on 0-forms and we have shown above that they furthermore agree on 1-forms. Moreover

$$\begin{aligned} D(\omega \wedge \eta) &= \sum_{i=1}^n e_i \wedge \nabla_i \omega \wedge \eta + e_i \wedge \omega \wedge \nabla_i \eta \\ &= (D\omega) \wedge \eta + (-1)^{|\omega|} \omega \wedge D\eta. \end{aligned}$$

So D and d also satisfy the same Leibniz rule. It follows that $D = d$. From this we can deduce the

formula for the codifferential by using Lemma 2.16. So let $\omega \in \Omega^k(M)$.

$$\begin{aligned}
\delta\omega &= (-1)^{n(k-1)+1} \star d \star \omega \\
&= (-1)^{n(k-1)+1} \sum_{i=1}^n \star(e_i \wedge \nabla_i \star \omega) \\
&= (-1)^{n(k-1)+1} \sum_{i=1}^n \star(e_i \wedge \star \nabla_i \omega) \\
&= (-1)^{n(k-1)+1+n(k+1)} \sum_{i=1}^n e_i \lrcorner \nabla_i \omega \\
&= - \sum_{i=1}^n e_i \lrcorner \nabla_i \omega.
\end{aligned}$$

In the third equality we used the fact that covariant differentiation and the Hodge star operator commute, as shown in the Lemma 2.20 below. $\square$

Lemma 2.20. *Let M be a an oriented Riemannian manifold, $\omega \in \Omega^*(M)$ and $X \in \mathcal{X}(M)$. Then*

$$\nabla_X \star \omega = \star \nabla_X \omega.$$

Proof. Let $\omega \in \Omega^k(M)$ and $\eta \in \Omega^{n-k}(M)$. Recall that the volume form $\text{vol} \in \Omega^n(M)$ is parallel, i.e. $\nabla_X \text{vol} = 0$. Hence

$$\nabla_X (\langle \eta, \star \omega \rangle \text{vol}) = (\langle \nabla_X \eta, \star \omega \rangle + \langle \eta, \nabla_X \star \omega \rangle) \text{vol}$$

On the other hand

$$\begin{aligned}
\nabla_X (\langle \eta, \star \omega \rangle \text{vol}) &= \nabla_X (\eta \wedge \star \star \omega) \\
&= (-1)^{k(n-k)} (\nabla_X \eta \wedge \star \star \omega + \eta \wedge \nabla_X \star \star \omega) \\
&= \nabla_X \eta \wedge \star \star \omega + \eta \wedge \star \star \nabla_X \omega \\
&= (\langle \nabla_X \eta, \star \omega \rangle + \langle \eta, \star \nabla_X \omega \rangle) \text{vol}.
\end{aligned}$$

Thus

$$\langle \eta, \nabla_X \star \omega \rangle = \langle \eta, \star \nabla_X \omega \rangle.$$

Since η was arbitrary the statement follows. $\square$

3 Riemannian Geometry

3.1 Product Manifolds

In the following let (M_1, g_1) and (M_2, g_2) be Riemannian manifolds. Then we can consider the product manifold

$$M := M_1 \times M_2$$

equipped with the product metric $g := g_1 \times g_2$ defined by

$$g(X_1 + X_2, Y_1 + Y_2) = g_1(X_1, Y_1) + g_2(X_2, Y_2), \quad (5)$$

where $X_i, Y_i \in \mathcal{X}(M_i)$. In this section we compute the Levi-Civita connection, curvature tensors, covariant derivative along curves and geodesics of (M, g) and see how they relate to the corresponding objects on (M_1, g_1) and (M_2, g_2) .

Let us establish some notation used in the discussion that follows. For a vector field $Z \in \mathcal{X}(M_1)$ we write $\tilde{Z} \in \mathcal{X}(M)$ for the horizontal lift of Z , that is

$$\tilde{Z}|_{(p_1, p_2)} := Z|_{p_1}$$

for $(p_1, p_2) \in M$. Moreover for a smooth function $f \in C^\infty(M_1)$ we define

$$\tilde{f} := f \circ \pi_1 \in C^\infty(M),$$

where $\pi_1: M \rightarrow M_1$ denotes the projection. The same notation is used for vector fields and smooth functions on M_2 . So to be precise, equation (5) should read

$$g(\tilde{X}_1 + \tilde{X}_2, \tilde{Y}_1 + \tilde{Y}_2) = \widetilde{g_1(X_1, Y_1)} + \widetilde{g_2(X_2, Y_2)}.$$

For the sake of notational simplicity we will often suppress these subtleties, if there is no risk of confusion and only write $X_1 + X_2$ instead of $\tilde{X}_1 + \tilde{X}_2$ or f instead of $\tilde{f}$. It should always be clear from the context what is meant. Let us moreover denote by $\nabla^1, D_t^1, \text{Rm}_1, \text{Ric}_1, \text{scal}_1$ the Levi-Civita connection, covariant derivative along curves, Riemannian curvature tensor, Ricci tensor and scalar curvature of (M_1, g_1) respectively. Similarly we denote the same objects on (M_2, g_2) by $\nabla^2, D_t^2, \text{Rm}_2, \text{Ric}_2, \text{scal}_2$ and on (M, g) simply by $\nabla, D_t, \text{Rm}, \text{Ric}, \text{scal}$. Finally we define $m_1 := \dim(M_1)$ and $m_2 := \dim(M_2)$.

Of course our first step will be to compute ∇ in terms of ∇^1 and ∇^2 . For this we use the characterisation of the canonical connection on a product vector bundle shown in Proposition 3.5.

Proposition 3.1. *The Levi-Civita connection ∇ on (M, g) is given by*

$$\nabla = \nabla^1 \times \nabla^2.$$

Proof. To ensure clarity, we will be more precise with the notation in this proof, as explained at the beginning of the section. Let $X, Y \in \mathcal{X}(M)$ be of the form $X = \tilde{X}_1 + \tilde{X}_2, Y = \tilde{Y}_1 + \tilde{Y}_2$, where $X_i, Y_i \in \mathcal{X}(M_i)$. By Proposition 3.5 it suffices to show that the Levi Civita connection satisfies

$$\nabla_X Y = \widetilde{\nabla_{X_1}^1 Y_1} + \widetilde{\nabla_{X_2}^2 Y_2}.$$

We would like to use the Koszul formula to compute $\nabla_X Y$ and begin by simplifying some of the terms involved in this formula. For $Z \in \mathcal{X}(M_1), W \in \mathcal{X}(M_2)$ we claim that

$$[\tilde{Z}, \tilde{W}] = 0.$$

To see this choose coordinates $x^1, \dots, x^{m_1}$ of M_1 and $y^1, \dots, y^{m_2}$ of M_2 . Write

$$Z = f^i \frac{\partial}{\partial x^i}, \quad W = g^j \frac{\partial}{\partial y^j}$$

for smooth functions f^i and g^j defined on the chosen coordinate neighbourhoods of M_1 and M_2 respectively. Then $x^1, \dots, x^{m_1}, y^1, \dots, y^{m_2}$ define coordinates of M and

$$\tilde{Z} = \tilde{f}^i \frac{\partial}{\partial x^i}, \quad \tilde{W} = \tilde{g}^j \frac{\partial}{\partial y^j}.$$

We now compute

$$\begin{aligned} [\tilde{Z}, \tilde{W}] &= \tilde{f}^i \frac{\partial \tilde{g}^j}{\partial x^i} \frac{\partial}{\partial y^j} + \tilde{g}^j \left[\tilde{f}^i \frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^j} \right] \\ &= \tilde{f}^i \frac{\partial \tilde{g}^j}{\partial x^i} \frac{\partial}{\partial y^j} - \tilde{g}^j \frac{\partial \tilde{f}^i}{\partial y^j} \frac{\partial}{\partial x^i} - \tilde{g}^j \tilde{f}^i \left[\frac{\partial}{\partial y^j}, \frac{\partial}{\partial x^i} \right]. \end{aligned}$$

Since $\tilde{g}^j$ does not depend on x^i we get

$$\frac{\partial \tilde{g}^j}{\partial x^i} = 0.$$

Similiarly

$$\frac{\partial \tilde{f}^i}{\partial y^j} = 0,$$

such that $[\tilde{Z}, \tilde{W}] = 0$ follows. For Z as above the Koszul formula now reads

$$\begin{aligned} g(\nabla_X Y, \tilde{Z}) &= \frac{1}{2} \left(Xg(Y, \tilde{Z}) + Yg(\tilde{Z}, X) - \tilde{Z}g(X, Y) \right. \\ &\quad \left. - g(Y, [X, \tilde{Z}]) - g(\tilde{Z}, [Y, X]) + g(X, [\tilde{Z}, Y]) \right) \\ &= \frac{1}{2} \left(\tilde{X}_1 g(\tilde{Y}_1, \tilde{Z}) + \tilde{Y}_1 g(\tilde{Z}, \tilde{X}_1) - \tilde{Z}g(\tilde{X}_1, \tilde{Y}_1) \right. \\ &\quad \left. - g(\tilde{Y}_1, [\tilde{X}_1, \tilde{Z}]) - g(\tilde{Z}, [\tilde{Y}_1, \tilde{X}_1]) + g(\tilde{X}_1, [\tilde{Z}, \tilde{Y}_1]) \right) \end{aligned} \quad (6)$$

Besides the claim about the Lie bracket shown above we have further used, that $g(Y, \tilde{Z}) = g(\tilde{Y}_1, \tilde{Z})$ is constant in the M_2 -direction, such that $\tilde{X}_2 g(\tilde{Y}_1, \tilde{Z}) = 0$, hence

$$Xg(Y, \tilde{Z}) = \tilde{X}_2 g(\tilde{Y}_1, \tilde{Z}).$$

An analogous statement holds for the second and third term. Moreover $[\tilde{X}_1, \tilde{Z}] = \widetilde{[X, Z]}$, such that $g(\tilde{Y}_2, [\tilde{X}_1, \tilde{Z}]) = 0$, hence

$$g(Y, [X, \tilde{Z}]) = g(\tilde{Y}_1, [\tilde{x}_1, \tilde{Z}]).$$

Again analogous statements hold for the last two terms. Applying the Koszul formula to $\nabla_{X_1}^1 Y_1$ we get

$$\begin{aligned} g_1(\nabla_{X_1}^1 Y_1, Z) &= \frac{1}{2} \left(X_1 g_1(Y_1, Z) + Y_1 g_1(Z, X_1) - Z g_1(X_1, Y_1) \right. \\ &\quad \left. - g_1(Y_1, [X_1, Z]) - g_1(Z, [Y_1, X_1]) + g_1(X_1, [Z, Y_1]) \right) \end{aligned} \quad (7)$$

To compare this expression to (6) we note the following identities for $U, V \in \mathcal{X}(M_1)$ and $f \in C^\infty(M_1)$:

$$g(\tilde{U}, \tilde{V}) = \widetilde{g(U, V)}$$

just by definition of g and

$$\tilde{V} \tilde{f} = \widetilde{Vf}.$$

So (6) and (7) now yield

$$g(\nabla_X Y, \tilde{Z}) = \widetilde{g_1(\nabla_{X_1}^1 Y_1, Z)} = g(\nabla_{X_1}^1 Y_1, \tilde{Z}).$$

Similiarly we get for $W \in \mathcal{X}(M_2)$

$$g(\nabla_X Y, \tilde{Z}) = g(\widetilde{\nabla_{X_2}^2 Y_2}, \tilde{W}).$$

Together this implies

$$g(\nabla_X Y, \tilde{Z} + \tilde{W}) = g(\widetilde{\nabla_{X_1}^1 Y_1} + \widetilde{\nabla_{X_2}^2 Y_2}, \tilde{Z} + \tilde{W}).$$

From which the assertion follows. □

Knowing the Levi-Civita connection on M , it is now easy to compute the curvature, which behaves just as expected.

Corollary 3.2. *Let $X = X_1 + X_2$, $Y = Y_1 + Y_2$, $Z = Z_1 + Z_2$, $W = W_1 + W_2$ be smooth vector fields on M , where $X_i, Y_i, Z_i, W_i \in \mathcal{X}$. Then the following holds*

- (i) $\text{Rm}(X, Y, Z, W) = \text{Rm}_1(X_1, Y_1, Z_1, W_1) + \text{Rm}_2(X_2, Y_2, Z_2, W_2)$
- (ii) $\text{Ric}(X, Y) = \text{Ric}_1(X_1, Y_1) + \text{Ric}_2(X_2, Y_2)$
- (iii) $\text{scal} = \text{scal}_1 + \text{scal}_2$

Proof. By Proposition 3.1

$$\nabla_X \nabla_Y Z = \nabla_X (\nabla_{Y_1}^1 Z_1 + \nabla_{Y_2}^2 Z_2) = \nabla_{X_1}^1 \nabla_{Y_1}^1 Z_1 + \nabla_{X_2}^2 \nabla_{Y_2}^2 Z_2.$$

An analogous equation holds for $\nabla_Y \nabla_X Z$. Moreover

$$[X, Y] = [X_1, Y_1] + [X_2, Y_2]$$

as we have seen in the proof of Proposition 3.1, such that

$$\nabla_{[X, Y]} Z = \nabla_{[X_1, Y_1]}^1 Z_1 + \nabla_{[X_2, Y_2]}^2 Z_2.$$

Then (i) follows from the definition of Rm and the product metric g . For (ii) write

$$\text{Ric}(X, Y) = \sum_{i=1}^{m_1} \text{Rm}(X, E_i^1, E_i^1, Y) + \sum_{j=1}^{m_2} \text{Rm}(X, E_j^2, E_j^2, Y)$$

for local orthonormal frames $E_1^1, \dots, E_{m_1}^1$ of TM_1 and $E_1^2, \dots, E_{m_2}^2$ of TM_2 . By (i)

$$\begin{aligned} \text{Rm}(X, E_i^1, E_i^1, Y) &= \text{Rm}_1(X_1, E_i^1, E_i^1, Y_1) + \text{Rm}_2(X_2, 0, 0, Y_2) \\ &= \text{Rm}_1(X_1, E_i^1, E_i^1, Y_1). \end{aligned}$$

An analogous statement holds for the second term, such that

$$\begin{aligned} \text{Ric}(X, Y) &= \sum_{i=1}^{m_1} \text{Rm}_1(X_1, E_i^1, E_i^1, Y_1) + \sum_{j=1}^{m_2} \text{Rm}_2(X_2, E_j^2, E_j^2, Y_2) \\ &= \text{Ric}_1(X_1, Y_1) + \text{Ric}_2(X_2, Y_2). \end{aligned}$$

Finally (iii) follows from (ii) in the same way. $\square$

Finally we compute the covariant derivative along curves and geodesics of M , which again behave in exactly the way one would expect.

Corollary 3.3. *Let $\gamma = (\gamma_1, \gamma_2): I \rightarrow M$ be a smooth curve and $V: I \rightarrow \gamma^* TM \cong \gamma_1^* TM_1 \oplus \gamma_2^* TM_2$ a smooth vector field along γ . Write $V = V_1 + V_2$ for $V_i: I \rightarrow \gamma_i^* TM$. Then*

$$D_t V = D_t^1 V_1 + D_t^2 V_2.$$

Proof. As in the proof of Proposition 3.1, we will take some care with the notation in this proof. To show the statement we compute both sides locally. To this end let $E_1^1, \dots, E_{m_1}^1$ be a local frame of TM_1 and $E_1^2, \dots, E_{m_2}^2$ a local frame of TM_2 . Write

$$V_1(t) = V_1^i(t) E_i^1|_{\gamma_1(t)}$$

and

$$V_2(t) = V_2^j(t) E_j^2|_{\gamma_2(t)}.$$

Then

$$V(t) = V_1^i(t) \tilde{E}_i^1|_{\gamma(t)} + V_2^j(t) \tilde{E}_j^2|_{\gamma(t)}$$

and hence

$$\begin{aligned} D_t V &= \dot{V}_1^i(t) \tilde{E}_i^1|_{\gamma(t)} + V_1^i(t) \nabla_{\dot{\gamma}(t)} \tilde{E}_i^1 \\ &\quad + \dot{V}_2^j(t) \tilde{E}_j^2|_{\gamma(t)} + V_2^j(t) \nabla_{\dot{\gamma}(t)} \tilde{E}_j^2. \end{aligned}$$

Moreover

$$D_t^1 V_1 = \dot{V}_1^i(t) E_i^1|_{\gamma_1(t)} + V_1^i(t) \nabla_{\dot{\gamma}_1(t)}^1 E_i^1$$

and similarly for $D_t^2 V_2$. By Proposition 3.1

$$\nabla_{\dot{\gamma}(t)} \tilde{E}_i^1 = \widetilde{\nabla_{\dot{\gamma}_1(t)}^1 E_i^1} \quad \text{and} \quad \nabla_{\dot{\gamma}(t)} \tilde{E}_j^2 = \widetilde{\nabla_{\dot{\gamma}_2(t)}^2 E_j^2}$$

and thus

$$D_t V = \widetilde{D_t^1 V_1} + \widetilde{D_t^2 V_2}.$$

□

Corollary 3.4. *Let $\gamma = (\gamma_1, \gamma_2): I \rightarrow M$ be a smooth curve. Then γ is a geodesic if and only if γ_1 and γ_2 are geodesics.*

Proof. This now follows immediately from the formula from Corollary 3.3. □

3.2 Induced connections

Given any vector bundle there are many ways to construct new ones from it. In this section we show how a connection on the new bundle is induced by a connection on the bundle we started with.

In the following proposition we shall use the same notational convention for sections in the product bundle as for vector fields introduced at the beginning of the last section.

Proposition 3.5. *Let $E_1 \rightarrow M_1$ and $E_2 \rightarrow M_2$ be smooth vector bundles and let ∇^1, ∇^2 be connections on E_1, E_2 respectively. Then there exists a unique connection ∇ on the product bundle*

$$E_1 \times E_2 \rightarrow M_1 \times M_2,$$

such that for sections $s \in \Gamma(E_1 \times E_2)$ of the form $s = s_1 + s_2$, where $s_i \in \Gamma(E_i)$, and vector fields $X \in \mathcal{X}(M_1 \times M_2)$ of the form $X = X_1 + X_2$, where $X_i \in \mathcal{X}(M_i)$, the following holds:

$$\nabla_X s = \nabla_{X_1}^1 s_1 + \nabla_{X_2}^2 s_2.$$

We call ∇ the product connection of ∇^1 and ∇^2 and write $\nabla = \nabla^1 \times \nabla^2$.

Proof. We first show uniqueness. Let $X \in \mathcal{X}(M_1 \times M_2)$ be an arbitrary vector field and $s \in \Gamma(E_1 \times E_2)$ an arbitrary section. For $(p_1, p_2) \in M_1 \times M_2$ we can find open neighbourhoods $U_i \subseteq M_i$ of p_i , such that there exist local frames

$$\sigma_1^i, \dots, \sigma_{n_i}^i \in \Gamma(E_i|_{U_i}),$$

where n_i is the rank of E_i and there exist local frames

$$E_1^i, \dots, E_{m_i}^i \in \Gamma(TM_i|_{U_i}),$$

On $U := U_1 \times U_2$ we can write

$$s = s_1^{j_1} \sigma_{j_1}^1 + s_2^{j_2} \sigma_{j_2}^2$$

and

$$X = X_1^{k_1} E_{k_1}^1 + X_2^{k_2} E_{k_2}^2$$

for smooth functions $s_1^{j_1}, s_2^{j_2}, X_1^{k_1}, X_2^{k_2} \in C^\infty(U)$. Then

$$\begin{aligned} \nabla_X s &= X_1^{k_1} \left(\nabla_{E_{k_1}^1} (s_1^{j_1} \sigma_{j_1}^1) + \nabla_{E_{k_1}^1} (s_2^{j_2} \sigma_{j_2}^2) \right) + X_2^{k_2} \left(\nabla_{E_{k_2}^2} (s_1^{j_1} \sigma_{j_1}^1) + \nabla_{E_{k_2}^2} (s_2^{j_2} \sigma_{j_2}^2) \right) \\ &= X_1^{k_1} \left((E_{k_1}^1 \cdot s_1^{j_1}) \sigma_{j_1}^1 + s_1^{j_1} \nabla_{E_{k_1}^1}^1 \sigma_{j_1}^1 + (E_{k_1}^1 \cdot s_2^{j_2}) \sigma_{j_2}^2 \right) \\ &\quad + X_2^{k_2} \left((E_{k_2}^2 \cdot s_1^{j_1}) \sigma_{j_1}^1 + (E_{k_2}^2 \cdot s_2^{j_2}) \sigma_{j_2}^2 + s_2^{j_2} \nabla_{E_{k_2}^2}^2 \sigma_{j_2}^2 \right) \end{aligned} \tag{8}$$

Since the last expression does not depend on ∇ this shows uniqueness. To show existence define a connection ∇^U on U by formula (8). It is easy to see that this does in fact define a connection on $E_1 \times E_2|_U$, which is compatible with the restricted connections $\nabla^1|_{U_1}$ and $\nabla^2|_{U_2}$ in the sense of the statement of the proposition. So by the uniqueness result this connection does not depend on the chosen frames, thus we get a well-defined connection ∇ on $E_1 \times E_2$ by

$$(\nabla_X s)|_U := \nabla_{X|_U}^U s|_U.$$

□

The next proposition is very similar and concerned with the induced connection on a tensor bundle.

Proposition 3.6. *Let $E_1 \rightarrow M$ and $E_2 \rightarrow M$ be smooth vector bundles and let ∇^1, ∇^2 be connections on E_1, E_2 respectively. Then there exists a unique connection ∇ on the tensor product bundle*

$$E_1 \otimes E_2 \rightarrow M,$$

such that for any vector field $X \in \mathcal{X}(M)$ and sections $s_1 \in \Gamma(E_1)$ and $s_2 \in \Gamma(E_2)$ the following holds:

$$\nabla_X(s_1 \otimes s_2) = \nabla_X^1 s_1 \otimes s_2 + s_1 \otimes \nabla_X^2 s_2.$$

Proof. The proof is analogous to the proof of Proposition 3.5, so we only summarize the idea briefly. One notes that any section $s \in \Gamma(E_1 \otimes E_2)$ can be written locally as

$$s = \sum_{i,j} s_1^i \otimes s_2^j,$$

where s_1^i and s_2^j are local sections of E_1 and E_2 respectively. This then shows uniqueness and moreover gives us a way to define ∇ locally, which can be pasted to together to define ∇ globally just as in Proposition 3.5. □

Finally we discuss the induced connection on the pullback bundle f^*E of some vector bundle $E \rightarrow M$ under a smooth map $f: N \rightarrow M$. For this we introduce the following notation: given a section s of E we can define a section f^*s of f^*E by $(f^*s)(p) := s(f(p))$.

Proposition 3.7. *Let $E \rightarrow M$ be a vector bundle with connection ∇ and let $f: N \rightarrow M$ be a smooth map. Then there exists a unique connection $f^*\nabla$ on the pullback bundle f^*E , such that for all sections $s \in \Gamma(E)$ and tangent vectors $v \in T_q N$ the following holds:*

$$(f^*\nabla)_v(f^*s) = \nabla_{f_*v} s,$$

where $f_*v = d_q f(v)$.

Proof. As usual we begin by showing uniqueness. So suppose we are given a connection $f^*\nabla$ with this property. Let $s \in \Gamma(f^*E)$ be any section and choose a local frame $e_1, \dots, e_k$ of E . Then by definition of the pullback bundle $f^*e_1, \dots, f^*e_k$ is a local frame of f^*E . Write $s = s^i f^*e_i$. Then

$$(f^*\nabla)_v s = v \cdot s^i f^*e_i + s^i \nabla_{f_*v} e_i.$$

Since the right hand side does not depend on the connection on f^*E , this shows uniqueness. Now use the above formula to define $f^*\nabla$. As this is a pointwise definition we still need to check smoothness, i.e. that $(f^*\nabla)_X s$ is a smooth section for $X \in \mathcal{X}(N)$ a smooth vector field. As a composition of smooth maps the map $f_*X = df \circ X: N \rightarrow TM$ is smooth, hence can be written locally as

$$f_*X = g^j f^* \partial_j$$

for smooth functions $g^j: N \rightarrow \mathbb{R}$ and a local frame $\partial_1, \dots, \partial_n$ of TM . Then

$$\nabla_{f_*X} e_i = g^j \nabla_{f^* \partial_j} e_i = g^j f^*(\nabla_{\partial_j} e_i),$$

which is clearly smooth. The other properties of a connection are also evidently satisfied by $f^*\nabla$, such that the statement follows. □

To conclude this section we present two useful formulas for induced connections.

Lemma 3.8. *Let M be a Riemannian manifold, $\omega, \eta \in \Omega^*(M)$ differential forms and $X \in \mathcal{X}(M)$ a vector field. Then*

$$\nabla_X(\omega \wedge \eta) = \nabla_X \omega \wedge \eta + \omega \wedge \nabla_X \eta,$$

where ∇_X denotes the connection on the bundle $\Lambda^* T^* M$ induced by the Levi-Civita connection on M .

Proof. By definition of the wedge product it suffices to show that ∇_X commutes with the anti-symmetrization map

$$\text{Alt}: (T^* M)^{\otimes k} \rightarrow (T^* M)^{\otimes k}.$$

To see this let $\alpha_1, \dots, \alpha_k \in \Omega^1(M)$. Then

$$\begin{aligned} \nabla_X(\text{Alt}(\alpha_1 \otimes \dots \otimes \alpha_k)) &= \nabla_X \left(\frac{1}{k!} \sum_{\sigma} \text{sgn}(\sigma) \alpha_{\sigma(1)} \otimes \dots \otimes \alpha_{\sigma(k)} \right) \\ &= \frac{1}{k!} \sum_{\sigma} \sum_i \text{sgn}(\sigma) \alpha_{\sigma(1)} \otimes \dots \otimes \nabla_X \alpha_{\sigma(i)} \otimes \dots \otimes \alpha_{\sigma(k)} \end{aligned}$$

and

$$\begin{aligned} \text{Alt}(\nabla_X(\alpha_1 \otimes \dots \otimes \alpha_k)) &= \text{Alt} \left(\sum_i \alpha_1 \otimes \dots \otimes \nabla_X \alpha_i \otimes \dots \otimes \alpha_k \right) \\ &= \sum_i \sum_{\sigma} \frac{1}{k!} \text{sgn}(\sigma) \alpha_{\sigma(1)} \otimes \dots \otimes \nabla_X \alpha_{\sigma(i)} \otimes \dots \otimes \alpha_{\sigma(k)}. \end{aligned}$$

Since locally any section $\omega \in \Gamma((T^* M)^{\otimes k})$ can be written as a sum of tensor products of 1-forms the statement follows. $\square$

Lemma 3.9. *Let $E \rightarrow M$ be a smooth vector bundle equipped with a metric and compatible connection. Then for any $X \in \mathcal{X}(M)$ the following diagram commutes*

$$\begin{array}{ccc} \Gamma(E) & \xrightarrow{\nabla_X} & \Gamma(E) \\ \downarrow \cong & & \downarrow \cong \\ \Gamma(E^*) & \xrightarrow{\nabla_X} & \Gamma(E^*) \end{array}$$

where the vertical isomorphisms are given by dualizing and the connection on E^* is the unique connection defined by

$$X \cdot (\alpha(s)) = (\nabla_X \alpha)(s) + \alpha(\nabla_X s)$$

for $s \in \Gamma(E), \alpha \in \Gamma(E^*)$.

Proof. This follows immediately from the definition of the connection on E^* and the compatibility of the metric. $\square$

3.3 Miscellaneous

In the following let M, N be Riemannian manifolds (without boundary).

Lemma 3.10. *Let $\gamma: (-\varepsilon, \varepsilon) \rightarrow M$ be a smooth curve. Then*

$$\lim_{t \rightarrow 0} \frac{d(\gamma(t), \gamma(0))}{|t|} = \|\dot{\gamma}(0)\|,$$

where d denotes the Riemannian distance function.

Proof. Choose normal coordinates around $p = \gamma(0)$ and write $\gamma = (\gamma_1, \dots, \gamma_n)$ in these coordinates. Then the distance from $\gamma(t)$ to $\gamma(0)$ is given by

$$d(\gamma(t), \gamma(0)) = \sqrt{\gamma_1^2(t) + \dots + \gamma_n^2(t)} = |(\gamma_1(t), \dots, \gamma_n(t))|,$$

where $|\cdot|$ denotes the euclidian norm. Thus

$$\begin{aligned}\lim_{t \rightarrow 0} \frac{d(\gamma(t), \gamma(0))}{|t|} &= \lim_{t \rightarrow 0} \frac{|(\gamma_1(t), \dots, \gamma_n(t))|}{|t|} \\ &= |(\dot{\gamma}_1(0), \dots, \dot{\gamma}_n(0))| \\ &= \|\dot{\gamma}(0)\|,\end{aligned}$$

where we used the fact that the metric on M is euclidian at p in normal coordinates. $\square$

Lemma 3.11. *A smooth map $f: M \rightarrow N$ is λ -Lipschitz if and only if*

$$\|d_p f(v)\| \leq \lambda \|v\|$$

for all $(p, v) \in TM$.

Proof. Suppose that f is λ -Lipschitz and assume without loss of generality that $\|v\| = 1$. Let $\gamma: (-\varepsilon, \varepsilon) \rightarrow M$ be a geodesic with $\gamma(0) = p$ and $\gamma'(0) = v$ and let d_M, d_N be the Riemannian distance functions on M, N respectively. Then

$$\begin{aligned}d_N(f(\gamma(t)), f(\gamma(0))) &\leq \lambda d_M(\gamma(t), \gamma(0)) \\ &= \lambda \operatorname{sgn}(t) \int_0^t \|\gamma'(s)\| ds \\ &= \lambda |t|,\end{aligned}$$

where we used $\|\gamma'(s)\| = \|v\| = 1$ in the last equality, which holds since γ is a geodesic. Thus

$$\frac{d_N(f(\gamma(t)), f(\gamma(0)))}{|t|} \leq \lambda$$

Letting $t \rightarrow 0$ on both sides we conclude from Lemma 3.10, that

$$\|d_p f(v)\| = \|(f \circ \gamma)'(0)\| \leq \lambda,$$

which shows the claim. For the converse let $x, y \in M$ and let $\gamma: [0, 1] \rightarrow M$ be any piecewise smooth curve from x to y . Then

$$\begin{aligned}d_N(f(x), f(y)) &\leq \int_0^1 \|(f \circ \gamma)'(t)\| dt \\ &= \int_0^1 \|d_{\gamma(t)} f(\gamma'(t))\| dt \\ &\leq \lambda \int_0^1 \|\gamma'(t)\| dt.\end{aligned}$$

Taking the infimum over all such curves γ shows

$$d_N(f(x), f(y)) \leq \lambda d_M(x, y).$$

$\square$

4 Chern Weil Theory

In this section we review the construction of the Chern-Weil homomorphism. This is largely based on [Kon26] but we add the missing details, fix some mistakes and give more precise definitions. In the following let $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$. All manifolds will be real but we consider real as well as complex vector bundles over them. If $\mathbb{K} = \mathbb{R}$ the vector bundles are understood to be real and if $\mathbb{K} = \mathbb{C}$ they are understood to be complex. There is virtually no difference between the complex and the real case in what follows.

4.1 Construction of the Chern Weil Homomorphism

Let us recall the definition of the connection and curvature form. Let $E \rightarrow M$ be a vector bundle with connection ∇ . For a local frame $e_1, \dots, e_k$ of E over $U \subseteq M$ define the **connection form** as the unique matrix valued 1-form

$$\Theta = (\Theta_{ij})_{i,j=1}^k \in \Omega^1(U, \mathbb{K}^{k \times k})$$

with

$$\nabla e_j = \sum_{i=1}^k \Theta_{ij} \otimes e_i.$$

Using the connection form we can compute the covariant derivative of any section $s \in \Omega^0(M, E)$ as follows. In the chosen frame write $s = s^j e_j$ for smooth functions $s^j \in C^\infty(M, \mathbb{K})$, then

$$\nabla s = ds^j \otimes e_j + \sum_{i=1}^k s^j \Theta_{ij} \otimes e_i = (d + \Theta)s,$$

where d acts by differentiation with respect to the local frame as written above and Θ acts by matrix multiplication in the coordinates of this frame.

The **curvature form** $\Omega \in \Omega^2(U, \mathbb{K}^{k \times k})$ is then defined as

$$\Omega := d\Theta + \Theta \wedge \Theta.$$

A computation shows that Ω transforms nicely under a change of frame and thus yields a globally defined object $\Omega \in \Omega^2(M, \text{End}(E))$. We sometimes write $\Omega = \Omega_\nabla$ to denote the dependence on the connection ∇ . The curvature form behaves well under pullback:

Lemma 4.1. *Let $E \rightarrow M$ be a vector bundle with connection ∇ and $f: N \rightarrow M$ a smooth map. Then*

$$f^* \Omega_\nabla = \Omega_{f^* \nabla},$$

where $f^* \nabla$ denotes the pullback connection on $f^* E$.

Proof. Let $e_1, \dots, e_k$ be a local frame, let Θ be the associated connection form and let $\tilde{e}_j := f^* e_j$ be the pullback frame. Then by definition of the pullback connection (cf. Proposition 3.7) we compute for any tangent vector $v \in T_q N$

$$(f^* \nabla)_v \tilde{e}_j = \nabla_{f_* v} e_j = \sum_{i=1}^k (\Theta_{ij})_{f(q)} (f_* v) e_i(f(q)) = \sum_{i=1}^k (f^* \Theta_{ij})_q(v) \tilde{e}_i(q).$$

Thus the connection form on $f^* E$ with respect to the frame $\tilde{e}_1, \dots, \tilde{e}_k$ is given by $f^* \Theta$. Since pullback commutes with the exterior derivative and the wedge product the statement now follows from the definition of the curvature form. $\square$

Chern-Weil theory now constructs cohomology classes (so called characteristic classes) of M out of the curvature form using certain invariant homogeneous polynomials, which we shall introduce next.

Definition 4.2. Let V be an n -dimensional (real or complex) vector space. An r -homogeneous polynomial on V is a map $\eta: V \rightarrow \mathbb{K}$, such that for all vectors $e_1, \dots, e_n \in V$ the map

$$\mathbb{K}^n \rightarrow \mathbb{K}, (t_1, \dots, t_n) \mapsto \eta(t_1 e_1 + \dots + t_n e_n)$$

is a (multivariate) r -homogeneous polynomial in the variables $t_1, \dots, t_n$. The space of all r -homogeneous polynomials on V is denoted by $\text{Pol}_r(V)$.

By choosing $e_1, \dots, e_n$ as a basis, it is easy to see that a map $\eta: V \rightarrow \mathbb{K}$ is an r -homogeneous polynomial if and only if for one (then all) linear isomorphisms $A: \mathbb{K}^n \rightarrow V$, the composition $\eta \circ A: \mathbb{K}^n \rightarrow \mathbb{K}$ is an r -homogeneous polynomial. Thus we can think of such polynomials as r -homogeneous polynomials in the entries of a vector $v = (v_1, \dots, v_n)$ with respect to some (then all) basis. With this definition it is easy to construct examples:

Example 4.3. The determinant $\det: \mathbb{K}^{n \times n} \rightarrow \mathbb{K}$ is an n -homogeneous polynomial and the trace $\text{tr}: \mathbb{K}^{n \times n} \rightarrow \mathbb{K}$ is a 1-homogeneous polynomial.

Yet another useful characterization of $\text{Pol}_r(V)$, which we will make use of later, can be given as follows.

Lemma 4.4. Let $\Sigma_{\mathbb{K}}^r(V^*)$ be the space of r -linear symmetric maps $V \times \dots \times V \rightarrow \mathbb{K}$. Then

$$\begin{aligned} \Phi_r: \Sigma_{\mathbb{K}}^r(V^*) &\rightarrow \text{Pol}_r(V) \\ \bar{\eta} &\mapsto (v \mapsto \bar{\eta}(v, \dots, v)) \end{aligned}$$

is an isomorphism of vector spaces.

Proof. This can be seen using a polarisation argument: If $\eta = \Phi_r(\bar{\eta})$ then

$$\eta(t_1 v_1 + \dots + t_r v_r) = \bar{\eta} \left(\sum_{i=1}^r t_i v_i, \dots, \sum_{i=1}^r t_i v_i \right)$$

for vectors $v_i \in V$ and scalars $t_i \in \mathbb{K}$. Using multilinearity and symmetry of $\bar{\eta}$ we see that this is a polynomial in $t_1, \dots, t_r$, where the coefficient in front of $t_1 \cdots t_r$ is given by $r! \bar{\eta}(v_1, \dots, v_r)$. So if $\eta = 0$ we conclude that $\bar{\eta}(v_1, \dots, v_r) = 0$. Hence Φ_r is injective. To show surjectivity we argue similarly. Let $\eta \in \text{Pol}_r(V)$ and let $e_1, \dots, e_n \in V$ be a basis. Then $\eta(t_1 e_1 + \dots + t_n e_n)$ is a polynomial in $t_1, \dots, t_n$. Let $a_{i_1, \dots, i_r}$ be the coefficient of this polynomial in front of $t_{i_1} \cdots t_{i_r}$, where we assume that $i_1 \leq \dots \leq i_r$. As above we see by comparing coefficients, that if $\eta = \Phi_r(\bar{\eta})$ then

$$\bar{\eta}(e_{i_1}, \dots, e_{i_r}) = \frac{1}{r!} a_{i_1, \dots, i_r}.$$

So we are led to define $\bar{\eta}$ on ordered tuples of basis vectors by this formula and extend it uniquely to a symmetric r -linear map. Then by definition the coefficients of the polynomials $\eta(t_1 e_1 + \dots + t_n e_n)$ and

$$\bar{\eta} \left(\sum_{i=1}^n t_i e_i, \dots, \sum_{i=1}^n t_i e_i \right)$$

agree, thus they are equal. Since $e_1, \dots, e_n$ is a basis, this shows

$$\eta(v) = \bar{\eta}(v, \dots, v).$$

□

We now turn our attention to homogeneous polynomials on the vector space $\mathbb{K}^{k \times k}$ of matrices, which are used to define characteristic classes in Chern-Weil theory.

Definition 4.5. An r -homogeneous polynomial $\eta: \mathbb{K}^{k \times k} \rightarrow \mathbb{K}$ is called invariant if

$$\eta(A^{-1}BA) = \eta(B)$$

for all $B \in \mathbb{K}^{k \times k}$ and $A \in \text{GL}(k, \mathbb{K})$. We denote the space of all such polynomials by $I_r(k) \subseteq \text{Pol}_r(\mathbb{K}^{k \times k})$.

Under the isomorphism Φ_r the invariant polynomials correspond to the invariant r -linear symmetric maps, where an r -linear symmetric map $\bar{\eta}: \mathbb{K}^{k \times k} \times \dots \times \mathbb{K}^{k \times k} \rightarrow \mathbb{K}$ is called invariant if

$$\bar{\eta}(A^{-1}B_1A, \dots, A^{-1}B_rA) = \bar{\eta}(B_1, \dots, B_r)$$

for all $B_1, \dots, B_r \in \mathbb{K}^{k \times k}$ and $A \in \text{GL}(k, \mathbb{K})$. Denote the space of all such maps by $S_r(k)$. We can then extend $\bar{\eta}$ to an r -linear map (here we don't need invariance yet)

$$\bar{\eta}: \Omega^1(M, \mathbb{K}^{k \times k}) \times \dots \times \Omega^1(M, \mathbb{K}^{k \times k}) \rightarrow \Omega^1(M, \mathbb{K}),$$

where $l = l_1 + \dots + l_r$ by setting

$$\bar{\eta}(\Omega_1, \dots, \Omega_r)(X_1, \dots, X_l) := \frac{1}{l_1! \dots l_r!} \sum_{\sigma} \text{sgn}(\sigma) \bar{\eta}(\Omega_1(X_{\sigma(1)}, \dots, X_{\sigma(l_1)}), \dots, \Omega_r(X_{\sigma(l-l_r+1)}, \dots, X_{\sigma(l)}))$$

where the sum is taken over all permutations σ of l elements. The coefficient in the above definition is chosen, such that if $\Omega_i = \omega_i \otimes A_i$ for some $\omega_i \in \Omega^{l_i}(M, \mathbb{K})$ and $A_i \in \Omega^0(M, \mathbb{K}^{k \times k})$ then

$$\bar{\eta}(\Omega_1, \dots, \Omega_r) = \omega_1 \wedge \dots \wedge \omega_r \bar{\eta}(A_1, \dots, A_r).$$

So this definition agrees with the one in [Kon26], but has the benefit of being coordinate independent. We will mostly work with this formula. Note that Ω_i can always be written as a sum of differential forms of the required form, where the A_i are matrices with only one non-zero entry being equal to 1. If $\Omega_1 = \dots = \Omega_r =: \Omega$ and Ω has even degree, one can also think of $\bar{\eta}(\Omega, \dots, \Omega)$ as the l -form obtained by evaluating the polynomial $\eta = \Phi_r(\bar{\eta})$ on the differential form valued entries of the matrix Ω , where multiplication is given by the wedge product (the even degree of Ω is needed for the wedge product to be commutative).

Now given a vector bundle $E \rightarrow M$, the invariance of $\bar{\eta}$ allows us to define an r -linear map

$$\bar{\eta}: \Omega^{l_1}(M, \text{End}(E)) \times \dots \times \Omega^{l_r}(M, \text{End}(E)) \rightarrow \Omega^l(M, \mathbb{K})$$

as follows: Choose any local frame of E over $U \subseteq M$. Then under this frame $\Omega_i \in \Omega^{l_i}(M, \text{End}(E))$ corresponds to a matrix valued form $\tilde{\Omega}_i \in \Omega^{l_i}(U, \mathbb{K}^{k \times k})$ and we set

$$\bar{\eta}(\Omega_1, \dots, \Omega_r) := \bar{\eta}(\tilde{\Omega}_1, \dots, \tilde{\Omega}_r)$$

on U . Since $\bar{\eta}$ is invariant, this definition does not depend on the chosen frame, hence we get a well-defined global l -form. For a polynomial $\eta = \Phi_r(\bar{\eta})$ we write

$$\eta(\Omega) := \bar{\eta}(\Omega, \dots, \Omega).$$

We can now state the central theorem.

Theorem 4.6 (Chern-Weil). *Let $E \rightarrow M$ be a rank k vector bundle with connection ∇ and curvature form Ω and let $\eta \in I_r(k)$. Then the following holds*

- (i) $\eta(\Omega)$ is closed
- (ii) If ∇' is another connection with curvature form Ω' , then $[\eta(\Omega)] = [\eta(\Omega)]$ in $H_{dR}^{2r}(M, \mathbb{K})$.

We can thus define a map

$$\chi_E^r: I_r(k) \rightarrow H_{dR}^{2r}(M, \mathbb{K})$$

by choosing any connection on E and defining $\chi_E^r(\eta) := [\eta(\Omega)]$ for the corresponding curvature form Ω . In fact if we define the algebra

$$I(k) := \bigoplus_{r=0}^{\infty} I_r(k),$$

where multiplication is defined pointwise as $(\eta_0 \cdot \eta_1)(A) := \eta_0(A)\eta_1(A)$, the maps χ_E^r define a homomorphism of commutative algebras

$$\chi_E: I(k) \rightarrow H_{dR}^{\text{even}}(M, \mathbb{K}),$$

called the **Chern-Weil homomorphism**. This homomorphism satisfies the following naturality property.

Corollary 4.7. *Let $E \rightarrow M$ be a vector bundle and $f: N \rightarrow M$ a smooth map. Then*

$$\chi_{f^*E} = f^* \chi_E.$$

Proof. From the definition of $\eta(\Omega)$ it follows directly, that $\eta(f^*\Omega) = f^*\eta(\Omega)$, such that the statement follows together with Lemma 4.1. $\square$

4.2 Proof of the Chern-Weil Theorem

We will now proof Theorem 4.6. The hard part of the proof is to show that $\eta(\Omega)$ is closed. For this we basically do the computation outlined in [Kon26] but add the missing details and fix a few mistakes. We need the following two Lemmas.

Lemma 4.8. *Let $\bar{\eta} \in I_r(k)$ and $X_1, \dots, X_r, Y \in \mathbb{K}^{k \times k}$. Then*

$$\sum_{i=1}^r \bar{\eta}(X_1, \dots, [X_i, Y], \dots, X_r) = 0,$$

where $[X_i, Y] = X_i Y - Y X_i$ is the commutator.

Proof. Let $A(t) \in \text{GL}(k, \mathbb{K})$ be a smooth curve with $A(0) = \text{id}$ and $\dot{A}(0) = Y$. Then

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} A(t)^{-1} X_i A(t) &= \frac{d}{dt} \Big|_{t=0} A(t)^{-1} X_i + \frac{d}{dt} \Big|_{t=0} X_i A(t) \\ &= -Y X_i + X_i Y \\ &= [X_i, Y], \end{aligned}$$

where we used that

$$0 = \frac{d}{dt} \Big|_{t=0} A(t) A(t)^{-1} = \frac{d}{dt} \Big|_{t=0} A(t) + \frac{d}{dt} \Big|_{t=0} A(t)^{-1} = Y + \frac{d}{dt} \Big|_{t=0} A(t)^{-1},$$

such that

$$\frac{d}{dt} \Big|_{t=0} A(t)^{-1} = -Y.$$

The invariance of $\bar{\eta}$ now yields

$$\begin{aligned} 0 &= \frac{d}{dt} \Big|_{t=0} \bar{\eta}(A(t)^{-1} X_1 A(t), \dots, A(t)^{-1} X_r A(t)) \\ &= \sum_{i=1}^r \bar{\eta}(X_1, \dots, \frac{d}{dt} \Big|_{t=0} A(t)^{-1} X_i A(t), \dots, X_r) \\ &= \sum_{i=1}^r \bar{\eta}(X_1, \dots, [X_i, Y], \dots, X_r). \end{aligned}$$

□

Before stating the second Lemma let us define the commutator of two differential forms $\Omega_1 \in \Omega^{l_1}(M, \mathbb{K}^{k \times k})$ and $\Omega_2 \in \Omega^{l_2}(M, \mathbb{K}^{k \times k})$ by

$$[\Omega_1, \Omega_2] := \Omega_1 \wedge \Omega_2 - (-1)^{l_1 l_2} \Omega_2 \wedge \Omega_1.$$

If $\Omega_i = \omega_i \otimes A_i$ for some $\omega_i \in \Omega^{l_i}(M, \mathbb{K})$ and $A_i \in \Omega^0(M, \mathbb{K}^{k \times k})$ then this can also be written as

$$[\Omega_1, \Omega_2] = \omega_1 \wedge \omega_2 \otimes [A_1, A_2],$$

such that this definition agrees with the one in [Kon26], but again has the benefit of being coordinate independent.

Lemma 4.9 (Bianchi Identity). *Let $E \rightarrow M$ be a vector bundle with a connection, let Θ be the associated connection form (with respect to some local frame) and let Ω be the curvature form. Then*

$$d\Omega + [\Theta, \Omega] = 0.$$

Proof. This is just a matter of writing out the definitions. □

We are now ready to proof the Chern-Weil Theorem.

Proof of Theorem 4.6. Since closedness is a local property we can directly use the local definition of $\eta(\Omega)$, i.e. we can assume that Ω is a matrix valued 2-form. Let $\bar{\eta} = \Phi_r^{-1}(\eta) \in S_r(k)$. In order to compute $d\eta(\Omega) = d\bar{\eta}(\Omega, \dots, \Omega)$, let us more generally show that

$$d\bar{\eta}(\Omega_1, \dots, \Omega_r) = \sum_{i=1}^r \bar{\eta}(\Omega_1, \dots, d\Omega_i, \dots, \Omega_r) \quad (9)$$

for matrix valued 2-forms $\Omega_i \in \Omega^2(M, \mathbb{K}^{k \times k})$. By linearity we may assume that all except for one entry in the matrix $\Omega_i = (\Omega_i^{jl})_{j,l=1}^k$ of 2-forms vanish, so in particular

$$\Omega_i = \omega_i \otimes E_i,$$

where E_i is a constant matrix (with only one non-zero entry being equal to 1). We then compute

$$\begin{aligned} d\bar{\eta}(\Omega_1, \dots, \Omega_r) &= d \left(\sum_{i=1}^r \omega_1 \wedge \dots \wedge \omega_r \bar{\eta}(E_1, \dots, E_r) \right) \\ &= \sum_{i=1}^r \omega_1 \wedge \dots \wedge d\omega_i \wedge \dots \wedge \omega_r \bar{\eta}(E_1, \dots, E_r) \\ &= \sum_{i=1}^r \bar{\eta}(\omega_1 \otimes E_1, \dots, d\omega_i \otimes E_i, \dots, \omega_r \otimes E_r) \\ &= \sum_{i=1}^r \bar{\eta}(\Omega_1, \dots, d\Omega_i, \dots, \Omega_r) \end{aligned}$$

Note that $dE_i = 0, d\bar{\eta}(E_1, \dots, E_r) = 0$ since both are constant. Next we claim that

$$\sum_{i=1}^r \bar{\eta}(\Omega_1, \dots, [\Theta, \Omega_i], \dots, \Omega_r) = 0, \quad (10)$$

where Θ is the connection form (or any matrix valued 1-form for that matter). Again we may assume by linearity that $\Theta = \theta \otimes E$ for some constant matrix E , such that

$$\begin{aligned} \bar{\eta}(\Omega_1, \dots, [\Theta, \Omega_i], \dots, \Omega_r) &= \bar{\eta}(\omega_1 \otimes E_1, \dots, \theta \wedge \omega_i \otimes [E, E_i], \dots, \omega_r \otimes E_r) \\ &= \theta \wedge \omega_1 \wedge \dots \wedge \omega_r \bar{\eta}(E_1, \dots, [E, E_i], \dots, E_r), \end{aligned}$$

where we used commutativity of the wedge product for 2-forms. Taking the sum over i the claim follows from Lemma 4.8. Now we plug in $\Omega = \Omega_i$ for the curvature form Ω and compute

$$\begin{aligned} d\eta(\Omega) &= d\bar{\eta}(\Omega, \dots, \Omega) \\ &\stackrel{(9)}{=} \sum_{i=1}^r \bar{\eta}(\Omega, \dots, d\Omega, \dots, \Omega) \\ &\stackrel{(10)}{=} \sum_{i=1}^r \bar{\eta}(\Omega, \dots, d\Omega + [\Theta, \Omega], \dots, \Omega) = 0, \end{aligned}$$

where we used the Bianchi identity (Lemma 4.9) in the last equation.

Next we show the independence of the chosen connection. This is basically taken from [Rah17] but with a few more details added. Let ∇^0, ∇^1 be two different connections on E with curvature forms Ω_0, Ω_1 and consider the pullback bundle π^*E under the projection $\pi: M \times \mathbb{R} \rightarrow M$. We want to define a connection ∇ on π^*E that smoothly interpolates between the connections ∇^0 and ∇^1 . To this end let $\lambda: M \times \mathbb{R} \rightarrow \mathbb{R}$ be the projection and set

$$\nabla := \lambda \pi^* \nabla^1 + (1 - \lambda) \pi^* \nabla^0.$$

It is easily seen that a convex combination of connections is again a connection (even if the coefficients are smooth functions). Now let $\iota_t: M \rightarrow M \times \mathbb{R}, p \mapsto (p, t)$ be the inclusions. We claim that $\iota_0^* \nabla = \nabla^0$ and $\iota_1^* \nabla = \nabla^1$. For this we have to check the characteristic property of pullback connections (cf. 3.7). So let $s \in \Gamma(\pi^*E)$ and $v \in T_p M$. The claim now is that

$$\nabla_v^0(\iota_0^* s) = \nabla_{(\iota_0)_* v} s.$$

Let us abbreviate $w := (\iota_0)_*v$. For the right hand side we have

$$\nabla_w s = (\pi^* \nabla^0)_w s,$$

where we used that w is a tangent vector at $(p, 0)$ and $\lambda(p, 0) = 0$. We would like to replace s with the section $\pi^* \iota_0^* s$, since then the claim just follows from the definition of the pullback connection:

$$(\pi^* \nabla^0)_w (\pi^* \iota_0^* s) = \nabla_v^0 \iota_0^* s.$$

In order to do this choose a local frame $e_1, \dots, e_k$ of E and write $s = s^i \pi^* e_i$ for smooth functions $s^i: M \times \mathbb{R} \rightarrow \mathbb{K}$. Then $\pi^* \iota_0^* s = (s^i \circ \iota_0 \circ \pi) \pi^* e_i$ and thus

$$\begin{aligned} (\pi^* \nabla^0)_w (\pi^* \iota_0^* s) &= w \cdot (s^i \circ \iota_0 \circ \pi) e_i(p) + s^i(p, 0) \nabla_v^0 e_i \\ &= w \cdot s^i e_i(p) + s^i(p, 0) \nabla_v^0 e_i \\ &= (\pi^* \nabla^0)_w s. \end{aligned}$$

The same argument holds for ι_1 and ∇^1 . In fact $\iota_t^* \nabla = \nabla^t$, where $\nabla^t = t \nabla^1 + (1-t) \nabla^0$. Now let Ω be the curvature form of ∇ , then by Lemma 4.1

$$\iota_0^* \Omega = \Omega_0, \quad \iota_1^* \Omega = \Omega_1.$$

Thus $\iota_0^* \eta(\Omega) = \eta(\Omega_0)$ and $\iota_1^* \eta(\Omega) = \eta(\Omega_1)$. Since the maps ι_0 and ι_1 are homotopic they induce the same map on cohomology, such that the statement follows. $\square$

4.3 Chern and Pontryagin Classes

Let $E \rightarrow M$ be a complex vector bundle of rank d . For any matrix $A \in \mathbb{C}^{d \times d}$ we can consider the characteristic polynomial in t :

$$\det \left(\text{id} - \frac{t}{2\pi i} A \right) = \sum_{k=0}^d f_k(A) t^k,$$

where $f_k: \mathbb{C}^{d \times d} \rightarrow \mathbb{C}$ are the coefficients of this polynomial and depend on A . Using the Leibniz formula for the determinant one can see that f_k is a k -homogeneous polynomial and it moreover is invariant since the characteristic polynomial is. Furthermore it follows from the Leibniz formula that

$$\begin{aligned} f_0(A) &= 1, \\ f_1(A) &= \text{tr} \left(-\frac{1}{2\pi i} A \right), \\ f_d(A) &= \det \left(-\frac{1}{2\pi i} A \right). \end{aligned}$$

Definition 4.10. The k -th Chern class of E is defined as

$$c_k(E) := \chi_E(f_k) \in H_{\text{dR}}^{2k}(M; \mathbb{C}).$$

If we instead consider a real vector bundle $E \rightarrow M$ we can similarly define k -homogeneous polynomials $g_k: \mathbb{R}^{d \times d} \rightarrow \mathbb{R}$ by

$$\det \left(\text{id} - \frac{t}{2\pi} A \right) = \sum_{k=0}^d g_k(A) t^k,$$

and define:

Definition 4.11. The k -th Pontryagin class of E is defined as

$$p_k(E) := \chi_E(g_{2k}) \in H_{\text{dR}}^{4k}(M; \mathbb{R}).$$

Of course there is a connection between these two definitions:

Lemma 4.12. *Let $E \rightarrow M$ be a real vector bundle. Then*

$$p_k(E) = (-1)^k c_{2k}(E \otimes \mathbb{C}),$$

where $E \otimes \mathbb{C}$ is the complexified bundle and the equality is understood by considering a real differential as the real part of a complex differential form, that is $H_{\text{dR}}^{4k}(M; \mathbb{R}) \subseteq H_{\text{dR}}^{4k}(M; \mathbb{C})$.

Proof. Choose any connection ∇ on E and let $\nabla_{\mathbb{C}}$ be the induced connection on the complexified bundle $E \otimes \mathbb{C}$. Then by definition any frame of E also defines a frame of $E \otimes \mathbb{C}$ and the corresponding connection 1-forms are equal. Thus the curvature form Ω of E agrees with the curvature form $\Omega_{\mathbb{C}}$ of $E \otimes \mathbb{C}$. From the definition of the polynomials f_k and g_k it is moreover clear that for any real matrix $A \in \mathbb{R}^{d \times d}$ we have

$$f_k(A) = (-i)^k g_k(A).$$

Thus

$$c_{2k}(E) = [f_{2k}(\Omega_{\mathbb{C}})] = [f_{2k}(\Omega)] = (-i)^{2k} [g_{2k}(\Omega)] = (-1)^k p_k(E).$$

□

Now assume we are given a real oriented vector bundle $E \rightarrow M$ of rank $2n$. Choose a metric on E together with a compatible connection. A simple computation shows that the connection 1-form Θ with respect to any oriented orthonormal frame is skew-symmetric. Thus

$$\begin{aligned} (\Theta \wedge \Theta)^T(X, Y) &= (\Theta(X)\Theta(Y) - \Theta(Y)\Theta(X))^T \\ &= (-\Theta(Y))(-\Theta(X)) - (-\Theta(X))(-\Theta(Y)) \\ &= -(\Theta \wedge \Theta)(X, Y). \end{aligned}$$

So the curvature form Ω is also skew symmetric:

$$\Omega^T = (d\Theta)^T + (\Theta \wedge \Theta)^T = -d\Theta - \Theta \wedge \Theta = -\Omega.$$

Recall that the Pfaffian is a n -homogeneous polynomial

$$\text{Pf}: \text{Skew}_{2n} \rightarrow \mathbb{R}$$

on the space of skew symmetric $2n \times 2n$ matrices and satisfies

$$\text{Pf}(B^T A B) = \det(B) \text{Pf}(A)$$

for $A \in \text{Skew}_{2n}$ and $B \in \mathbb{R}^{n \times n}$ an arbitrary matrix. Thus we can define the $2n$ -form $\text{Pf}(\Omega)$ on the neighbourhood where the oriented orthonormal frame was defined. Under a change of oriented orthonormal frame the curvature form Ω transforms as $S\Omega S^{-1}$ for a special orthogonal matrix $S \in \text{SO}(2n)$. Since $S^{-1} = S^T$ and $\det(S) = 1$ we conclude that $\text{Pf}(\Omega)$ does not depend on the chosen frame and thus defines a global object. Just as in the proof of Theorem 4.6 we see that $\text{Pf}(\Omega)$ is closed (only the statement of Lemma 4.8 changes, where instead of arbitrary X_i and Y we only allow for skew symmetric matrices. Note that the tangent space of $\text{SO}(2n)$ is precisely the space of skew symmetric matrices, such that we can realize any such Y as the velocity vector of a curve through id). Moreover the cohomology class does not depend on the chosen connection and metric. Again the proof is similar to Theorem 4.6. Thus we can define

Definition 4.13. The **Euler class** of E is defined as

$$e(E) := \left[\text{Pf} \left(\frac{1}{2\pi} \Omega \right) \right] \in H_{\text{dR}}^{2n}(M; \mathbb{R}).$$

It is no coincidence that the above construction of the Euler class works and is just an example of the following more general construction. Instead of considering homogeneous polynomials on $\mathbb{K}^{k \times k}$ which are invariant under conjugation with elements in $\text{GL}(k, \mathbb{K})$ we can more generally consider polynomials $\eta: \mathfrak{g} \rightarrow \mathbb{K}$ defined on the Lie-Algebra $\mathfrak{g}$ of some matrix Lie-group $G \subseteq \text{GL}(k, \mathbb{K})$, which are invariant under conjugation with elements in G . Suppose we are then given a vector bundle $E \rightarrow M$ with structure group G and some connection, such that the corresponding (locally defined) curvature form Ω takes values in $\mathfrak{g}$. Under a change of local frame this curvature form transforms as $S\Omega S^{-1}$ for a matrix $S \in G$, thus $\eta(\Omega)$ is a well defined global object and the proofs above are easily generalized to show that $\eta(\Omega)$ is closed and does depend on the chosen connection (as long as the connection yields local curvature forms with values in $\mathfrak{g}$). For this more general approach to be useful we need to guarantee the existence of such connections, which is the statement of the following proposition.

Proposition 4.14. *Let $E \rightarrow M$ be a vector bundle with structure group $G \subseteq \mathrm{GL}(k, \mathbb{K})$. Then there exists a connection on E , such that the curvature form with respect to any local frame ³takes values in $\mathfrak{g}$.*

Proof. Let (e_i^α) be local frames defined on some neighbourhoods $U_\alpha \subseteq M$ which cover M . On $E|_{U_\alpha}$ define a connection by $\nabla^\alpha e_i = 0$. In other words for a local section $s = s^i e_i^\alpha$ we set $\nabla^\alpha s := ds^i \otimes e_i^\alpha$. Now let (ψ_α) be a partition of unity subordinate to the cover (U_α) and define a connection ∇ on E by

$$\nabla s := \sum_\alpha \psi_\alpha \nabla^\alpha s$$

for any section $s \in \Gamma(E)$. We claim that the connection one forms of ∇ already take values in $\mathfrak{g}$. To this end let (e_i) be any other frame defined on $U \subseteq M$ and let $A_\alpha = (a_{ij}^\alpha) \in C^\infty(U \cap U_\alpha, G)$ be the change of basis matrix from (e_i) to (e_i^α) , that is

$$e_j = \sum_i a_{ij}^\alpha e_i^\alpha.$$

Moreover we write $A_\alpha^{-1} = (a_\alpha^{ij})$ for the inverse, that is

$$e_j^\alpha = \sum_i a_\alpha^{ij} e_i.$$

We now compute

$$\nabla^\alpha e_j = \sum_i da_{ij}^\alpha \otimes e_i^\alpha = \sum_{i,k} a_\alpha^{ki} da_{ij}^\alpha \otimes e_k = \sum_{k,i} a_\alpha^{ik} da_{kj}^\alpha \otimes e_i$$

In the last equality we only exchanged the roles of k and i . So

$$\nabla e_j = \sum_{\alpha,k,i} \psi_\alpha a_\alpha^{ik} da_{kj}^\alpha \otimes e_i,$$

such that for some vector field X the connection one form is given by

$$\Theta(X) = \sum_\alpha \psi_\alpha \left(\sum_k a_\alpha^{ik} da_{ki}^\alpha(X) \right)_{ij} = \sum_\alpha \psi_\alpha A_\alpha^{-1} dA_\alpha(X)$$

Note that for some point $p \in M$, where both frames (e_i) and (e_i^α) are defined we have

$$d_p A_\alpha(X_p) \in T_{A_\alpha(p)} G,$$

such that $A_\alpha^{-1}(p) d_p A_\alpha(X_p) = d_p(A_\alpha^{-1}(p) A_\alpha)(X_p) \in T_e G = \mathfrak{g}$. Thus $\Theta(X) \in \mathfrak{g}$. Next we show that the corresponding curvature form

$$\Omega = \Theta \wedge \Theta + d\Theta$$

also takes values in $\mathfrak{g}$. Firstly note that for vector fields X, Y we have

$$(\Theta \wedge \Theta)(X, Y) = \Theta(X)\Theta(Y) - \Theta(Y)\Theta(X) = [\Theta(X), \Theta(Y)]$$

and it follows from general Lie theory of matrix groups, that the corresponding Lie algebra is closed under taking the commutator. For the second term we apply Corollary 2.7 component wise to get

$$d\Theta(X, Y) = X \cdot \Theta(Y) - Y \cdot \Theta(X) - \Theta([X, Y]).$$

The first two terms are just directional derivatives of smooth functions with values in the vector space $\mathfrak{g}$ and thus take again values in $\mathfrak{g}$. The statement follows. $\square$

³That is any local frame determined by a local trivialisation from the maximal trivialising atlas making $E \rightarrow M$ into a vector bundle with structure group G . Compare Definition 7.1.

5 Algebraic Topology

5.1 General Statements about Homology

In the following let $(H_\bullet, \partial_\bullet)$ be any homology theory.

Lemma 5.1. *Let $A \subseteq X$ be a retract. Then $H_\bullet(X) \cong H_\bullet(X, A) \oplus H_\bullet(A)$.*

Proof. Let $r: X \rightarrow A$ be a retraction and $i: A \rightarrow X$ the inclusion. Then $H_\bullet(r) \circ H_\bullet(i) = \text{id}_{H_\bullet(A)}$, in particular $H_\bullet(i)$ is injective. Hence the long exact sequence for the pair (X, A) breaks up into short split exact sequences

$$0 \rightarrow H_n(A) \xrightarrow{H_n(r)} H_n(X) \rightarrow H_n(X, A) \rightarrow 0$$

and the splitting lemma yields the desired isomorphism. $\square$

Lemma 5.2. *Let $f: (X, A) \rightarrow (Y, B)$ be a continuous map of pairs, such that $f: X \rightarrow Y$ and $f: A \rightarrow B$ are homotopy equivalences. Then f induces an isomorphism*

$$H_\bullet(f): H_\bullet(X, A) \rightarrow H_\bullet(Y, B).$$

Remark 5.3. Note that $f: (X, A) \rightarrow (Y, B)$ is not necessarily a homotopy equivalence of pairs. See chapter 3 in [Ste25] for a counterexample.

Proof. Consider the map between the long exact sequences of the pairs (X, A) and (Y, B) induced by f :

$$\begin{array}{ccccccccc} \cdots & \rightarrow & H_n(A) & \rightarrow & H_n(X) & \rightarrow & H_n(X, A) & \rightarrow & H_{n-1}(A) & \rightarrow & H_{n-1}(X) & \rightarrow & \cdots \\ & & \cong \downarrow H_n(f) & & \cong \downarrow H_n(f) & & \downarrow H_n(f) & & \cong \downarrow H_{n-1}(f) & & \cong \downarrow H_{n-1}(f) & & \\ \cdots & \rightarrow & H_n(B) & \rightarrow & H_n(Y) & \rightarrow & H_n(Y, B) & \rightarrow & H_{n-1}(B) & \rightarrow & H_{n-1}(Y) & \rightarrow & \cdots \end{array}$$

The outer maps are isomorphisms by homotopy invariance and so the five lemma implies that the middle map is an isomorphism as well. $\square$

Lemma 5.4. *Let $A \subseteq X$ be a closed neighbourhood-deformation retract. Then the projection $X \rightarrow X/A$ induces a natural isomorphism*

$$H_\bullet(X, A) \rightarrow H_\bullet(X/A, *).$$

Proof. Applying excision in the push-out

$$\begin{array}{ccc} A & \longrightarrow & X \\ \downarrow & & \downarrow \\ * & \longrightarrow & X/A \end{array}$$

shows that the projection induces an isomorphism. Furthermore, any map $(X, A) \rightarrow (Y, B)$ descends to a map $X/A \rightarrow Y/B$ and naturality then just follows from functoriality. $\square$

5.2 Seifert-Van Kampen for Pushouts

Theorem 5.5 (Seifert-Van Kampen for Pushouts). *Let*

$$\begin{array}{ccc} X_0 & \xrightarrow{i_2} & X_2 \\ i_1 \downarrow & & \downarrow j_2 \\ X_1 & \xrightarrow{j_1} & X \end{array}$$

be a pushout of topological spaces, where X_0 is path connected and i_1 and i_2 are inclusions of neighbourhood deformation retracts. Then for any base point $x \in X_0$ the following diagram is a pushout

$$\begin{array}{ccc} \pi_1(X_0, x) & \xrightarrow{(i_2)_*} & \pi_1(X_2, i_2(x)) \\ (i_1)_* \downarrow & & \downarrow (j_2)_* \\ \pi_1(X_1, i_1(x)) & \xrightarrow{(j_1)_*} & \pi_1(X, j(x)) \end{array}$$

where $j = j_1 \circ i_1 = j_2 \circ i_2$.

Proof. Under these assumptions we can find open neighbourhoods $U_1, U_2 \subseteq X$ of $j_1(X_1), j_2(X_2)$ respectively, such that $j_i: X_i \rightarrow U_i$ are deformation retractions. From the construction of these open neighbourhoods it is clear that $j: X_0 \rightarrow U_1 \cap U_2$ is also a deformation retraction. From the ordinary Seifert-Van Kampen Theorem it follows that

$$\begin{array}{ccc} \pi_1(U_1 \cap U_2, j(x)) & \longrightarrow & \pi_1(U_2, j(x)) \\ \downarrow & & \downarrow \\ \pi_1(U_1, j(x)) & \longrightarrow & \pi_1(X, j(x)) \end{array}$$

is a pushout, where all maps are induced by inclusions. The theorem now follows from the commutativity of the following diagram

$$\begin{array}{ccccc} & & \pi_1(X_2, i_2(x)) & \xrightarrow{\cong} & \pi_1(U_2, j(x)) \\ & \nearrow & \downarrow & & \nearrow \\ \pi_1(X_0, x) & \xrightarrow{\cong} & \pi_1(U_1 \cap U_2, j(x)) & & \downarrow \\ & \searrow & \downarrow & & \downarrow \\ & & \pi_1(X, j(x)) & \xrightarrow{\cong} & \pi_1(X, j(x)) \\ & \nearrow & \downarrow & & \nearrow \\ \pi_1(X_1, i_1(x)) & \xrightarrow{\cong} & \pi_1(U_1, j(x)) & & \end{array}$$

where the right side square is the pushout from above, the left side square is the supposed pushout and the horizontal maps in between are induced by the deformation retractions and in particular isomorphisms. $\square$

5.3 CW-Complexes

Proposition 5.6. *Let (X, A) be a relative CW-complex. If A is Hausdorff (resp. normal) then X is Hausdorff (resp. normal) as well. In particular every absolute CW-complex is Hausdorff and normal.*

Proof. We show the statement for the case that A is normal. The Hausdorff case follows completely analogously by replacing the sets to be separated by point sets. So let $B, C \subseteq X$ be closed disjoint sets. We will show the following claim inductively:

Claim. There exist open disjoint sets $U_n, V_n \subseteq X_n$, such that $B_n := B \cap X_n \subseteq U_n$ and $C_n := C \cap X_n \subseteq V_n$ as well as $U_{n-1} \subseteq U_n, V_{n-1} \subseteq V_n$.

The case $n = -1$ follows from the assumption on A . For the induction step choose a push-out

$$\begin{array}{ccc} \coprod_{i \in I} S^{n-1} & \xrightarrow{(q_i)} & X_{n-1} \\ \downarrow & & \downarrow \\ \coprod_{i \in I} D^n & \xrightarrow{(Q_i)} & X_n \end{array}$$

and set $U'_{n-1} := (q_i)_{i \in I}^{-1}(U_{n-1})$, $V'_{n-1} := (q_i)_{i \in I}^{-1}(V_{n-1})$. By Lemma 5.7 shown below we can find open neighbourhoods $U'_n, V'_n \subseteq \coprod_I D^n$ with the following properties:

- (i) $(Q_i)_I^{-1}(B_n) \subseteq U'_n$ and $(Q_i)_I^{-1}(C_n) \subseteq V'_n$
- (ii) $U'_n \cap V'_n = \emptyset$
- (iii) $U'_{n-1} = U'_n \cap \coprod_I S^{n-1}$ and $V'_{n-1} = V'_n \cap \coprod_I S^{n-1}$

Then define

$$U_n := U_{n-1} \cup (Q_i)_I(U'_n), \quad V_n := V_{n-1} \cup (Q_i)_I(V'_n)$$

By (iii) $U_n \cap X_{n-1} = U_{n-1}$ and $(Q_i)_I^{-1}(U_n) = U'_n$, so $U_n \subseteq X_n$ is open. The same holds for V_n . Furthermore, $B_n \subseteq U_n$, $C_n \subseteq V_n$ by (i) and $U_n \cap V_n = \emptyset$ by (ii). This shows the claim. Now consider

$$U := \bigcup_n U_n, \quad V := \bigcup_n V_n.$$

By construction $U \cap X_n \subseteq$ is open for all $n \geq -1$ and hence $U \subseteq X$ is open by definition of the colimit topology. The same holds for V . Moreover, $B \subseteq U, C \subseteq V$ and $U \cap V = \emptyset$. $\square$

It remains to proof the lemma mentioned in the proof of the above proposition.

Lemma 5.7. *Let $B, C \subseteq D^n$ be closed disjoint sets and $U', V' \subseteq S^{n-1}$ open disjoint neighbourhoods of $B \cap S^{n-1}, C \cap S^{n-1}$, respectively. Then there exist open sets $U, V \subseteq D^n$ with the following properties:*

- (i) $B \subseteq U, C \subseteq V$
- (ii) $U \cap V = \emptyset$
- (iii) $U' = U \cap S^{n-1}, V' = V \cap S^{n-1}$

Proof. Consider the open disjoint sets

$$U'' := \{rx \mid x \in U', 0 < r \leq 1\}$$

and

$$V'' := \{rx \mid x \in V', 0 < r \leq 1\}.$$

Moreover, since B and $\overline{V'}$ are disjoint and closed we can find an open neighbourhood W_B of B , such that

$$\overline{W_B} \cap \overline{V'} = \emptyset.$$

Similialry we can find an open neighbourhood W_C of C with

$$\overline{W_C} \cap \overline{U'} = \emptyset.$$

After potentially shrinking W_B and W_C we may also assume, that they are disjoint. Now set

$$U := (U'' - \overline{W_C}) \cup (W_B - S^{n-1})$$

and

$$V := (V'' - \overline{W_B}) \cup (W_C - S^{n-1}).$$

It is then easy to verify, that these sets satisfy the statement. $\square$

5.4 Remarks about Urs' lecture on Algebraic Topology

In this section we present an alternative proof on how to extend the fact that de-Rham cohomology and singular cohomology of open subsets of $\mathbb{R}^n$ are isomorphic to manifolds with or without boundary, while at the same time formulating the statement in more category theoretical terms.

Lemma 5.8. *The map*

$$\begin{aligned}\Gamma: \Omega^k(M) &\rightarrow C_k^\infty(M)^* \\ \omega &\mapsto (\sigma \mapsto \int_\sigma \omega := \int_{\Delta^k} \sigma^* \omega)\end{aligned}$$

defines a natural map of cochain complexes

$$(\Omega^\bullet(M), d) \rightarrow (C_k^\infty(M)^*, \partial^*),$$

i.e. a natural transformation of functors from the category of smooth manifolds to the category of cochain complexes over $\mathbb{R}$.

Proof. For smooth manifolds M, N and a smooth map $f: M \rightarrow N$ we have to show the commutativity of the following diagrams:

$$\begin{array}{ccc} \Omega^k(M) & \xrightarrow{\Gamma} & C_k^\infty(M)^* \\ \downarrow d & & \downarrow \partial^* \\ \Omega^{k+1}(M) & \xrightarrow{\Gamma} & C_{k+1}^\infty(M)^* \end{array} \quad \begin{array}{ccc} \Omega^k(N) & \xrightarrow{\Gamma} & C_k^\infty(N)^* \\ \downarrow f^* & & \downarrow f^* \\ \Omega^k(M) & \xrightarrow{\Gamma} & C_k^\infty(M)^* \end{array}$$

For the left diagram we compute:

$$(\Gamma \circ d)(\omega) = \left(\sigma \mapsto \int_\sigma d\omega \right) = \left(\sigma \mapsto \int_{\partial\sigma} \omega \right) = (\partial^* \circ \Gamma)(\omega)$$

where we used Stokes' Theorem for singular chains in the second equality. Moreover, we compute for the right diagram:

$$\begin{aligned}(\Gamma \circ f^*)(\omega) &= \left(\sigma \mapsto \int_\sigma f^* \omega \right) \\ (f^* \circ \Gamma)(\omega) &= \left(\sigma \mapsto \int_{f \circ \sigma} \omega \right)\end{aligned}$$

But by definition:

$$\int_\sigma f^* \omega = \int_{\Delta^k} \sigma^* f^* \omega = \int_{\Delta^k} (f \circ \sigma)^* \omega = \int_{f \circ \sigma} \omega$$

This completes the proof. □

Lemma 5.9. *The inclusion $C_k^\infty(M) \rightarrow C_k(M)$ induces a natural map of cochain complexes*

$$(C_k^\infty(M)^*, \partial^*) \rightarrow (C_k(M)^*, \partial^*)$$

Proof. Obvious. □

In the lecture we have shown that the cochain maps from the above two lemmas induce isomorphisms in homology at least for open subsets of $\mathbb{R}^n$. A short proof of how this generalizes to all manifolds with or without boundary is given in the following theorem (which notably uses facts that we have not established in the lecture).

Theorem 5.10. *The cochain maps from Lemma 5.8 and 5.9 induce isomorphisms in homology.*

Proof. We first show the statement if $\partial M = \emptyset$. Begin by choosing an embedding

$$\iota: M \hookrightarrow U$$

into a tubular neighbourhood $U \subseteq \mathbb{R}^n$. In particular, U is open and ι a homotopy equivalence. Then by Lemma 5.8 the following diagram commutes:

$$\begin{array}{ccc} H_{dR}^k(U) & \xrightarrow{\Gamma} & H_{\infty}^k(U) \\ \downarrow \iota^* & & \downarrow \iota^* \\ H_{dR}^k(M) & \xrightarrow{\Gamma} & H_{\infty}^k(M) \end{array}$$

The vertical maps are isomorphisms due to homotopy invariance and in the lecture we have shown that the top horizontal map is an isomorphism as well. The claim follows.

If $\partial M \neq \emptyset$ we can reduce this to the case $\partial M = \emptyset$ by considering the homotopy equivalence

$$\text{int}(M) \hookrightarrow M$$

and using a completely analogous argument to above. The same argument shows the statement for the map from Lemma 5.9. $\square$

Composing the two maps we get a natural isomorphism

$$H_{dR}^k(M) \rightarrow H^k(M).$$

Under the identification $H^k(M) \cong H_k^*(M)$ (which is also natural, see Lemma 5.11 below) this isomorphism is given by

$$[\omega] \mapsto \left([\eta] \mapsto \int_{\tilde{\eta}} \omega \right),$$

where $\tilde{\eta}$ is a smooth representative of the class $[\eta]$.

Lemma 5.11. *Let $(C_{\bullet}, d_{\bullet})$ be a chain complex over $\mathbb{R}$ (or any other field). Then*

$$\begin{aligned} D: H^k(C_{\bullet}^*, d_{\bullet}^*) &\rightarrow H_k^*(C_{\bullet}, d_{\bullet}) \\ [\phi] &\mapsto ([c] \mapsto \phi(c)) \end{aligned}$$

defines an isomorphism, which is natural for maps between chain complexes over $\mathbb{R}$.

Proof. First we show well-definedness, so let $\phi' = \phi + d^*\psi$ for some $\psi \in C_{k-1}^*$. But then

$$\phi'(c) - \phi(c) = d^*\psi(c) = \psi(dc) = 0$$

since c is closed. Moreover, if $c' = c + da$ for some $a \in C_{k+1}$ we get

$$\phi(c' - c) = \phi(da) = d^*\phi(a) = 0$$

since ϕ is closed. Next we show injectivity, i.e. for closed $\phi \in C_k^*$ with $\phi(c) = 0$ for all closed $c \in C_k$ we have to construct $\psi \in C_{k-1}^*$ with $d^*\psi = \phi$. For $a = dc \in C_{k-1}$ we define ψ by

$$\psi(a) := \phi(c)$$

which is well defined since for $a = dc'$ we have $d(c - c') = 0$ such that

$$\phi(c - c') = 0.$$

Now extend ψ arbitrarily to a linear map $C_{k-1} \rightarrow \mathbb{R}$ (which is possible since C_{k-1} is a vector space.) To show surjectivity let $\tau \in H_k^*(C_{\bullet}, d_{\bullet})$. We have to construct a closed $\phi \in C_k^*$ with

$$\phi(c) = \psi([c])$$

for all closed $c \in C_k$. To this end write

$$C_k = \ker d_k \oplus V$$

for some subspace $V \subseteq C_k$ and define ϕ by

$$\begin{aligned}\phi(c) &:= \psi([c]), \quad c \in \ker d_k \\ \phi|_V &:= 0\end{aligned}$$

By construction $d^*\phi = 0$ and $D([\phi]) = \psi$. Finally, we will show naturality. Let

$$f_\bullet: (C_\bullet, d_\bullet) \rightarrow (B_\bullet, b_\bullet)$$

be a chain map. We have to show commutativity of the following diagram:

$$\begin{array}{ccc} H^k(C_\bullet^*, d_\bullet^*) & \xrightarrow{D} & H_k^*(C_\bullet, d_\bullet) \\ \uparrow H^k(f_\bullet) & & \uparrow H_k^*(f_\bullet) \\ H^k(B_\bullet^*, b_\bullet^*) & \xrightarrow{D} & H_k^*(B_\bullet, b_\bullet) \end{array}$$

For $\phi \in H^k(B_\bullet^*, b_\bullet^*)$ we compute:

$$\begin{aligned}(H_k^*(f_\bullet) \circ D)([\phi]) &= (H_k(f_\bullet))^*([b] \mapsto \phi(b)) \\ &= ([c] \mapsto \phi(f_k c))\end{aligned}$$

as well as

$$\begin{aligned}(D \circ H^k(f_\bullet))([\phi]) &= D([\phi \circ f_k]) \\ &= ([c] \mapsto \phi(f_k c)).\end{aligned}$$

□

5.5 Local Computation of the Degree

In this section we proof the following theorem.

Theorem 5.12. *Let $f: S^n \rightarrow S^n$ be smooth around a point $x \in S^n$, $y = f(x)$ and $T_x f: T_x S^n \rightarrow T_y S^n$ an isomorphism. Then for the local degree of f at x we have*

$$\deg_x(f) = +1$$

if $T_x f$ is orientation preserving and

$$\deg_x(f) = -1$$

if $T_x f$ is orientation reversing.

Proof. First assume $T_x f$ is orientation preserving. We have to show that for an open neighbourhood $U \subseteq S^n$ of x the map

$$H_n(S^n) \xrightarrow{\cong} H_n(S^n, S^n - y_1) \xleftarrow{\cong} H_n(U, U - x) \xrightarrow[\cong]{H_n(f|_U)} H_n(S^n, S^n - y_1) \xleftarrow{\cong} H_n(S^n)$$

is the identity (where all other maps are induced by inclusions). This follows immediately from the following claim.

Claim. For orientation preserving embeddings $g_i: U \rightarrow S^n$ with $g_i(x) = y_i$, $i = 1, 2$ the map

$$H_n(S^n) \xrightarrow{\cong} H_n(S^n, S^n - y_1) \xleftarrow[\cong]{H_n(g_1)} H_n(U, U - x) \xrightarrow[\cong]{H_n(g_2)} H_n(S^n, S^n - y_1) \xleftarrow{\cong} H_n(S^n) \quad (\star)$$

is the identity

Firstly we may assume, that $y_1 = y_2$, otherwise replace g_1 with $R \circ g_1$, where $R: S^n \rightarrow S^n$ is a rotation with $R(y_1) = y_2$. Then the map $(\star)$ is unchanged as seen by the commutativity of the following diagramm

$$\begin{array}{ccccc}
H_n(S^n) & \xrightarrow{\cong} & H_n(S^n, S^n - y_1) & & \\
\downarrow H_n(R) = id & & \downarrow H_n(R) & \swarrow H_n(g_1) & \\
H_n(S^n) & \xrightarrow{\cong} & H_n(S^n, S^n - y_2) & \swarrow H_n(R \circ g_1) & \\
& & & \swarrow H_n(U, U - x) &
\end{array}$$

Note that $R \simeq id_{S^n}$ since $SO(n)$ is path-connected. Moreover we may assume $U \cong \mathbb{R}^n$, otherwise shrink U to some $U' \subseteq U$ and excision yields $H_n(U', U' - x) \xrightarrow{\cong} H_n(U, U - x)$. So if we consider g_1 and g_2 in suitable charts we get orientation preserving embeddings

$$\tilde{g}_1, \tilde{g}_2: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

with $\tilde{g}_1(0) = 0 = \tilde{g}_2(0)$. So by Lemma 8.1 $\tilde{g}_1$ and $\tilde{g}_2$ are isotopic rel $\{0\}$. In particular we get a homotopy

$$H: U \times I \rightarrow S^n$$

from g_1 to g_2 with $H((U - x) \times I) \subseteq S^n - y_1$, i.e. a homotopy of *pairs*

$$g_1, g_2: (U, U - x) \rightarrow (S^n, S^n - y_1)$$

and thus

$$H_n(g_1) = H_n(g_2): H_n(U, U - x) \rightarrow H_n(S^n, S^n - y_1)$$

which proofs the claim. It remains to consider the case where $T_x f$ is orientation reversing. For this consider the following commutative diagramm

$$\begin{array}{ccccccc}
H_n(S^n) & \xrightarrow{\cong} & H_n(S^n, S^n - x) & \xleftarrow{\cong} & H_n(U, U - x) & \xrightarrow[\cong]{H_n(f|_U)} & H_n(S^n, S^n - y) & \xleftarrow{\cong} & H_n(S^n) \\
& & & & \searrow H_n(r \circ f|_U) & & \downarrow \cong H_n(r) & & \downarrow \cong H_n(r) \\
& & & & & & H_n(S^n, S^n - r(y)) & \xleftarrow[\cong]{} & H_n(S^n)
\end{array}$$

where $r: S^n \rightarrow S^n$ is the reflection along the equator. Then $T_x(r \circ f)$ is orientation preserving and the statement follows from what has been shown already and the fact that $\deg(r) = -1$. $\square$

6 Point Set Topology

Lemma 6.1. *Let $\pi: X \rightarrow Y$ be a quotient map and $A \subseteq X$ with $\pi^{-1}(\pi(A)) = A$. If $A \subseteq X$ is either closed or open then the restriction*

$$\pi|_A: A \rightarrow \pi(A)$$

is also a quotient map.

Proof. Let $p := \pi|_A: A \rightarrow \pi(A)$ be the restriction and A be open. Given $V \subseteq \pi(A)$ such that $U := p^{-1}(V) \subseteq A$ is open we get from our assumptions on A that U is also open in X and that $U = \pi^{-1}(V)$, hence V is also open in Y and thus V is open in $\pi(A)$. The same argument shows the statement if A is closed, since we can equivalently describe the quotient topology in terms of closed sets. $\square$

Lemma 6.2. *Let $(X_i)_{i \in I}$ be a family of connected topological spaces. Then the product $\prod_{i \in I} X_i$ is again connected.*

Proof. We first show that the statement holds if I is finite. So let X_1 and X_2 be connected and let

$$f: X_1 \times X_2 \rightarrow \{0, 1\}$$

be continuous. Let $(p_1, p_2), (p'_1, p'_2) \in X_1 \times X_2$. Since the subspaces $X_1 \times \{p_2\}$ and $\{p'_1\} \times X_2$ are connected (they are images of inclusion maps $X_i \rightarrow X_1 \times X_2$) and both contain the point (p'_1, p_2) it follows that

$$Z := X_1 \times \{p_2\} \cup \{p'_1\} \times X_2$$

is connected and thus f is constant on Z . But the points (p_1, p_2) and (p'_1, p'_2) were chosen arbitrarily, thus f is constant. It follows inductively, that the lemma holds for all finite I . Now let I be arbitrary and again

$$f: \prod_{i \in I} X_i \rightarrow \{0, 1\}$$

continuous and $(p_i), (p'_i) \in \prod_{i \in I} X_i$. Consider the set

$$D := \left\{ \prod_{i \in I} U_i \mid p'_i \in U_i \text{ and } U_i = X_i \text{ for all but finitely many } i \in I \right\}$$

of open product neighbourhoods of (p'_i) . Together with the relation

$$\prod_{i \in I} U_i \supseteq \prod_{i \in I} V_i \Leftrightarrow \prod_{i \in I} U_i \subseteq \prod_{i \in I} V_i$$

D has the structure of a directed set. For $\alpha = \prod_{i \in I} U_i \in D$ define

$$x_i^\alpha := \begin{cases} p_i & , \text{ if } U_i = X_i \\ p'_i & , \text{ otherwise} \end{cases}$$

By construction the net $(x_i^\alpha)_\alpha$ converges to (p'_i) . It moreover follows from the case where I is finite, that

$$f((x_i^\alpha)_i) = f((p_i)_i)$$

for all $\alpha \in D$, since for $\alpha = \prod_{i \in I} U_i \in D$ both points lie in the connected subspace

$$A := \prod_{i \in I} A_i,$$

where $A_i = \{p_i\}$ if $U_i = X_i$ and $A_i = X_i$ for the finitely many exceptions. Thus

$$f((p'_i)_i) = \lim_{\alpha \in D} f((x_i^\alpha)_i) = f((p_i)_i).$$

$\square$

Lemma 6.3. *Let K be a compact topological space, (Y, d) a metric space and $f: K \rightarrow Y$ continuous. Then for any open neighbourhood $U \subseteq K \times Y$ of the graph $\Gamma_f \subseteq K \times Y$ of f , there exists $\varepsilon > 0$, such that the ε -tube*

$$\{(x, y) \in K \times Y \mid d(f(x), y) < \varepsilon\}$$

is contained in U .

7 Fiber Bundles

7.1 Construction of Fiber Bundles

Definition 7.1 (Fiber Bundle). Let G be a topological group acting effectively on a space F . A fiber bundle E over B with fiber F and structure group G consists of a surjective continuous map $p: E \rightarrow B$ and a collection of maps

$$(\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha))_\alpha,$$

called a trivialising atlas⁴, for open sets $U_\alpha \subseteq B$, such that

- (i) The maps ϕ_α are homeomorphisms and the following diagrams commute:

$$\begin{array}{ccc} U_\alpha \times F & \xrightarrow{\phi_\alpha} & p^{-1}(U_\alpha) \\ & \searrow \pi \quad \swarrow p & \\ & U_\alpha & \end{array}$$

where π is the usual projection onto the first factor.

- (ii) For any two maps ϕ_α, ϕ_β there exists a continuous map

$$\tau_{\alpha,\beta}: U_\alpha \cap U_\beta \rightarrow G,$$

such that

$$\phi_\alpha^{-1} \circ \phi_\beta: (U_\alpha \cap U_\beta) \times F \rightarrow (U_\alpha \cap U_\beta) \times F$$

is given by $(b, f) \mapsto (b, \tau_{\alpha,\beta}(b) \cdot f)$.

- (iii) $(U_\alpha)_\alpha$ is an open cover of B .

- (iv) The collection (ϕ_α) is maximal with respect to the first 3 conditions.

The maps ϕ_α are called local trivialisations, the sets U_α are called trivialising neighbourhoods and the map $\tau_{\alpha,\beta}$ is called the transition function for ϕ_α, ϕ_β . We write (U_α, ϕ_α) as a shorthand for the local trivialisation $\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha)$. Furthermore, we call E the total space, B the base space and F the fiber. We denote the fiber bundle by

$$F \hookrightarrow E \xrightarrow{p} B$$

or sometimes just by $E \xrightarrow{p} B$.

Fiber bundles with structure group also come with an appropriate notion of morphism:

Definition 7.2. Let $F \hookrightarrow E \xrightarrow{p} B$ and $F \hookrightarrow E' \xrightarrow{p'} B'$ be fiber bundles with structure group G . A G -morphism is a pair (H, h) of continuous maps $H: E \rightarrow E'$ and $h: B \rightarrow B'$, such that

$$\begin{array}{ccc} E & \xrightarrow{H} & E' \\ \downarrow & & \downarrow \\ B & \xrightarrow{h} & B' \end{array}$$

commutes and in local trivialisations H is given by the action of elements in G . That is for any local trivialisations (U_α, ϕ_α) of E and (V_β, ψ_β) of E' with $h(U_\alpha) \subseteq V_\beta$ there exists a continuous map

$$g_{\alpha\beta}: U_\alpha \rightarrow G,$$

such that the composition $\psi_\beta^{-1} \circ H \circ \phi_\alpha: U_\alpha \times F \rightarrow V_\beta \times F$ is given by

$$(x, f) \mapsto (h(x), g_{\alpha\beta}(x) \cdot f).$$

We usually denote such a morphism only by $H: E \rightarrow E'$. Note that h is completely determined by f , justifying this notation. If $B = B'$ we generally implicitly require h to be the identity.

⁴Wikipedia calls this a *G-Atlas*

Note that with the above definition any G -morphism is a fiber wise homeomorphism. Sometimes we may want to relax this condition by dropping the requirement that locally H is given by the action of group elements and call the resulting morphism simply a morphism of fiber bundles. For category theoretical aspects it is however nicer to work with G -morphisms. Thus we define the category $\text{Fib}_{F,G}$ to be the category whose objects are given by fiber bundles with fiber F and structure group G and whose morphisms are given by G -morphisms.

Remark 7.3. In Definition 7.1 we require surjectivity of p to exclude the case $F = \emptyset$. The requirement, that G acts effectively guarantees for the transition maps to be unique. However, this is not a strong requirement to impose, since if G does not act effectively, i.e.

$$\Phi: G \rightarrow \text{Aut}(F)$$

is not injective we get an effective group action by taking a quotient of G :

$$G/\ker\Phi \rightarrow \text{Aut}(F).$$

It follows from the universal property of the quotient topology, that $G/\ker\Phi$ still acts continuously on F , i.e. the induced map $G/\ker\Phi \times F \rightarrow F$ is still continuous (if F is locally compact we can define the exponential topology on $\text{Aut}(F)$ and the statement is equivalent to $G/\ker\Phi \rightarrow \text{Aut}(F)$ being continuous).

Remark 7.4. The maximality condition for the trivialising atlas is well-defined: Given any trivialising atlas $\mathcal{F} = (\phi_\alpha)_\alpha$ satisfying conditions (i) - (iii) of Definition 7.1 we can define

$$\begin{aligned} \hat{\mathcal{F}} := \{ & \phi: U \times F \rightarrow p^{-1}(U) \mid U \subseteq B \text{ open, } \phi \text{ homeomorphism, for any } \phi_\alpha \in \mathcal{F} \\ & \text{exists a map } \tau_{\alpha,\phi}: U \cap U_\alpha \rightarrow G, \text{ with } (\phi_\alpha^{-1} \circ \phi)(b, f) = (b, \tau_{\alpha,\phi}(b) \cdot f) \}. \end{aligned}$$

This is the maximal trivialising atlas induced by $\mathcal{F}$ satisfying (i) - (iii). Note however, that there is no unique trivialising atlas, since beginning with a different $\mathcal{F}$ may yield a different $\hat{\mathcal{F}}$ (as is the case for smooth atlases of smooth manifolds).

Proof of above remark. Given $\phi: U \times F \rightarrow p^{-1}(U)$, $\phi': U' \times F \rightarrow p^{-1}(U') \in \hat{\mathcal{F}}$, we claim that there exists $\tau: U \cap U' \rightarrow G$, such that

$$(\phi^{-1} \circ \phi')(b, f) = (b, \tau(b) \cdot f).$$

For $b_0 \in B$ choose $(\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha)) \in \mathcal{F}$ with $b_0 \in U_\alpha$. Then there exists

$$\tau_{\alpha,\phi}: U \cap U_\alpha \rightarrow G, \tau_{\alpha,\phi'}: U' \cap U_\alpha \rightarrow G$$

with

$$(\phi_\alpha^{-1} \circ \phi)(b, f) = (b, \tau_{\alpha,\phi}(b) \cdot f), (\phi_\alpha^{-1} \circ \phi')(b, f) = (b, \tau_{\alpha,\phi'}(b) \cdot f).$$

Then

$$(\phi^{-1} \circ \phi_\alpha) \circ (\phi_\alpha^{-1} \circ \phi'): (U_\alpha \cap U \cap U') \times F \rightarrow (U_\alpha \cap U \cap U') \times F$$

is given by

$$(b, f) \mapsto (b, \tau_{\alpha,\phi}(b)^{-1} \tau_{\alpha,\phi'}(b) \cdot f).$$

We conclude that around $\{b_0\} \times F$ we have

$$(\phi^{-1} \circ \phi')(b, f) = (b, \tau(b) \cdot f)$$

for $\tau(b) := \tau_{\alpha,\phi}(b)^{-1} \tau_{\alpha,\phi'}(b)$. Since group inversion and multiplication are continuous, τ is continuous. As b_0 was chosen arbitrary, such a map τ can be defined on all of $U \cap U'$. Furthermore, the group action of G is effective, hence τ is unique and the above argument shows that it is continuous. $\square$

For the transition functions we have the following Lemma, where again the effectiveness of the group action is crucial.

Lemma 7.5. *Let $F \hookrightarrow E \xrightarrow{p} B$ be a fiber bundle with structure group G . Then (using the notation in Definition 7.1):*

- (i) $\tau_{\alpha,\alpha}(b) = e$
- (ii) $\tau_{\alpha,\beta}^{-1}(b) = \tau_{\beta,\alpha}(b)$
- (iii) $\tau_{\gamma,\alpha}(b)\tau_{\alpha,\beta}(b) = \tau_{\gamma,\beta}(b)$

Proof. Let us first proof (iii), then (i) and (ii) follow directly from the group axioms. We compute:

$$\begin{aligned} (b, \tau_{\gamma,\beta}(b) \cdot f) &= (\phi_\gamma^{-1} \circ \phi_\beta)(b, f) \\ &= (\phi_\gamma^{-1} \circ \phi_\alpha \circ \phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\gamma,\alpha}(b)\tau_{\alpha,\beta}(b) \cdot f). \end{aligned}$$

Since the transition functions are uniquely determined, the statement follows. $\square$

Interestingly, the transition functions already uniquely determine the fiber bundle, in a sense made precise by the following theorem.

Theorem 7.6 (Fiber bundle construction A). *Let B, F be (non-empty) topological spaces and G a topological group acting effectively on F . Suppose we are given the following data:*

- (i) *An open cover $\mathcal{U} := (U_\alpha)_\alpha$ of B (where α ranges over some index set A)*
- (ii) *A collection $(\tau_{\alpha,\beta})$ of continuous maps*

$$\tau_{\alpha,\beta}: U_\alpha \cap U_\beta \rightarrow G$$

for each pair $U_\alpha, U_\beta \in \mathcal{U}$

- (iii) *$\tau_{\gamma,\alpha}(b)\tau_{\alpha,\beta}(b) = \tau_{\gamma,\beta}(b)$ for all $b \in U_\alpha \cap U_\beta \cap U_\gamma$*

Then there exists a unique (up to isomorphism) fiber bundle $F \hookrightarrow E \xrightarrow{p} B$ with structure group G , such that the maximal trivialising atlas is induced by a trivialising atlas of the form

$$(\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha))_\alpha,$$

such that $(\phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\alpha,\beta}(b) \cdot f)$.

Proof. Consider the space

$$E := \left(\coprod_{\alpha \in A} U_\alpha \times F \right) / \sim$$

where the equivalence relation $\sim$ is defined by

$$(b, f, \beta) \sim (b, \tau_{\alpha,\beta}(b) \cdot f, \alpha)$$

and let $p: E \rightarrow B, [(b, f, \alpha)] \mapsto b$. Furthermore, let $\pi: \coprod_\alpha U_\alpha \times F \rightarrow E$ be the projection map.

Step 1. $\sim$ is an equivalence relation.

Transitivity follows directly from condition (iii). Reflexivity and symmetry then follow from the group axioms.

Step 2. p is well-defined, continuous and surjective.

Clearly p is well-defined and surjective as $F \neq \emptyset$. Continuity then follows directly from the universal property of the quotient topology.

Step 3. Defining the trivialising atlas

Consider

$$\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha), (b, f) \mapsto [(b, f, \alpha)].$$

Clearly ϕ_α is well-defined ($[(b, f, \alpha)] \in p^{-1}(U_\alpha)$) and continuous (the projection map π is continuous). The inverse is given by

$$\phi_\alpha^{-1}([(b, f, \beta)]) = (b, \tau_{\alpha,\beta}(b) \cdot f).$$

To show that ϕ_α^{-1} is continuous, we show that ϕ_α is an open map. To this end, let $U \times V \subseteq U_\alpha \times F$ be open and consider

$$\theta_{\alpha,\beta}: (U_\alpha \cap U_\beta) \times F \rightarrow (U_\alpha \cap U_\beta) \times F, (b, f) \mapsto (b, \tau_{\alpha,\beta}(b) \cdot f).$$

Clearly $\theta_{\alpha,\beta}$ is continuous (in fact it is a homeomorphism), hence

$$V_\beta := \theta_{\alpha,\beta}^{-1}((U \cap U_\beta) \times V) \subseteq (U_\alpha \cap U_\beta) \times F$$

is open and thus

$$\pi^{-1}(\phi_\alpha(U \times V)) = \coprod_{\beta \in A} V_\beta \subseteq \coprod_{\beta \in A} U_\beta \times F$$

is open. Thus $\phi_\alpha(U \times V)$ is open in E and since $p^{-1}(U_\alpha)$ is also open, we conclude that ϕ_α is an open map. Furthermore, the diagram

$$\begin{array}{ccc} U_\alpha \times F & \xrightarrow{\phi_\alpha} & p^{-1}(U_\alpha) \\ & \searrow & \swarrow p \\ & U_\alpha & \end{array}$$

commutes, such that ϕ_α is a local trivialisation. Moreover,

$$(\phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\alpha,\beta}(b) \cdot f).$$

So the collection of maps $(\phi_\alpha)_\alpha$ is a trivialising atlas with the desired transition functions, making $F \hookrightarrow E \xrightarrow{p} B$ into a fiber bundle with structure group G .

Step 4. Uniqueness

Given two fiber bundles

$$\begin{aligned} \delta &:= (F \hookrightarrow E \xrightarrow{p} B) \\ \varepsilon &:= (F \hookrightarrow \tilde{E} \xrightarrow{\tilde{p}} B) \end{aligned}$$

with structure group G and trivialising atlases

$$\begin{aligned} (\phi_\alpha: U_\alpha \times F &\rightarrow p^{-1}(U_\alpha))_{\alpha \in A} \\ (\tilde{\phi}_\alpha: U_\alpha \times F &\rightarrow \tilde{p}^{-1}(U_\alpha))_{\alpha \in A} \end{aligned}$$

respectively, such that

$$(\phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\alpha,\beta}(b) \cdot f) = (\tilde{\phi}_\alpha^{-1} \circ \tilde{\phi}_\beta)(b, f).$$

We can then construct an isomorphism $\Gamma: \delta \rightarrow \varepsilon$ as follows: for $x \in E$ choose $U_\alpha \in \mathcal{U}$ with $p(x) \in U_\alpha$. Then define

$$\Gamma(x) := (\tilde{\phi}_\alpha \circ \phi_\alpha^{-1})(x).$$

Well-definedness of Γ follows from the commutativity of the following diagram:

$$\begin{array}{ccccc} & & (U_\alpha \cap U_\beta) \times F & & \\ & \swarrow \phi_\beta & \downarrow \tau & \searrow \tilde{\phi}_\beta & \\ p^{-1}(U_\alpha \cap U_\beta) & & & & \tilde{p}^{-1}(U_\alpha \cap U_\beta) \\ & \swarrow \phi_\alpha & \downarrow & \searrow \tilde{\phi}_\alpha & \\ & & (U_\alpha \cap U_\beta) \times F & & \end{array}$$

where $\tau(b, f) := (b, \tau_{\alpha,\beta}(b) \cdot f)$. Locally Γ is given by a continuous map, so it is continuous. Analogously we define the inverse map. Furthermore,

$$\begin{array}{ccc} E & \xrightarrow{\Gamma} & \tilde{E} \\ & \searrow p \quad \swarrow \tilde{p} & \\ & B & \end{array}$$

commutes, so Γ is a bundle isomorphism. This completes the proof. $\square$

We may also formulate Theorem 7.6 slightly different, if we are given the fibers $F_x = p^{-1}(x)$ as sets. This is usually more practical, as the description of the bundle is more explicit:

Theorem 7.7 (Fiber bundle construction B). *Let B, F be (non-empty) topological spaces and G a topological group acting effectively on F . Given sets F_x for each $x \in B$ we define*

$$E := \coprod_{x \in B} F_x$$

and $p: E \rightarrow B$ by sending each element in F_x to x . Suppose we are given the following data:

- (i) An open cover $\mathcal{U} := (U_\alpha)_\alpha$ of B (where α ranges over some index set A)
- (ii) A collection of bijections $\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha)$, that restrict to bijections $\{x\} \times F \rightarrow F_x$
- (iii) A collection $(\tau_{\alpha,\beta})$ of continuous maps

$$\tau_{\alpha,\beta}: U_\alpha \cap U_\beta \rightarrow G$$

for each pair $U_\alpha, U_\beta \in \mathcal{U}$, such that $(\phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\alpha,\beta}(b) \cdot f)$

Then E has a unique topology making $F \hookrightarrow E \xrightarrow{p} B$ into a fiber bundle with structure group G , where the maps ϕ_α are local trivialisations.

Proof. Define the topology on E as the final topology with respect to the maps ϕ_α , i.e.

$$V \subseteq E \text{ is open} \Leftrightarrow \phi_\alpha^{-1}(V) \subseteq U_\alpha \times F \text{ is open for each } \alpha$$

We show that the maps ϕ_α are open with this topology. To this end, let $W \subseteq U_\alpha \times F$ be open. Then

$$\phi_\beta^{-1}(\phi_\alpha(W)) = (\phi_\beta^{-1} \circ \phi_\alpha)(W \cap ((U_\alpha \cap U_\beta) \times F)) \subseteq U_\beta \times F$$

is open, since $\phi_\beta \circ \phi_\alpha$ is a homeomorphism by condition (iii). Thus ϕ_α is a homeomorphism. In particular, for any open set $U \subseteq U_\alpha$, we get that

$$p^{-1}(U) = \phi_\alpha(U \times F)$$

is open, since by (ii) the following diagram commutes:

$$\begin{array}{ccc} U_\alpha \times F & \xrightarrow{\phi_\alpha} & p^{-1}(U_\alpha) \\ & \searrow p & \swarrow \\ & U_\alpha & \end{array}$$

So p is also continuous. Moreover, as $F \neq \emptyset$, we get that p is surjective. It remains to show uniqueness. So let E be equipped with any topology, such that p is continuous and ϕ_α are homeomorphisms. Let $V \subseteq E$, such that

$$W_\alpha := \phi_\alpha^{-1}(V) \subseteq U_\alpha \times F$$

is open for all α . Then

$$V = \bigcup_{\alpha} W_\alpha,$$

since the U_α cover B (and hence $E = \bigcup_{\alpha} \text{im } \phi_\alpha$). As the ϕ_α are homeomorphisms and $p^{-1}(U_\alpha)$ is open, we conclude that V is open. Furthermore, the topology can not be finer than the final topology defined above, such that the two must coincide. $\square$

Example 7.8. Suppose we are given a (topological) rank n vector bundle $E \xrightarrow{\pi} B$. We can use Theorem 7.7 to define the **frame bundle** $\text{GL}(E)$ of E as follows. Let $\text{GL}(E_x)$ denote the set of all ordered bases of the vector space $E_x := \pi^{-1}(x)$ and set

$$\text{GL}(E) := \coprod_{x \in B} \text{GL}(E_x).$$

Now choose an open cover (U_α) of B over which E is trivial, i.e. there exist local frames $X_1^\alpha, \dots, X_n^\alpha$ over U_α . We define the local trivialisations for the bundle by

$$\begin{aligned} \phi_\alpha: U_\alpha \times \mathrm{GL}(n) &\rightarrow p^{-1}(U_\alpha) \\ (x, A) &\mapsto \left(\sum_{i=1}^n a_{i1} X_i^\alpha(x), \dots, \sum_{i=1}^n a_{in} X_i^\alpha(x) \right), \end{aligned}$$

where $A = (a_{ij})$, i.e. $\phi_\alpha(x, A)$ is the basis of E_x , whose coordinates in the frame (X_i^α) are precisely the columns of the matrix A . Clearly ϕ_α is a bijection. Moreover for $x \in U_\alpha \cap U_\beta$ we have

$$(\phi_\alpha^{-1} \circ \phi_\beta)(x, A) = (x, \tau_{\alpha,\beta}(x)A),$$

where $\tau_{\alpha,\beta}(x) \in \mathrm{GL}(n)$ is the change of basis matrix from $(X_i^\beta(x))$ to $(X_i^\alpha(x))$, which depends continuously on x . So applying Theorem 7.7 we can see $\mathrm{GL}(E)$ as principal $\mathrm{GL}(n)$ -bundle over B . Using Theorem 7.11 the same construction works in the smooth category. It is interesting to note that E is trivial if and only if $\mathrm{GL}(E)$ admits a global section.

If E carries a fiber metric we may similarly construct the **orthonormal frame bundle** $O(M)$ as a principal $O(n)$ -bundle and if E is oriented we can construct the **oriented frame bundle** $\mathrm{GL}^+(M)$ as a principal $\mathrm{GL}^+(n)$ -bundle. Of course we can also combine the two to get the **oriented orthonormal frame bundle** $\mathrm{SO}(M)$.

Theorem 7.6 describes a construction which can be applied to objects in $\mathrm{Fib}_{F,G}$. The following theorem describes an analogous construction for the morphisms, which will be the key to the functoriality result of Theorem 7.17 below.

Proposition 7.9. *Let $E \rightarrow B$ and $E' \rightarrow B'$ be fiber bundles with fiber F and structure group G . Suppose we are given the following data:*

- (i) *A continuous map $h: B \rightarrow B'$,*
- (ii) *Trivialisations $(U_\alpha, \phi_\alpha)_\alpha$ of E and $(V_\beta, \psi_\beta)_\beta$ of E' , such that the U_α cover B and for each α there exists some β with $h(U_\alpha) \subseteq V_\beta$,*
- (iii) *For each pair α, β with $h(U_\alpha) \subseteq V_\beta$ a continuous map $g_{\alpha\beta}: U_\alpha \rightarrow G$, which transforms as follows: for some other α', β' with $h(U_{\alpha'}) \subseteq V_{\beta'}$ it holds that*

$$g_{\alpha\beta}(x) = \tau_{\beta'\beta}^{-1}(x) g_{\alpha'\beta'}(x) \tau_{\alpha'\alpha}(x)$$

for all $x \in U_\alpha \cap U_{\alpha'}$, where $\tau_{\alpha'\alpha}$ and $\tau_{\beta'\beta}$ are the respective transition functions.

Then there exists a unique G -morphism $E \rightarrow E'$, which corresponds to the maps

$$U_\alpha \times F \rightarrow V_\beta \times F, \quad (x, f) \mapsto (h(x), g_{\alpha\beta}(x) \cdot f)$$

under the local trivialisations ϕ_α and ψ_β for all α, β with $h(U_\alpha) \subseteq V_\beta$.

Proof. By definition any G -morphism is locally determined by maps $g_{\alpha\beta}: U_\alpha \rightarrow G$ and these maps transform as in (iii) by definition of the transition functions. So conversely we may define a G -morphism locally by these $g_{\alpha\beta}$ and since they transform correctly we get a globally defined object. $\square$

Smooth Bundles

If we are in the smooth category, we can analogously define smooth fiber bundles by requiring the spaces E, B and F to be smooth manifolds (where either F or B has no boundary), the group G to be a Lie group and all maps to be smooth. In particular, the local trivialisation are diffeomorphisms. All of the above results also hold in the smooth category.

Theorem 7.10 (Fiber bundle construction A'). *The conclusions of Theorem 7.6 remain true in the smooth category.*

Proof. The only difficulty is in defining the smooth structure on

$$E := \left(\coprod_{\alpha \in A} U_\alpha \times F \right) / \sim$$

This smooth structure is defined via the local trivialisations (ϕ_α) as follows: Choose an open cover (V_α) of B , such that

- $V_\alpha \subseteq U_\alpha$
- there exist smooth charts $h_\alpha: W_\alpha \rightarrow V_\alpha$

Furthermore, choose a smooth atlas

$$(k_i: W_i^F \rightarrow V_i^F)_i$$

of F . Then

$$(\phi_\alpha \circ (h_\alpha \times k_i): W_\alpha \times W_i^F \rightarrow E)_{i,\alpha}$$

defines a smooth atlas for E , since the maps $\phi_\alpha^{-1} \circ \phi_\beta$ are diffeomorphisms. It is then easy to verify, that this atlas satisfies the second countability and Hausdorff axioms. Furthermore, this smooth structure is uniquely determined by requiring the local trivialisations to be diffeomorphisms. $\square$

Theorem 7.11 (Fiber bundle construction B'). *The conclusions of Theorem 7.7 remain true in the smooth category.*

Proof. As in the above theorem, the local trivialisations determine a unique smooth structure on E . $\square$

Pullback Bundles

Given a fiber bundle $F \hookrightarrow E \xrightarrow{p} B$ and a map $g: B' \rightarrow B$ we can construct the pullback bundle g^*E using Theorem 7.7 as the fiber bundle over B' , where the fiber over $b' \in B'$ is given by the fiber of E over $g(b')$, i.e.

$$g^*E = \coprod_{b' \in B'} p^{-1}(g(b')).$$

Given a local trivialisation $\phi: U \times F \rightarrow E$ we define a local trivialisation of g^*E by

$$g^{-1}(U) \times F \rightarrow g^*E, (b', f) \mapsto (b', \phi(g(b'), f)).$$

By projecting onto the second factor we get a bundle homomorphism $G: g^*E \rightarrow E$, making the following diagram commute:

$$\begin{array}{ccc} g^*E & \xrightarrow{G} & E \\ \downarrow & & \downarrow p \\ B' & \xrightarrow{g} & B \end{array}$$

In fact it is easily seen, that this diagram is a pullback diagram in the category **Top** of topological spaces, which gives a useful characterisation of the pullback bundle. The same construction works in the smooth category using Theorem 7.11 as long as either F has no boundary or B and B' have no boundary.

Lemma 7.12. *In the above situation the following holds:*

(i) *If g is an immersion, so is G*

(ii) *If g is an embedding, so is G*

Proof. In local trivialisations we can write G as

$$g^{-1}(U) \times F \rightarrow U \times F, (b, f) \mapsto (g(b'), f),$$

such that (i) follows. Now let g be an embedding. We have $\text{im}G = p^{-1}(g(B'))$, so we can define

$$H: \text{im}G \rightarrow f^*E, x \mapsto (g^{-1}(p(x)), x).$$

Clearly H is continuous and $G \circ H = \text{id}_{\text{im}G}$ as well as $H \circ G = \text{id}_{g^*E}$, so G is in fact an embedding. $\square$

One usecase for pullback bundles is to define vector fields along curves $\gamma: I \rightarrow M$ for a smooth manifold M as sections in the pullback bundle γ^*TM . The following Lemma says, that we can equivalently define them as maps

$$V: I \rightarrow TM$$

with $V(t) \in T_{\gamma(t)}M$ (as is done in [Lee18]).

Lemma 7.13. *Let $F \hookrightarrow E \rightarrow M$ be a smooth fiber bundle and $g: N \rightarrow M$ a smooth map. Then a section $N \xrightarrow{\sigma} g^*E$ is smooth if and only if the composition*

$$N \xrightarrow{\sigma} g^*E \xrightarrow{G} E$$

*is smooth, where G is the induced bundle homomorphism $g^*E \rightarrow E$ discussed above.*

Proof. This follows directly from the definition of the smooth structure of the pullback bundle given above. $\square$

7.2 Changing the fiber

Given a fiber bundle, Theorem 7.6 allows us to change the fiber:

Corollary 7.14 (Changing the fiber). *Let $F \hookrightarrow E \xrightarrow{p} B$ be a fiber bundle with structure group G . Given a space F' upon which G acts effectively, there exists a unique (up to isomorphism) fiber bundle*

$$F' \hookrightarrow E' \xrightarrow{p'} B$$

which has the same trivialising neighbourhoods (U_α) and the same transition functions $(\tau_{\alpha,\beta})$ as $F \hookrightarrow E \xrightarrow{p} B$. We denote this bundle by $E \times_G F'$ (see Lemma 7.25 for the origin of this notation).

Theorem 7.6 tells us, that the new bundle still retains all of the information of the old bundle, as it is uniquely determined by the transition functions, i.e. we can simply change the fiber again to get the original bundle back.

Example 7.15. Given any fiber bundle $F \hookrightarrow E \xrightarrow{p} B$ with structure group G we can apply the above corollary with $F' := G$ upon which G acts effectively by left-translations to get a principal G -bundle over B . This bundle is called the **underlying principal bundle** of $E \rightarrow B$ (see Definition 7.19).

Example 7.16. Using the above corollary we can also construct the frame bundle from Example 7.8 more abstractly by defining it as the underlying principal bundle of $E \rightarrow F$. Note that the uniqueness results in Theorem 7.6 shows, that the bundle constructed in this way is in fact isomorphic to $\text{GL}(E)$. The construction in Example 7.8 of course has the benefit of being more concrete.

We can use Proposition 7.9 to get a much stronger form of Corollary 7.14:

Theorem 7.17 (The Change of Fiber Functor). *Let G be a topological group acting effectively on spaces F and F' . Then changing the fiber from F to F' defines a functor*

$$\times_G F': \text{Fib}_{F,G} \rightarrow \text{Fib}_{F',G}.$$

Moreover the composition $\times_G F' \circ \times_G F$ is naturally isomorphic to the identity on $\text{Fib}_{F,G}$. Thus $\times_G F$ defines an equivalence of categories.

Proof. The proof is straightforward but takes some care to write down precisely. For every bundle E in $\text{Fib}_{F,G}$ choose a bundle $E \times_G F'$ in $\text{Fib}_{F',G}$ with the same trivialising neighbourhoods and transition functions as E (using Corollary 7.14). On morphisms we define the functor as follows. Let

$$\begin{array}{ccc} E_1 & \xrightarrow{H_1} & E_2 \\ \downarrow & & \downarrow \\ B_1 & \xrightarrow{h_1} & B_1 \end{array}$$

be a G -morphism in $\text{Fib}_{F,G}$. Choose a trivialising cover (U_α) of E_1 and (V_β) of E_2 , such that for all α there exists β with $h_1(U_\alpha) \subseteq V_\beta$. Then in local trivialisations H is given by

$$U_\alpha \times F \rightarrow V_\beta \times F, \quad (x, f) \mapsto (h_1(x), g_{\alpha\beta}(x) \cdot f)$$

for some $g_{\alpha\beta}: U_\alpha \rightarrow G$, which transform as required in Proposition 7.9. Since by definition (U_α) is also a trivialising cover of $E_1 \times_G F'$ with the same transition functions as E_1 and similarly for $E_2 \times_G F'$, we can apply Proposition 7.9 to construct a G -morphism $K: E_1 \times_G F' \rightarrow E_2 \times_G F'$. We claim that K does not depend on the chosen covers (U_α) and (V_β) . So let $(U'_{\alpha'})$ and $(V'_{\beta'})$ be different trivialising covers with the same property and define $K': E_1 \times_G F' \rightarrow E_2 \times_G F'$ in the same way. Then we can also use the covers $(U_\alpha) \cup (U'_{\alpha'})$ and $(V_\beta) \cup (V'_{\beta'})$ to construct a G -morphism $K'': E_1 \times_G F' \rightarrow E_2 \times_G F'$, which by definition agrees with K on $E_1|_{U_\alpha}$ for all α , such that $h_1(U_\alpha) \subseteq V_\beta$ for some β and similarly for K' . Thus $K = K'' = K'$. We may then define

$$H_1 \times_G F' := K.$$

To see that this is functorial let

$$\begin{array}{ccc} E_2 & \xrightarrow{H_2} & E_3 \\ \downarrow & & \downarrow \\ B_2 & \xrightarrow{h_2} & B_3 \end{array}$$

be another G -morphism. Let $(U_\alpha), (V_\beta)$ and (W_γ) be trivialising covers of E_1, E_2 and E_3 respectively, such that for all α there exists β with $h_1(U_\alpha) \subseteq V_\beta$ and for all β there exists γ with $h_2(V_\beta) \subseteq W_\gamma$. Then the covers (U_α) and (W_γ) may be used to construct $(H_2 \circ H_1) \times_G F'$ as above and for α, β, γ with $h_1(U_\alpha) \subseteq V_\beta$ and $h_2(V_\beta) \subseteq W_\gamma$ this map clearly agrees with the composition $H_2 \times_G F' \circ H_1 \times_G F'$. Moreover it is clear that the identity gets mapped to the identity under $\times_G F'$ by using $(V_\beta) = (U_\alpha)$ for the construction.

Next we show the statement about the equivalence of categories. First we define a canonical isomorphism

$$\eta_E: E \rightarrow (E \times_G F') \times_G F$$

as follows. Let (U_α) be a trivialising cover for E with transition functions $\tau_{\beta\alpha}: U_\alpha \cap U_\beta \rightarrow G$. Then by definition (U_α) is also a trivialising cover for $(E \times_G F') \times_G F$ with the same transition functions. For α, β with $U_\alpha \subseteq U_\beta$ set $G_{\alpha\beta} := \tau_{\beta\alpha}$ (the reversed order of α and β only has to do with the notational conventions used for the transition functions). Then these transform as in Proposition 7.9, and thus define a G -morphism $\eta_E: E \rightarrow (E \times_G F') \times_G F$. This is an isomorphism, since it corresponds to the identity

$$U_\alpha \times F \rightarrow U_\alpha \times F$$

under local trivialisations and as before we see that η_E does not depend on the cover (U_α) . Finally for the naturality let $H_1: E_1 \rightarrow E_2$ be as above and again choose trivialising covers (U_α) of E_1

and (V_β) of E_2 , such that for all α there exists β with $h_1(U_\alpha) \subseteq V_\beta$. Then for fixed α, β with $h_1(U_\alpha) \subseteq V_\beta$ the maps H_1 and $(H_1 \times_G F') \times_G F$ both correspond to the same map

$$U_\alpha \times F \rightarrow V_\beta \times F.$$

Moreover since η_{E_1} and η_{E_2} correspond to the identity maps on $U_\alpha \times F$ and $V_\beta \times F$, respectively, the statement follows. $\square$

Changing the fiber satisfies the following naturality condition with respect to the pullback operation.

Proposition 7.18. *Let $F \hookrightarrow E \xrightarrow{p} B$ be a fiber bundle with structure group G, F' a space upon which G acts effectively and $f: B' \rightarrow B$ a continuous map. Then*

$$(f^*E) \times_G F' \cong f^*(E \times_G F').$$

Proof. For the sake of brevity let us call an open cover (U_α) of some base space X of a fiber bundle $Q \rightarrow X$ a trivialising cover for Q , if the U_α are trivialising neighbourhoods for Q . Now let (U_α) be a trivialising cover for E with transition functions $\tau_{\alpha, \beta}: U_\alpha \cap U_\beta \rightarrow G$. Then by definition $(f^{-1}(U_\alpha))$ is a trivialising cover for f^*E with transition functions

$$\tau_{\alpha, \beta} \circ f: f^{-1}(U_\alpha) \cap f^{-1}(U_\beta) \rightarrow G.$$

Thus again by definition $(f^{-1}(U_\alpha))$ is a trivialising cover of $(f^*E) \times_G F'$ with transition functions $\tau_{\alpha, \beta} \circ f$. Similarly we see that this same cover is a trivialising cover for $f^*(E \times_G F')$ with $\tau_{\alpha, \beta} \circ f$ as transition functions. Thus the statement follows from the uniqueness result of Theorem 7.6. $\square$

More abstractly we can formulate this proposition as follows. Let $\mathcal{F}_{F,G}(B)$ denote the set of isomorphism classes of fiber bundles with fiber F and structure group G over a given base space B . Then $\mathcal{F}_{F,G}(-)$ defines a contravariant functor $\mathbf{Top} \rightarrow \mathbf{Set}$ and changing the fiber defines a natural isomorphism of functors $\mathcal{F}_{F,G}(-) \Rightarrow \mathcal{F}_{F',G}(-)$.

7.3 Principal Bundles

Definition 7.19. Let G be a topological group. A G -principal bundle is a fiber bundle with fiber G and structure group G acting by left translations.

The total space of a G -principal bundle $P \xrightarrow{\pi} B$ carries a natural right action of G defined as follows. Around $y \in P$ choose a local trivialisation

$$\phi: U \times G \rightarrow \pi^{-1}(U)$$

and let $\phi^{-1}(y) = (\pi(y), h)$. Then set

$$y \cdot g := \phi(\pi(y), hg).$$

This is well defined since by definition G acts on itself by *left* translations under a change of local trivialisation. We then have the following Lemma.

Lemma 7.20. *Let $P \xrightarrow{\pi} B$ be a G -principal bundle. Then π descends to a homeomorphism $P/G \rightarrow B$.*

Proof. Clearly π descends to a continuous bijection

$$\hat{\pi}: P/G \rightarrow B.$$

To show that $\hat{\pi}$ is in fact a homeomorphism let us first assume that P is trivial. Let $\tau: B \times G \rightarrow P$ be a trivialisation. Consider the commutative diagram

$$\begin{array}{ccccc} B \times G & \xrightarrow[\cong]{\tau} & P & \xrightarrow{\pi} & B \\ \downarrow & & \downarrow & \nearrow \hat{\pi} & \\ B & \xleftarrow[\cong]{\hat{p}} & (B \times G)/G & \xrightarrow[\cong]{\hat{\tau}} & P/G \end{array}$$

where the action of G on $B \times G$ corresponds to the action of G on P under τ and is explicitly given by

$$(x, h) \cdot g = (x, hg).$$

Moreover $\hat{\tau}$ is the map induced on the quotients by τ and $\hat{p}$ is the homeomorphism induced by the projection $p: B \times G \rightarrow B$. Then for the lower route in the diagram we have

$$\hat{\pi} \circ \hat{\tau} \circ \hat{p}^{-1} = \text{id}_B.$$

Since $\hat{\tau}$ and $\hat{p}^{-1}$ are homeomorphisms, so is $\hat{\pi}$. Now for the general case note that the local triviality of P together with the above shows that $\hat{\pi}$ is a local homeomorphism. Since it is bijective the statement follows. $\square$

A morphism of principal bundles should respect this right action, thus we define:

Definition 7.21. A morphism from a G -principal bundle $P \rightarrow B$ to an H -principal bundle $P' \rightarrow B'$ is a morphism $(F, f): (P \rightarrow B) \rightarrow (P' \rightarrow B')$ of the respective fiber bundles (see Definition 7.2) together with a homomorphism $\phi: G \rightarrow H$ of topological groups such that F is ϕ -equivariant, that is

$$F(x \cdot g) = F(x) \cdot \phi(g)$$

for all $x \in P$ and $g \in G$. If $G = H$ we usually assume ϕ to be the identity and if $B = B'$ we assume f to be the identity.

The added condition of equivariance is actually quite strong as illustrated by the following proposition.

Proposition 7.22. *Any morphism of G -principal bundles over the same base space is an isomorphism.*

Proof. First suppose we are given a morphism

$$F: B \times G \rightarrow B \times G$$

of trivial bundles. Then by definition F is of the form

$$F(x, g) = (x, \phi_x(g))$$

for a continuous family of equivariant maps $\phi_x: G \rightarrow G$. By equivariance $\phi_x(g) = \phi_x(e)g$ and thus an inverse of F is given by

$$F^{-1}(x, g) := (x, \phi_x(e)^{-1}g).$$

Since every G -principal bundle is locally trivial this shows that we can always find local inverses in the general case and since inverses are unique they fit together to define a global inverse. $\square$

Corollary 7.23. *A G -principal bundle is trivial if and only if it admits a section.*

Proof. Suppose we are given a section $s: B \rightarrow P$ of $P \rightarrow B$. Then $B \times G \rightarrow P$, $(x, g) \mapsto s(x) \cdot g$ defines a morphism of G -principal bundles and hence is an isomorphism by Proposition 7.22. $\square$

Remark 7.24. In the definition of G -principal bundles we may also require G to act by right translations instead of left translations on itself. Correspondingly we then get an induced left G -action on the total space. Let us call those bundles right G -principal bundles and the bundles from Definition 7.19 left G -principal bundles. Given a right G -principal bundle $P \rightarrow B$ we can always turn it into a left G -principal bundle as follows. Let $\tau: U \times G \rightarrow P$ be any local trivialisation and set

$$\tau' := U \times G \xrightarrow{\text{id} \times \text{inv}} U \times G \xrightarrow{\tau} P,$$

where $\text{inv}: G \rightarrow G$ is the inversion map. Then the maps τ' constructed in this way give $P \rightarrow B$ the structure of a left G -principal bundle. The new right G -action on P can be expressed by the left G -action as

$$y \cdot g = g^{-1} \cdot y$$

Note that any morphism of right principal bundles also respects this new right action and thus defines a morphism of the corresponding left principal bundles. In other words, this construction defines an isomorphism between the category of right principal bundles and the category of left principal bundles. Thus there is virtually no difference between these two notions.

Given any fiber bundle we have seen in Corollary 7.14, that we can change the fiber. If we start with a principal bundle we can make this construction quite explicit:

Lemma 7.25 (Borel Construction). *Let $P \xrightarrow{p} B$ be a G -principal bundle and F a space upon which G acts effectively from the left. Then*

$$P \times_G F := (P \times F) / \sim, \quad (y, f) \sim (y \cdot g, g^{-1}f)$$

together with the map $\pi: P \times_G F \rightarrow B$ given by $[(y, f)] \mapsto p(y)$ is a fiber bundle with fiber F and structure group G . Moreover this construction yields the same bundle as Corollary 7.14, hence the same notation.

Proof. The map π is induced by $P \times F \rightarrow B$, $(y, f) \mapsto p(y)$ and hence continuous. To see that $P \times_G F$ is locally trivial let $\lambda: U \times G \rightarrow P$ be a local trivialisation. Then the homeomorphism

$$\lambda \times \text{id}_F: U \times G \times F \rightarrow p^{-1}(U) \times F$$

descends to a homeomorphism

$$\tilde{\kappa}: (U \times G \times F) / \sim \rightarrow (p^{-1}(U) \times F) / \sim = \pi^{-1}(U),$$

where the equivalence relation on $U \times G \times F$ is given by $(x, g, f) \sim (x, gh, h^{-1}f)$. Moreover the map

$$U \times G \times F \rightarrow U \times F, \quad (x, g, f) \mapsto (x, gf)$$

descends to a homeomorphism $(U \times G \times F) / \sim \rightarrow U \times F$ with inverse given by $(x, f) \mapsto [(x, e, f)]$. Then under this homeomorphism $\tilde{\kappa}$ corresponds to a homeomorphism $\kappa: U \times F \rightarrow \pi^{-1}(U)$, which is explicitly given by $(x, f) \mapsto [(\lambda(x, e), f)]$ and

$$\begin{array}{ccc} U \times F & \xrightarrow{\kappa} & \pi^{-1}(U) \\ & \searrow \text{proj.} & \downarrow \pi \\ & & U \end{array}$$

commutes. So $P \times_G F$ is locally trivial. Next we have to compute the transition functions. Let $U_\alpha, U_\beta \subseteq B$ be two trivialising neighbourhoods for P with local trivialisations $\lambda_\alpha, \lambda_\beta$ and transition function $\tau_{\alpha, \beta}: U_\alpha \cap U_\beta \rightarrow G$. Moreover let $\kappa_\alpha, \kappa_\beta$ be the corresponding local trivialisations for $P \times_G F$ as constructed above. Then we compute

$$(\kappa_\alpha^{-1} \circ \kappa_\beta)(x, f) = \kappa_\alpha^{-1}([(\lambda_\beta(x, e), f)]) = (x, \tau_{\alpha, \beta}(x)f).$$

So $P \times_G F$ has structure group G and from the uniqueness result of Theorem ?? it follows that this construction agrees with the one from Corollary 7.14. $\square$

Given a group homomorphism $\phi: G \rightarrow H$ of topological groups we can view H as a left G -space by $g \cdot h := \phi(g)h$. Of course this action may not be effective anymore but the Borel construction can still be used to construct an H -principal bundle out of a G -principal bundles using this action:

Proposition 7.26. *Let $P \xrightarrow{p} B$ be a G -principal bundle and $\phi: G \rightarrow H$ a homomorphism of topological groups. Then*

$$P \times_{\phi} H := (P \times H) / \sim, \quad (y, h) \sim (y \cdot g, \phi(g)^{-1}h)$$

together with the map $\pi: P \times_{\phi} H \rightarrow B$ given by $[(y, h)] \mapsto p(y)$ is an H -principal bundle. Moreover this construction is natural in the sense that if $f: B' \rightarrow B$ is a continuous map then

$$(f^*P) \times_{\phi} H \cong f^*(P \times_{\phi} H)$$

and it is also functorial in the sense that if $\psi: H \rightarrow K$ is another homomorphism of topological groups then

$$(P \times_{\phi} H) \times_{\psi} K \cong P \times_{\psi \circ \phi} K.$$

Proof. The proof is the same as the one for Lemma 7.25, for the special case $F = H$ and the specific left action of G on H determined by ϕ . We only used the effectiveness of the group action at the very end to make use of Theorem ??, which only makes sense if the group action is effective, such that the transition functions are uniquely determined (and besides that we required the group action to be effective in the definition of a fiber bundle, so the statement of the Lemma wouldn't even technically make sense). The difference in the statement of this proposition is that $P \times_{\phi} H$ will be a bundle with structure group H and not G and H does again act effectively on itself. Using the notation of Lemma 7.25 we then see that changing local trivialisations is given by

$$(\kappa_{\alpha}^{-1} \circ \kappa_{\beta})(x, h) = (x, \phi(\tau_{\alpha, \beta}(x))h),$$

such that $P \times_{\phi} H$ is indeed an H -principal bundle. The naturality and functoriality statements are shown analogously to Proposition 7.18 and we shall use the same terminology of a trivialising cover introduced in that proof. Then the construction of the local trivialisations for $P \times_{\phi} H$ shows that any trivialising cover (U_{α}) for P with transition functions $\tau_{\alpha, \beta}$ is also a trivialising cover for $P \times_{\phi} H$ with transition functions

$$\phi \circ \tau_{\alpha, \beta}: U_{\alpha} \cap U_{\beta} \rightarrow H.$$

Hence we see as in Proposition 7.18 that both $(f^*P) \times_{\phi} H$ and $f^*(P \times_{\phi} H)$ have $(f^{-1}(U_{\alpha}))$ as a trivialising cover with transition functions

$$\phi \circ \tau_{\alpha, \beta} \circ f: f^{-1}(U_{\alpha}) \cap f^{-1}(U_{\beta}) \rightarrow H,$$

so the naturality result follows again from Theorem ??. For the functoriality note that both bundles have (U_{α}) as a trivialising cover with transition functions $\psi \circ \phi \circ \tau_{\alpha, \beta}$. $\square$

If $\mathcal{P}_G(B)$ denotes the set of isomorphism classes of G -principal bundles over a given base space B then $\mathcal{P}_G(-)$ defines a contravariant functor $\mathbf{Top} \rightarrow \mathbf{Set}$. Then the above proposition says that any $\phi: G \rightarrow H$ defines a natural transformation $\mathcal{P}\phi := \times_{\phi} H$ of functors $\mathcal{P}_G(-) \Rightarrow \mathcal{P}_H(-)$ and does so in a functorial way, that is $\mathcal{P}\psi \circ \mathcal{P}\phi = \mathcal{P}(\psi \circ \phi)$. Thus we get a functor from the category of topological groups into the functor category of functors $\mathbf{Top} \rightarrow \mathbf{Set}$.

7.4 Vector Bundles

Whitney Sums

Given vector bundles E_1, E_2 over the same base B , we can define their **Whitney Sum** $E_1 \oplus E_2$ using Theorem 7.7 as the vector bundle whose fibers are given by $E_1|_p \oplus E_2|_p$ with the obvious choice as local trivialisations. The following Lemma characterizes the Whitney sum and at the same time provides the technical grounds for an intuition.

Lemma 7.27 (Characterisation of the Whitney sum). *Let E_1 and E_2 be vector bundles over the same base space B of rank k_1 and k_2 respectively. A rank $k_1 + k_2$ vector bundle E over B is isomorphic to $E_1 \oplus E_2$ if and only if E_1 and E_2 can be embedded into E as subbundles and $E_1|_p + E_2|_p = E|_p$ for all $p \in B$.*

Let us first clarify what we mean by embedding. A vector bundle homomorphism $F: E \rightarrow E'$ is called an **embedding**, if it is injective. Then $\text{im}(F)$ is a subbundle of E' and $F: E \rightarrow \text{im}(F)$ is a vector bundle isomorphism. This follows from [Lee11] Theorem 10.34 and Prop. 10.26.

Proof of Lemma 7.27. First let $E = E_1 \oplus E_2$. Then we can define embeddings by

$$F_1: E_1 \rightarrow E, v \mapsto (v, 0), F_2: E_2 \rightarrow E w \mapsto (0, w).$$

Now let $E_1, E_2 \subseteq E$ be subbundles. Let $(U_\alpha)_\alpha$ be an open cover of B , such that E_1 and E_2 are trivial over U_α . Choose local frames

$$\sigma_{\alpha,1}^i, \dots, \sigma_{\alpha,k_i}^i: U_\alpha \rightarrow E_i$$

and define local trivialisations

$$\phi_\alpha^i: U_\alpha \times \mathbb{R}^{k_i} \rightarrow E_i, (p, v) \mapsto \sum_{j=1}^{k_i} v_j \sigma_{\alpha,j}^i(p).$$

Let $\tau_{\alpha,\beta}^i: U_\alpha \cap U_\beta \rightarrow \text{GL}(k_i)$ be the transition functions defined by

$$((\phi_\alpha^i)^{-1} \circ \phi_\beta^i)(p, v) = (p, \tau_{\alpha,\beta}^i(p) \cdot v).$$

Since $E|_p$ is the direct sum of $E_1|_p$ and $E_2|_p$, we get that

$$\sigma_{\alpha,1}^1, \dots, \sigma_{\alpha,k_1}^1, \sigma_{\alpha,1}^2, \dots, \sigma_{\alpha,k_2}^2$$

is a local frame for E , thus defines a local trivialisation by

$$\psi_\alpha: U_\alpha \times \mathbb{R}^{k_1} \times \mathbb{R}^{k_2} \rightarrow E, (p, v, w) \mapsto \phi_\alpha^1(p, v) + \phi_\alpha^2(p, w).$$

Then

$$(\psi_\alpha^{-1} \circ \psi_\beta)(p, v, w) = (p, \tau_{\alpha,\beta}^1(p) \cdot v, \tau_{\alpha,\beta}^2(p) \cdot w).$$

Note that $E_1 \oplus E_2$ is also trivial over U_α via

$$\varphi_\alpha: U_\alpha \times \mathbb{R}^{k_1} \times \mathbb{R}^{k_2} \rightarrow E_1 \oplus E_2, (p, v, w) \mapsto (\phi_\alpha^1(p, v), \phi_\alpha^2(p, w))$$

and $\varphi_\alpha^{-1} \circ \varphi_\beta = \psi_\alpha^{-1} \circ \psi_\beta$, thus

$$E \cong E_1 \oplus E_2$$

by the uniqueness result in Theorem 7.6. □

This characterisation gives an intuitive reasoning behind the fact, that the Whitney sum of two Moebius bundles is trivial:




Figure 2: Illustration of the Whitney sum of two Moebius bundles [source: <https://math.stackexchange.com/questions/1009126/global-trivialization-of-m-oplus-m>]

Lemma 7.28. *Let E, E_1 and E_2 be vector bundles over a manifold M and $\sigma_i: M \rightarrow E_i$ smooth sections for $i = 1, 2$. Then the following maps are smooth:*

- (i) $(\sigma_1, \sigma_2): N \rightarrow E_1 \oplus E_2, p \mapsto (\sigma_1(p), \sigma_2(p))$
- (ii) The addition map $E \oplus E \rightarrow E, (v, w) \mapsto v + w$
- (iii) The scalar multiplication map $\mathbb{R} \times E \rightarrow E, (\lambda, v) \mapsto \lambda v$

Proof. All statements follow easily by writing the maps in local trivialisations. $\square$

The Normal Bundle

We can generalize Lemma 10.35 from [Lee11] to Riemannian manifolds:

Lemma 7.29 (Orthogonal complement bundles). *Let M be a Riemannian manifold, $S \subseteq M$ a submanifold and $E \subseteq TM|_S$ a subbundle. Then*

$$E^\perp := \{(p, v) \mid p \in S, v \in E_p^\perp\} \subseteq TM|_S$$

is a subbundle and there exists a smooth orthonormal local frame for $E^\perp$ around each point in S .

Proof. Choose any local frame

$$\sigma_1, \dots, \sigma_k: U \rightarrow E.$$

After possibly shrinking U , we can extend this frame to a frame

$$\sigma_1, \dots, \sigma_n: U \rightarrow TM|_S$$

of $TM|_S$. Write $\sigma_i(p) = (p, X_p^i)$. Applying Gram-Schmidt to the vectors X_p^i yields an orthonormal basis $Y_p^1, \dots, Y_p^n$ of $T_p M$ for each $p \in U$:

$$Y_p^i := \frac{X_p^i - \sum_{j=1}^{i-1} \langle X_p^i, Y_p^j \rangle Y_p^j}{\|X_p^i - \sum_{j=1}^{i-1} \langle X_p^i, Y_p^j \rangle Y_p^j\|}$$

Since this formula is smooth, we can define the smooth sections

$$\tilde{\sigma}_i(p) := (p, Y_p^i).$$

Notice that $\tilde{\sigma}_i(p) \in E^\perp$ for $i \geq k+1$ and $E_p^\perp = \text{span}\{Y_p^{k+1}, \dots, Y_p^n\}$. This shows the statement. $\square$

Given a Riemannian manifold M and a submanifold $S \subseteq M$, we can define the **normal bundle** of S :

$$NS := \{(p, v) \in TM \mid p \in S, v \in T_p S^\perp\}.$$

Corollary 7.30. *Let M be a Riemannian manifold, $S \subseteq M$ a submanifold. Then NS is a subbundle of $TM|_S$.*

Proof. We only need to show, that TS is a smooth subbundle of $TM|_S$ and then apply Lemma 7.29. To this end, choose a local frame (σ_i) of TS . Then $\iota \circ \sigma_i$ are sections of $TM|_S$, where $\iota: TS \hookrightarrow TM|_S$ is the inclusion. So we only have to verify, that ι is smooth. But this can easily be seen in local trivialisations of TS and $TM|_S$. $\square$

7.5 Universal Bundles

This section is mainly based on [Ebe12]. Other sources are [Mit11] and [td08].

Definition 7.31. A G -principal bundle $E \rightarrow B$ is called universal if E is contractible as a topological space.

The significance of this definition is established by the following theorem. We say that two morphisms of G -principal bundles are G -homotopic if they are homotopic through morphisms of G -principal bundles.

Theorem 7.32. *Let $E \rightarrow B$ be a universal G -principal bundle. Then for any other G -principal bundle $P \rightarrow X$ over a CW-complex X there exists a morphism $P \rightarrow E$ of G -principal bundles, which is unique up to G -homotopy.*

Before showing the Theorem let us discuss its implications. Firstly if we assume, that B is also a CW complex (which we may always do, see Remark 7.36) we can equivalently formulate the Theorem by saying that $E \rightarrow B$ is a terminal object in the homotopy category of G -principal bundles over CW -complexes and thus in particular uniquely determined (up to G -homotopy equivalence). This justifies writing $EG := E$ and $BG := B$. Since any bundle morphism $(F, f): P \rightarrow EG$ induces a morphism $P \rightarrow f^*EG$ (and vice versa, which is true for arbitrary fiber bundles) we moreover conclude from Proposition 7.22 that

$$P \cong f^*EG.$$

So by the homotopy invariance of the pullback we get a well-defined bijection

$$\phi_X: [X, BG] \rightarrow \mathcal{P}_G(X), [f] \mapsto [f^*EG],$$

where $\mathcal{P}_G(X)$ denotes the set of isomorphism classes of G -principal bundles over X . To see that ϕ_X is indeed bijective note that the existence result in Theorem 7.32 shows that ϕ_X is surjective and the uniqueness result shows that it is injective. Moreover ϕ_X is clearly natural in X . Let us note this important corollary.

Corollary 7.33. *The maps ϕ_X define a natural isomorphism of contravariant functors*

$$[-, BG], \mathcal{P}_G(-): \mathbf{HCW} \rightarrow \mathbf{Set},$$

where $\mathbf{HCW}$ denotes the homotopy category of CW -complexes.

Theorem 7.32 will be an application of the following Theorem.

Theorem 7.34. *Let $P \rightarrow X$ be a G -principal bundle over a CW -complex and $A \subseteq X$ a subcomplex. Then any morphism $P|_A \rightarrow EG$ can be extended to a morphism $P \rightarrow EG$.*

Proof. We follow the outline given in [Ebe12] but add quite a few details on the explicit construction and especially show that the constructed extension is in fact continuous. Let $F: P|_A \rightarrow EG$ be a given morphism. By a standard argument using Zorn's Lemma it suffices to show the statement in the case where X is obtained from A by attaching a single cell, that is we have a pushout

$$\begin{array}{ccc} S^{n-1} & \longrightarrow & A \\ \downarrow & & \downarrow \\ D^n & \xrightarrow{\psi} & X \end{array}$$

Using homotopy invariance of the pullback we get that ψ^*P is trivial. Let $\tau: D^n \times G \rightarrow \psi^*P$ be such a trivialisation and let $\Psi: \psi^*P \rightarrow P$ be the canonical map (cf. paragraph on pullback bundles). Then by equivariance the morphism

$$S^{n-1} \times G \xrightarrow{\tau} \psi^*P|_{S^{n-1}} \xrightarrow{\Psi} P|_A \xrightarrow{F} EG$$

is given by $(x, g) \mapsto \phi(x) \cdot g$ for some continuous map $\phi: S^{n-1} \rightarrow EG$. Since $\pi_{n-1}(EG) = 0$ we can extend ϕ to a map $\phi': D^n \rightarrow EG$ and thus we can extend

$$\psi^*P|_{S^{n-1}} \xrightarrow{\Psi} P|_A \xrightarrow{F} EG$$

to a morphism $H: \psi^*P \rightarrow EG$ by setting

$$H(\tau(x, g)) := \phi'(x) \cdot g.$$

With this we can now define an extension $F': P \rightarrow EG$ of F . To this end let $B^n := D^n - S^{n-1}$ be the open ball and note that $\psi|_{B^n}$ is an embedding and so in particular does Ψ define an isomorphism $\psi^*P|_{B^n} \rightarrow P|_{\psi(B^n)}$ (see Lemma 7.12). Now set

$$F'(y) := \begin{cases} H(\Psi^{-1}(y)) & , y \in P|_{\psi(B^n)}, \\ F(y) & , \text{otherwise.} \end{cases}$$

Clearly F' is equivariant and so it only remains to show continuity. For this let $y_n \in P$ be a sequence converging to some $y \in P$. Since $P|_{\psi(S^{n-1})} \subseteq P$ is closed the only non-trivial case is

wehn $y \in P|_{\psi(S^{n-1})}$ and infinitely many $y_n \in P|_{\psi(B^n)}$. Without loss of generality we may then of course assume that $y_n \in P|_{\psi(B^n)}$ for all n . We now claim that (after passing to a subsequence) the sequence

$$y'_n := \Psi^{-1}(y_n)$$

converges to some $y' \in \psi^*P$. Then $\Psi(y') = y$, $y' \in \psi^*P|_{S^{n-1}}$ and thus

$$F'(y_n) = H(y'_n) \longrightarrow H(y') = F(\Psi(y')) = F(y),$$

such that continuity of F' follows. To show the claim let $p: \psi^*P \rightarrow D^n$ and $\pi: P \rightarrow X$ be the bundle projections. By compactness of D^n we may assume that the sequence

$$x'_n := p(y'_n) \in D^n$$

converges to some $x' \in D^n$. Now choose some local trivialisation $\kappa: U \times G \rightarrow P$ around y . Then the y_n correspond to points

$$\kappa^{-1}(y_n) = (x_n, g_n) \in U \times G,$$

where $x_n = \psi(x'_n) = \pi(y_n)$. By definition of the pullback, κ induces a trivialisation

$$\kappa': \psi^{-1}(U) \times G \rightarrow \psi^*P$$

with $(\kappa')^{-1}(y'_n) = (x'_n, g_n)$. Note that for n sufficiently large $p(y'_n) = x'_n \in \psi^{-1}(U)$, since we assumed that this sequence converges to $x' \in D^n$ and $x := \psi(x') = \pi(y) \in U$. So we can in fact assume that $y'_n \in \text{im}(\kappa')$. Since $y_n \longrightarrow y$ we get that

$$(x_n, g_n) \longrightarrow (x, g) := \kappa^{-1}(y)$$

and so

$$(x'_n, g_n) \longrightarrow (x', g).$$

Thus $y'_n \longrightarrow \kappa'(x', g)$, showing the claim. $\square$

Proof of Theorem 7.32. Applying Theorem 7.34 to $A = \emptyset$ shows the existence of a morphism $P \rightarrow EG$. Now suppose we are given morphisms $F_0, F_1: P \rightarrow EG$. Consider the G -principal bundle

$$\pi \times \text{id}: P \times [0, 1] \rightarrow X \times [0, 1],$$

where $\pi: P \rightarrow X$ is the bundle projection (this is just the pullback bundle of P under the projection $X \times [0, 1] \rightarrow X$). Then F_0 and F_1 define a morphism of G -principal bundles

$$(F_0, F_1): P \times \{0, 1\} = (P \times [0, 1])|_{X \times \{0, 1\}} \rightarrow EG,$$

which by Theorem 7.34 can be extended to $P \times [0, 1]$, thus yielding a G -homotopy between F_0 and F_1 . $\square$

For any topological group G a universal bundle always exists. This can for example be shown using the Milnor construction, which does in fact construct universal bundles in a functorial way.

Theorem 7.35 (Milnor Construction). *For any topological group G there exists a universal G -principal bundle. Moreover this construction is functorial, such that we get a functor*

$$\text{TopGrp} \rightarrow \text{Prin},$$

from the category of topological groups to the category of principal bundles, which is given on objects by $G \mapsto (EG \rightarrow BG)$.

Proof. The construction of $EG \rightarrow BG$ is shown in [Ebe12] and we shall only indicate how to construct a morphism $EG \rightarrow EH$ from a homomorphism $\phi: G \rightarrow H$. Recall that EG was defined as the set of formal convex combinations of group elements with a suitable topology. That is an element in EG is of the form

$$\sum_{i=1}^{\infty} t_i g_i$$

for $t_i \in [0, 1]$ with only finitely many t_i not being equal to zero and $\sum_{i=1}^{\infty} t_i = 1$. Then

$$E\phi: EG \rightarrow EH, \quad \sum_{i=1}^{\infty} t_i g_i \mapsto \sum_{i=1}^{\infty} t_i \phi(g_i)$$

defines a continuous map which is ϕ -invariant and hence descends to a continuous map

$$B\phi: BG = EG/G \rightarrow BH = EH/H.$$

Then $(E\phi, B\phi)$ is a morphism of principal bundles and clearly this construction is functorial. $\square$

Remark 7.36. The Milnor construction may not yield a CW-complex BG (at least I don't think it does in general) but in [Mit11] it is shown that the existence of a universal G -principal bundle always implies the existence of another universal G -principal bundle over a CW-complex. The construction of universal bundles over CW-complexes can also be made functorial (though I do not have a clear reference for this).

Remark 7.37. Let $\phi: G \rightarrow H$ be a homomorphism of topological groups. Assuming that BG and BH are CW-complexes, we may also construct $B\phi: BG \rightarrow BH$ more abstractly as follows. Note that for any CW-complex X we have a natural map

$$[X, BG] \xrightarrow{\cong} \mathcal{P}_G(X) \xrightarrow{\times_{\phi}^H} \mathcal{P}_H(X) \xrightarrow{\cong} [X, BH],$$

where the isomorphisms are the natural isomorphisms from Corollary 7.33 and the middle morphism is the natural transformation from Proposition 7.26. Then for $X = BG$, the identity on BG corresponds to a map $B\phi: BG \rightarrow BH$, which is characterized by

$$(B\phi)^* EH \cong EG \times_\phi H.$$

To see that this definition agrees with the one made in Theorem 7.35 we have to show that the $B\phi$ constructed there satisfies this characteristic property. This is done by again using the uniqueness result from Theorem ??: If $U_\alpha, U_\beta \subseteq BG$ are trivialising neighbourhoods for EG as constructed in the Milnor construction (see [Ebe12]) with transition function $\tau_{\alpha,\beta}$, then $EG \times_\phi H$ will also be trivial over U_α, U_β with transition function $\phi \circ \tau_{\alpha,\beta}$ (cf. Proposition 7.26). Moreover $(B\phi)^*EH$ is also trivial over U_α, U_β , since EH is trivial over $B\phi(U_\alpha), B\phi(U_\beta)$ and a simple computation shows that the transition function is again $\phi \circ \tau_{\alpha,\beta}$.

Note however that the Milnor construction yields a map $BG \rightarrow BH$, while the alternative construction only yields a homotopy class of maps.

Using the theory developed above we can generalize the concept of a universal bundle to arbitrary fiber bundles. First let us define what a universal should be in this context.

Definition 7.38. A fiber bundle $E \rightarrow X$ over a CW-complex X with fiber F and structure group G is called universal if for any other fiber bundle $E' \rightarrow X'$ over a CW-complex X' with the same fiber and structure group, there exists a G -morphism $E' \rightarrow E$, which is unique up to homotopy of G -morphisms.

For principal bundles we called a bundle universal if the total space is contractible, but showed in Theorem 7.32 that such bundles are precisely the universal bundles in the sense of the definition above. Thus this definition generalizes our previous one. Again we remark that Definition 7.38 is equivalent to a saying that universal bundles are the terminal objects in the appropriate homotopy category. Using the theory developed above it is easy to show existence of universal bundles:

Theorem 7.39. *Let G be a topological group acting effectively on F . Then the bundle $EG \times_G F$ is universal.*

Proof. ???

Moreover we have an analogue of Theorem 7.34:

Theorem 7.40. *Let $E \rightarrow X$ be a fiber bundle with fiber F and structure group G over a CW-complex and let $A \subseteq X$ be a subcomplex. Then any G -morphism $E|_A \rightarrow EG \times_G F$ can be extended to a G -morphism $E \rightarrow EG \times_G F$.*

Proof. ???

□

Corollary 7.41. *Let G be a topological group acting effectively on a space F . Then for any CW-complex X the map*

$$[X, BG] \rightarrow \mathcal{F}_{F,G}(X), \quad [f] \mapsto [f^*(EG \times_G F)]$$

is a natural bijection, where $\mathcal{F}_{F,G}(X)$ denotes the set of isomorphism classes of fiber bundles with fiber F and structure group G over X .

Proof. We could argue as we did in Corollary 7.33 for principal bundles using Theorem 7.39 instead. Alternatively we can also show this by using Corollary 7.33 directly, which tells us that the map

$$[X, BG] \rightarrow \mathcal{P}_G(X), \quad [f] \mapsto [f^*EG]$$

is a natural bijection and by Proposition 7.18

$$\mathcal{P}_G(X) \rightarrow \mathcal{F}_{F,G}(X), \quad [P] \mapsto [P \times_G F]$$

is a natural bijection. Composing both thus yields the natural bijection

$$[X, BG] \rightarrow \mathcal{F}_{F,G}(X), \quad [f] \mapsto [(f^*EG) \times_G F].$$

So the statement follows by again applying Proposition 7.18.

□

8 Construction of Diffeomorphisms

8.1 Existence of Smooth Isotopies

In this section we proof a few statements on the existence of smooth isotopies. Lemma 8.1 and Theorem 8.2 follow [Mil65] quite closely. The first Lemma is a slight modification of Lemma 5.7 in [Mil65].

Lemma 8.1 (Isotopy Lemma). *Let $h: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an orientation preserving embedding with $h(0) = 0$. Then h is isotopic to the identity $\text{rel } \{0\}$.*

Proof. The proof is basically the same as in [Mil65]. Since $h(0) = 0$ we can write

$$h(x) = x_1 h_1(x) + \dots + x_n h_n(x)$$

for smooth functions $h_i: \mathbb{R}^n \rightarrow \mathbb{R}^n$ and then define an isotopy

$$\frac{1}{t} h(tx) = x_1 h_1(tx) + \dots + x_n h_n(tx), \quad 0 \leq t \leq 1$$

from the linear map

$$A: \mathbb{R}^n \rightarrow \mathbb{R}^n, x \mapsto x_1 h_1(0) + \dots + x_n h_n(0)$$

to h . Since $\det(A) > 0$ and the space of all $n \times n$ matrices with positive determinant is path connected we moreover get an isotopy from A to id . $\square$

Next we proof Theorem 5.8 in [Mil65] and give a slightly stronger formulation of it. The main idea behind the proof is beautifully explained in [Mil65].

Theorem 8.2 (Isotopy Extension Theorem). *Let $M \subseteq N$ be a submanifold with or without boundary of the boundaryless manifold N and let $h_t: M \rightarrow N, 0 \leq t \leq 1$ be an isotopy of the inclusion $M \hookrightarrow N$. Given any compact subset $K \subseteq M$ and an open set $U \subseteq N$, such that $h_t(K) \subseteq U$ for all $t \in [0, 1]$, there exists an isotopy*

$$\bar{h}_t: N \rightarrow N, 0 \leq t \leq 1$$

with the following properties

- i) $\bar{h}_0 = \text{id}_N$
- ii) $\bar{h}_t(x) = h_t(x)$ for all $x \in K$ and $t \in [0, 1]$
- iii) $\bar{h}_t(x) = x$ for all $x \in N \setminus U$ and $t \in [0, 1]$

Proof. We formulate the proof only for the case $\partial M = \emptyset$, see Remark 8.3. Let $V \subseteq M$ be a relatively compact neighbourhood of K . Then

$$\begin{aligned} H: V \times [0, 1] &\rightarrow N \times [0, 1], \\ (x, t) &\mapsto (h_t(x), t) \end{aligned}$$

is an embedding, since the map $M \times [0, 1] \rightarrow N \times [0, 1]$ defined by the same formula is an injective immersion. So H defines a submanifold

$$S := \text{im}(H) \subseteq N \times [0, 1]$$

as well as the smooth vector field $X := H_*\left(\frac{\partial}{\partial t}\right)$ on S , i.e. for $y = h_t(x)$ we have

$$X_{(y,t)} := D_{(x,t)} H \left(\frac{\partial}{\partial t} \right) = \left(\frac{\partial h_t}{\partial t}(x), \frac{\partial}{\partial t} \right).$$

We can then extend X to some open neighbourhood $W \subseteq U \times [0, 1] \subseteq N \times [0, 1]$ of S at first arbitrarily and then set the coordinate of the part pointing in the $\frac{\partial}{\partial t}$ direction equal to 1. Thus we obtain a smooth vector field Y on W of the form

$$Y_{(y,t)} = (\tilde{X}_{(y,t)}, \frac{\partial}{\partial t})$$

which agrees with X on S . Choose a cut-off function

$$\psi: N \times [0, 1] \rightarrow \mathbb{R}$$

which equals 1 on $H(K \times [0, 1])$ and has its support contained in W . Then we consider the smooth vector field Z on $N \times [0, 1]$ defined by

$$Z := \psi Y + (1 - \psi) \frac{\partial}{\partial t}.$$

The idea now is to define the required isotopy $\bar{h}_t$ using the flow of Z . For this we want to assume that U is relatively compact, otherwise we may shrink U . The reason for this will become clear shortly, but first let us record the crucial properties of Z :

i) Z is of the form

$$Z_{(y,t)} = (\tilde{Z}_{(y,t)}, \frac{\partial}{\partial t}),$$

ii) Z agrees with X on $H(K \times [0, 1])$,

iii) $Z = \frac{\partial}{\partial t}$ outside of the compact set $\bar{U} \times [0, 1]$.

We claim that for any $y \in N$ the unique maximal integral curve ψ_y of Z starting at $(y, 0) \in N \times [0, 1]$ is defined for all $t \in [0, 1]$. If $y \notin \bar{U}$ this is clear, since the ψ_y is explicitly given by

$$\psi_y(t) = (y, t), \quad 0 \leq t \leq 1.$$

For $y \in \bar{U}$ this follows from the fact, that ψ_y has to be contained in the compact set $\bar{U} \times [0, 1]$ for all time, where it is defined (due to uniqueness of integral curves and the case $y \notin \bar{U}$ above). Then using property (i) of Z we see that $\psi_y(t)$ is in fact defined for all $t \in [0, 1]$.

So after extending Z to $N \times \mathbb{R}$ (to avoid dealing with flows on manifolds with boundary) we can consider the flowout of $N \times \{0\}$, which is defined on $N \times [0, 1]$ by the preceding observations and hence defines a level-preserving diffeomorphism

$$\begin{aligned} \bar{H}: N \times [0, 1] &\rightarrow N \times [0, 1], \\ (y, t) &\mapsto \psi_y(t). \end{aligned}$$

Finally we define the required isotopy $\bar{h}_t$ by

$$\bar{H}(y, t) = (\bar{h}_t(y), t).$$

It remains to show that $\bar{h}_t$ agrees with h_t on K . But this follows from property (ii) of Z : for $x \in K$ let $\phi_x(t) := (x, t)$, $0 \leq t \leq 1$ be the integral curve of the vector field $\frac{\partial}{\partial t}$ on $M \times [0, 1]$. Then

$$H(x, t) = H(\phi_x(t)) = \psi_x(t) = \bar{H}(x, t).$$

The theorem follows. □

Remark 8.3. If M is a manifold with boundary we can define a smooth isotopy as a map

$$h: M \times [0, 1] \rightarrow N,$$

which extends to a smooth isotopy

$$\bar{h}: M \times \mathbb{R} \rightarrow N.$$

Then exactly the same proof can be carried out if M is allowed to have boundary by constructing the required vector fields on $N \times \mathbb{R}$ and then restricting everything back to $N \times [0, 1]$. Note that this definition of smooth isotopies does not change the isotopy classes of maps in the case $\partial M = \emptyset$ (cf. Lemma 12.6). Of course using the notion of a manifold with edges we can define smooth isotopies in the same way and the proof of Theorem 8.2 goes through unchanged.

Together the Isotopy Lemma and the Isotopy Extension Theorem can be used to extend locally defined diffeomorphisms to the whole manifold as follows: Begin with an orientation preserving embedding $\phi: U \rightarrow M$ of an open subset $U \subseteq M$. Then - after possibly restricting ϕ to a smaller domain - use Lemma 8.1 to construct an isotopy from ϕ to the inclusion $U \hookrightarrow M$. Lastly - after once again restricting the isotopy to a smaller domain - we can use Theorem 8.2 to extend this isotopy to an isotopy $\bar{h}_t: N \rightarrow N$ starting at id_N . Then $h_1: N \rightarrow N$ is a diffeomorphism extending ϕ .

Next we state a few interesting corollaries of Theorem 8.2.

Corollary 8.4. *Let M be a manifold without boundary, $p, q \in M$ and $\gamma: [0, 1] \rightarrow M$ a smooth path from p to q . Then there exists a diffeomorphism $\phi: M \rightarrow M$ with $\phi(p) = q$, which is smoothly isotopic to id_M . Moreover this isotopy can be chosen to be stationary outside a given open neighbourhood $U \subseteq M$ of the image of γ .*

Proof. The smooth path γ can be seen as an isotpy of the inclusion $\{p\} \hookrightarrow M$. Then the statement follows directly from Theorem 8.2. $\square$

Remark 8.5. If $\text{Aut}_{\text{id}}(M) \subset \text{Aut}(M)$ denotes the subgroup of diffeomorphisms which are smoothly isotopic to id_M , the above corollary in particular implies that $\text{Aut}_{\text{id}}(M)$ acts transitively on M .

We use Corollary 8.4 to show a corollary from [GP74].

Corollary 8.6. *Let M be a connected manifold without boundary of dimension at least 2. Given distinct points $x_1, \dots, x_k, y_1, \dots, y_k \in M$, there exists a diffeomorphism*

$$\psi: M \rightarrow M$$

with $\psi(x_i) = y_i$, which is smoothly isotopic to the identity. Moreover, we can choose this isotopy to be compactly supported.

Proof. We proof the statement inductively. For $k = 0$ there is nothing to show. So suppose we already have a diffeomorphism $\psi': M \rightarrow M$ smoothly isotopic to the identity, where the isotopy is compactly supported and $\psi'(x_i) = y_i$ for $1 \leq i \leq k-1$. Note that $M - \{y_1, \dots, y_{k-1}\}$ is connected, so Corollary 8.4 implies the existence of a diffeomorphism $\phi: M \rightarrow M$ with $\phi(\psi'(x_k)) = y_k$ and $\phi(y_i) = y_i$ for $1 \leq i \leq k-1$, such that ϕ is smoothly isotopy to the identity via a compactly supported isotopy. Now set $\psi := \phi \circ \psi'$ and compose the isotopies to get a compactly supported isotopy from ψ to the identity. $\square$

This next corollary essentially says, that there are many embeddings of small dimensional manifolds into large dimensional manifolds.

Corollary 8.7. *Let M be a connected manifold without boundary and S a manifold with or without boundary, such that there exists an embedding $S \hookrightarrow \mathbb{R}^{\dim(M)}$. Moreover we fix two distinct points $s_0, s_1 \in S$. Then for any two points $p, q \in M$ there exists an embedding $\iota: S \hookrightarrow M$, such that $\iota(s_0) = p$ and $\iota(s_1) = q$.*

Proof. By assumption we can find an embedding $\iota': S \hookrightarrow U \subseteq M$ into any coordinate neighbourhood U around p , such that $\iota'(s_0) = p$. We may assume $p \neq q$, so we can find a smooth path from $\iota'(s_1)$ to q , that does not hit p (if M is 1-dimensional we may arrange the embedding ι' in such a way, that $\iota'(s_1)$ lies in the same connected component of the manifold $M - \{p\}$ as q). Then Corollary 8.4 implies the existence of a diffeomorphism $\phi: M \rightarrow M$ with $\phi(\iota'(s_1)) = q$ and $\phi(p) = p$. Now set $\iota := \phi \circ \iota'$. $\square$

The above corollary should also be true if M is allowed to have boundary, but the proof fails, since Corollary 8.4 is clearly false if M has non-empty boundary. An interesting consequence of Corollary 8.7 is the existence of **embedded paths**: For any connected manifold M (without boundary) and any $p, q \in M$ there exists an embedding $\gamma: [0, 1] \rightarrow M$ with $\gamma(0) = p, \gamma(1) = q$.

By the Whitney embedding theorem any n -dimensional manifold can be embedded into $\mathbb{R}^{2n+1}$, but there might be different non-isotopic embeddings. Think for example of inequivalent knots in $\mathbb{R}^3$. The following theorem however says, that upon further increasing the dimension by one, there is only one embedding $M \hookrightarrow \mathbb{R}^{2n+2}$ up to isotopy (at least for M closed).

Theorem 8.8. *Let M be a closed n -dimensional manifold. Then any two embeddings $M \hookrightarrow \mathbb{R}^{2n+k}$ for $k \geq 2$ are isotopic.*

Proof. Let $N := 2n + k$ and let $f, g: M \hookrightarrow \mathbb{R}^N$ be two embeddings. The idea of the proof is to construct preliminary isotopies $f \simeq f'$ and $g \simeq g'$, such that the straight line homotopy

$$h_t := (1 - t)f' + tg'$$

is an isotopy. Let us observe what is necessary for h_t to be an isotopy. First of all injectivity of h_t for each time $t \in [0, 1]$ is equivalent to (note that for $t = 0, 1$ there is nothing to show)

$$(1 - t)(f'(x) - f'(y)) \neq t(g'(y) - g'(x))$$

for all $t \in (0, 1)$ and $x, y \in M$ with $x \neq y$. If we denote by $\Delta(X) \subseteq X \times X$ the diagonal of a manifold X this in turn is equivalent to

$$([f'(x) - f'(y)], [g'(x) - g'(y)]) \notin \Delta(\mathbb{R}P^{N-1}) \subseteq \mathbb{R}P^{N-1} \times \mathbb{R}P^{N-1}.$$

But this precisely means that the map

$$\begin{aligned} M \times M - \Delta(M) &\rightarrow \mathbb{R}P^{N-1} \times \mathbb{R}P^{N-1} \\ (x, y) &\mapsto ([f'(x) - f'(y)], [g'(x) - g'(y)]) \end{aligned}$$

is transverse to the diagonal $\Delta(\mathbb{R}P^{N-1})$, since

$$\begin{aligned} \dim(\Delta(\mathbb{R}P^{N-1})) + \dim(M \times M - \Delta(M)) &= 4n + k - 1 \\ &< 4n + 2(k - 1) = \dim(\mathbb{R}P^{N-1} \times \mathbb{R}P^{N-1}). \end{aligned}$$

Similarly injectivity of the differential of h_t is equivalent to

$$(1 - t)d_x f'(v) \neq -td_x g'(v)$$

for all $t \in (0, 1)$ and $(x, v) \in TM - M$, where we embed $M \subseteq TM$ as the zero section. Again we can formulate this as a transversality condition, namely that the map

$$\begin{aligned} TM - M &\rightarrow \mathbb{R}P^{N-1} \times \mathbb{R}P^{N-1} \\ (x, v) &\mapsto ([d_x f'(v)], [d_x g'(v)]) \end{aligned}$$

is transversal to the diagonal $\Delta(\mathbb{R}P^{N-1})$. In order to find such maps f' and g' we will be using the parametric transversality theorem. More precisely we will show that for almost all matrices $A, B \in \text{GL}(N)$ the maps

$$f' := A \circ f, \quad g' := A \circ g$$

satisfy the desired transversality conditions. Note that for $A, B \in \text{GL}^+(N)$ we have

$$f \simeq A \circ f, \quad g \simeq A \circ g,$$

since $\text{GL}^+(N)$ is the path connected component of $\text{GL}(N)$ containing the identity. Now consider the maps

$$\begin{aligned} \Phi: (M \times M - \Delta(M)) \times \text{GL}(N) \times \text{GL}(N) &\rightarrow (\mathbb{R}^N - 0) \times (\mathbb{R}^N - 0) \\ (x, y, A, B) &\mapsto (A(f(x) - f(y)), B(g(x) - g(y))) \end{aligned}$$

and

$$\begin{aligned} \Psi: (TM - M) \times \text{GL}(N) \times \text{GL}(N) &\rightarrow (\mathbb{R}^N - 0) \times (\mathbb{R}^N - 0) \\ (x, v, A, B) &\mapsto (A d_x f(v), B d_x g(v)). \end{aligned}$$

To compute the derivatives let us write L_w for the map $L_w: \mathbb{R}^{N \times N} \rightarrow \mathbb{R}^N$, $S \mapsto Sw$ for $w \in \mathbb{R}^N$, which is clearly surjective for $w \neq 0$. Then

$$d_{(x,y,A,B)} \Phi = \begin{pmatrix} * & L_{f(x)-f(y)} & 0 \\ * & 0 & L_{g(x)-g(y)} \end{pmatrix}$$

as a map

$$T_{(x,y)}(M \times M - \Delta(M)) \oplus \mathbb{R}^{N \times N} \oplus \mathbb{R}^{N \times N} \rightarrow \mathbb{R}^N \oplus \mathbb{R}^N,$$

where we denote by $*$ the entries irrelevant for our discussion. Since $f(x) - f(y), g(x) - g(y) \neq 0$ it follows that this map is surjective and thus Φ is a submersion. In particular, post composing with the projection

$$\pi: (\mathbb{R}^N - 0) \times (\mathbb{R}^N - 0) \rightarrow \mathbb{R}P^{N-1} \times \mathbb{R}P^{N-1}$$

shows that $\pi \circ \Phi$ is a submersion. Essentially the same calculation also shows that $\pi \circ \Psi$ is a submersion. Now the theorem follows from the parametric transversality theorem. $\square$

As a corollary of the above theorem and the isotopy extension theorem we can show:

Corollary 8.9. *Let W be an n -dimensional manifold with compact boundary. Then any embedding $\partial W \hookrightarrow \mathbb{R}^{2n+1}$ can be extended to an embedding $W \hookrightarrow \mathbb{R}^{2n+1}$.*

Proof. By the Whitney embedding theorem we can choose an embedding $\phi: W \hookrightarrow \mathbb{R}^{2n+1}$. Let $f := \phi|_{\partial W}$ be the restriction to the boundary and let $g: \partial W \hookrightarrow \mathbb{R}^{2n+1}$ be any other embedding. By Theorem 8.8 there exists an isotopy from the inclusion $f(\partial W) \hookrightarrow \mathbb{R}^{2n+1}$ to the embedding

$$g \circ f^{-1}: f(\partial W) \hookrightarrow \mathbb{R}^{2n+1}$$

and by the isotopy extension Theorem (Theorem 8.2) we can extend this isotopy to an isotopy $h_t: \mathbb{R}^{2n+1} \rightarrow \mathbb{R}^{2n+1}$. Then $\psi := h_1 \circ \phi: W \hookrightarrow \mathbb{R}^{2n+1}$ is an extension of g . $\square$

9 Some Linear Algebra

9.1 General Statements

Lemma 9.1. *Let $T: V \rightarrow W$ be a linear map of finite dimensional vector spaces and let A be the matrix representation of T with respect to some basis. Then the matrix representation of the dual map $T^*: W^* \rightarrow V^*$ with respect to the corresponding dual basis is given by A^T .*

Proof. Let $X: \mathbb{R}^n \rightarrow V$ and $Y: \mathbb{R}^m \rightarrow W$ be linear isomorphisms given by the chosen basis, i.e.

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \uparrow X & & \uparrow Y \\ \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \end{array}$$

commutes. Moreover, let $\phi: \mathbb{R}^n \rightarrow (\mathbb{R}^n)^*$, $\psi: \mathbb{R}^m \rightarrow (\mathbb{R}^m)^*$ be the canonical isomorphisms induced by the standard basis $e_1, \dots, e_n$ and $d_1, \dots, d_m$ on $\mathbb{R}^n$ and $\mathbb{R}^m$, respectively. We then get the following commutative diagram

$$\begin{array}{ccc} V^* & \xleftarrow{T^*} & W^* \\ \downarrow X^* & & \downarrow Y^* \\ (\mathbb{R}^n)^* & \xleftarrow{A^*} & (\mathbb{R}^m)^* \\ \uparrow \phi & & \uparrow \psi \\ \mathbb{R}^n & \xleftarrow{B} & \mathbb{R}^m \end{array}$$

where B is the desired matrix representation of T^* . Write $B = (b_{ij})$. Then b_{ij} is given as the i -th component of the vector Bd_j . By the above diagram this is precisely

$$b_{ij} = (d_j^* \circ A)(e_i) = a_{ji},$$

i.e. $B = A^T$. □

Lemma 9.2. *Let $V, V_1, \dots, V_k$ be finite dimensional vector spaces. There exist natural isomorphisms*

$$\begin{aligned} V_1^* \otimes \dots \otimes V_k^* &\xrightarrow{\cong} (V_1 \otimes \dots \otimes V_k)^* \\ \Lambda^k V^* &\xrightarrow{\cong} (\Lambda^k V)^* \\ \Sigma^k V^* &\xrightarrow{\cong} (\Sigma^k V)^* \end{aligned}$$

where $\Lambda^k V$ denotes the k -th exterior power and $\Sigma^k V$ the k -th symmetric power (see section 9.4).

Proof. Define the first isomorphism $V_1^* \otimes \dots \otimes V_k^* \rightarrow (V_1 \otimes \dots \otimes V_k)^*$ by

$$\phi_1 \otimes \dots \otimes \phi_k \mapsto \left(v_1 \otimes \dots \otimes v_k \mapsto \prod_{i=1}^k \phi_i(v_i) \right).$$

using the universal property of the tensor product. This map is clearly injective, hence indeed an isomorphism. Moreover this map induces the second isomorphism $\Lambda^k V^* \rightarrow (\Lambda^k V)^*$ by

$$\begin{array}{ccc} (V^*)^{\otimes k} & \xrightarrow{\cong} & (V^{\otimes k})^* \\ \uparrow & & \downarrow \\ \Lambda^k V^* & \longrightarrow & (\Lambda^k V)^* \end{array}$$

where the vertical maps are (induced by) the inclusions $\Lambda^k V^* \rightarrow (V^*)^{\otimes k}$ and $\Lambda^k V \rightarrow V^{\otimes k}$. To see that this does in fact define an isomorphism note that an inverse is explicitly given by

$$\begin{array}{ccc} (V^*)^{\otimes k} & \xrightarrow{\cong} & (V^{\otimes k})^* \\ \text{Alt} \downarrow & & \uparrow \text{Alt}^* \\ \Lambda^k V^* & \longleftarrow & (\Lambda^k V)^* \end{array}$$

where Alt denotes the antisymmetrization map (again see section 9.4). The third isomorphism $\Sigma^k V^* \rightarrow (\Sigma^k V)^*$ is defined analogously using the symmetrization map instead. Naturality of the defined isomorphisms follows directly from the definition by a straightforward computation. $\square$

9.2 Multilinear Algebra

Here are two statements about multilinear maps, which are generalizations of the corresponding statements for linear maps.

Lemma 9.3. *Let $V_1, \dots, V_k$ and W be normed vector spaces and*

$$A: V_1 \times \dots \times V_k \rightarrow W$$

be a multilinear map, then A is continuous (with respect to the product norm on $V_1 \times \dots \times V_k$) if and only if it is bounded, i.e. there exists $C > 0$ with

$$\|A(v_1, \dots, v_k)\| \leq C \prod_{i=1}^k \|v_i\|$$

for all $v_i \in V_i$.

Proof. The proof is analogous to the linear case. First assume that A is continuous, then in particular there exists $\delta > 0$, such that

$$\sum_{i=1}^k \|v_i\| \leq k\delta \quad \Rightarrow \quad \|A(v_1, \dots, v_k)\| \leq 1.$$

So for arbitrary $v_i \in V_i$, we have

$$\|A(v_1, \dots, v_k)\| = \frac{1}{\delta^k} \prod_{i=1}^k \|v_i\| \|A(\frac{\delta v_1}{\|v_1\|}, \dots, \frac{\delta v_k}{\|v_k\|})\| \leq \frac{1}{\delta^k} \prod_{i=1}^k \|v_i\|$$

Now assume, that A is bounded. Given any $v_i, w_i \in V_i$ we compute:

$$\begin{aligned} \|A(v_1, \dots, v_k) - A(w_1, \dots, w_k)\| &= \left\| \sum_{i=1}^k A(w_1, \dots, w_{i-1}, v_i - w_i, v_{i+1}, \dots, v_k) \right\| \\ &\leq C \sum_{i=1}^k \|w_1\| \cdots \|w_{i-1}\| \|v_i - w_i\| \|v_{i+1}\| \cdots \|v_k\| \end{aligned}$$

Since the left hand side vanishes for $v_i \rightarrow w_i$ the claim follows. $\square$

Lemma 9.4. *Multilinear maps on finite dimensional vector spaces are continuous (with respect to any norm).*

Proof. For ease of notation we only proof the Lemma for bilinear maps, but the proof is completely analogous for the general case. So let $A: V \times W \rightarrow Z$ be a bilinear map for finite dimensional vector spaces V, W and let $e_1, \dots, e_k$ and $d_1, \dots, d_m$ be a basis for V and W , respectively. Given $v \in V$ and $w \in W$ write

$$v = \sum_{i=1}^k v^i e_i, \quad w = \sum_{j=1}^m w^j d_j.$$

Define $C := \max_{i,j} \|A(e_i, d_j)\|$, then

$$\|A(v, w)\| \leq \sum_{i,j} |v^i| |w^j| \|A(e_i, d_j)\| \leq C \sum_{i,j} |v^i| |w^j| = C \left(\sum_i |v^i| \right) \left(\sum_j |w^j| \right).$$

Note that

$$\|v\| := \sum_i |v^i|$$

defines a norm, so A is bounded with respect to these norms on V, W , hence continuous by Lemma 9.3. Since all norms on finite dimensional vector spaces are equivalent, the statement follows. $\square$

The following lemma extends the universal property of the Tensor Product to multilinear maps.

Lemma 9.5. *For $i = 1, \dots, n$ let $V_1^i, \dots, V_{k_i}^i$ be vector spaces and*

$$\pi_i: V_1^i \times \dots \times V_{k_i}^i \rightarrow V_1^i \otimes \dots \otimes V_{k_i}^i$$

the canonical product maps. Then for every multilinear map $L: \prod_{i=1}^n V_1^i \times \dots \times V_{k_i}^i \rightarrow Z$ to a vector space Z , there exists a unique multilinear map $A: \prod_{i=1}^n V_1^i \otimes \dots \otimes V_{k_i}^i \rightarrow Z$ such that the following diagram commutes

$$\begin{array}{ccc} \prod_{i=1}^n V_1^i \times \dots \times V_{k_i}^i & \xrightarrow{L} & Z \\ \Pi_i \pi_i \downarrow & \nearrow A & \\ \prod_{i=1}^n V_1^i \otimes \dots \otimes V_{k_i}^i & & \end{array}$$

Proof. Consider the following commutative diagram

$$\begin{array}{ccccc} & \prod_{i=1}^n V_1^i \times \dots \times V_{k_i}^i & & & \\ & \downarrow \pi & \searrow L & & \\ \Pi_i \pi_i \swarrow & \otimes_{i=1}^n V_1^i \otimes \dots \otimes V_{k_i}^i & \xrightarrow{B} & & Z \\ & \uparrow p & \nearrow A & & \\ & \prod_{i=1}^n V_1^i \otimes \dots \otimes V_{k_i}^i & & & \end{array}$$

where p and π are the canonical product maps, the map B is induced by L by the universal property of the tensor product and A is defined as the composition of B and p . Clearly A is the unique map satisfying the statement of the lemma. $\square$

9.3 Symplectic Vector Spaces

Lemma 9.6. *Let (V, ω) be a symplectic vector space and $W \subseteq V$ an isotropic subspace. Then any basis of W can be extended to a symplectic basis of V .*

Proof. We proof this by induction over $\dim(V)$. If $\dim(V) = 2$ the statement follows directly from non-degeneracy of ω . Now assume the statement holds for $\dim(V) = 2(n-1)$ and let $\dim(V) = 2n$. Given a basis $u_1, \dots, u_k$ of W define

$$U := \text{span}(u_2, \dots, u_k) \subsetneq W$$

Then $W^\omega \subsetneq U^\omega$, hence there exists $v_1 \in U^\omega$ with $v_1 \notin W^\omega$, so by construction we must have

$$\omega(u_1, v_1) \neq 0.$$

After normalizing we may assume $\omega(u_1, v_1) = 1$. Now consider

$$Q := \text{span}(u_1, v_1).$$

Then Q is a symplectic subspace, thus

$$(Q^\omega, \sigma := \omega|_{Q^\omega})$$

again a symplectic vector space. Moreover $U \subseteq Q^\omega$ and

$$U \subseteq U^\sigma.$$

So by our induction hypothesis we can extend $u_2, \dots, u_k$ to a symplectic basis $u_2, \dots, u_n, v_2, \dots, v_n$ of (Q^ω, σ) . Then

$$u_1, \dots, u_n, v_1, \dots, v_n$$

is a symplectic basis of (V, ω) . $\square$

The following Lemma gives a useful characterisation of symplectic matrices analogous to orthogonal matrices.

Lemma 9.7. *A matrix $A \in \mathbb{R}^{2n \times 2n}$ is symplectic if and only if the column vectors of A form a symplectic basis of $(\mathbb{R}^{2n}, \omega_0)$.*

Proof. Write $A = (a_{x_1} \cdots a_{x_n} a_{y_1} \cdots a_{y_n})$ for $a_{x_i}, a_{y_i} \in \mathbb{R}^{2n}$. If A is symplectic we have

$$\omega_0(a_{x_i}, a_{y_j}) = \omega_0(Ae_{x_i}, Ae_{y_j}) = \omega_0(e_{x_i}, e_{y_j}) = \delta_{ij}$$

and similarly $\omega_0(a_{x_i}, a_{x_j}) = 0 = \omega_0(a_{y_i}, a_{y_j})$. For the other implication it suffices to show that $\omega_0(\cdot, \cdot) = \omega_0(A\cdot, A\cdot)$ holds on basis vectors, which follows from the same calculation. $\square$

9.4 The Exterior Algebra

This section is meant to give a quick overview of the definition and most essential properties of the exterior algebra Λ^*V of a finite dimensional vector space V (over some field $\mathbb{K}$). To define the k -fold exterior power $\Lambda^k V$ consider the permutation maps

$$P_\sigma: V^{\otimes k} \rightarrow V^{\otimes k}$$

for $\sigma \in S(k)$ ⁵ given by

$$P_\sigma(v_1 \otimes \cdots \otimes v_k) := \text{sgn}(\sigma) v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(k)}.$$

These are well-defined by the universal property of the tensor product. We then call a vector $\alpha \in V^{\otimes k}$ **alternating** if

$$P_\sigma(\alpha) = \alpha \quad \forall \sigma \in S(k)$$

and define $\Lambda^k V \subseteq V^{\otimes k}$ as the subspace of all alternating vectors. Given any vector in $V^{\otimes k}$ there is a canonical way to make it into an alternating one using the **antisymmetrization map**

$$\text{Alt}: V^{\otimes k} \rightarrow \Lambda^k V, \quad \alpha \mapsto \frac{1}{k!} \sum_{\sigma \in S(k)} P_\sigma(\alpha).$$

Note that this map is surjective and $\text{Alt}(\alpha) = \alpha$ if and only if α is alternating. Using the antisymmetrization map we can define the **exterior product** (or **wedge product**) as the bilinear map

$$\begin{aligned} \Lambda^k V \times \Lambda^l V &\rightarrow \Lambda^{k+l} V, \\ (\alpha, \beta) &\mapsto \alpha \wedge \beta := \frac{(k+l)!}{k!l!} \text{Alt}(\alpha \otimes \beta). \end{aligned}$$

Without proof we state the facts, that this product is associative and graded commutative. Moreover if $e_1, \dots, e_n \in V$ is a basis of V then

$$\{e_{i_1} \wedge \cdots \wedge e_{i_k} \mid 1 \leq i_1 \leq \cdots \leq i_k \leq n\}$$

is a basis of $\Lambda^k V$. If we identify $\Lambda^k V$ with the space of alternating maps $V^* \times \cdots \times V^* \rightarrow \mathbb{R}$, these basis vectors satisfy

$$(e_{i_1} \wedge \cdots \wedge e_{i_k})(\varepsilon^{j_1}, \dots, \varepsilon^{j_k}) = \begin{cases} 1 & , (i_1, \dots, i_k) = (j_1, \dots, j_k) \\ 0 & \text{otherwise} \end{cases}$$

where $\varepsilon^1, \dots, \varepsilon^n \in V^*$ is the dual basis of $e_1, \dots, e_n$ and $1 \leq j_1 < \cdots < j_k \leq n$ is any ordered tuple. This is the reason for the normalization factor in the definition of the wedge product. The exterior algebra is then defined as

$$\Lambda^* V := \bigoplus_{k=0}^{\infty} \Lambda^k V$$

with multiplication given by the wedge product (we set $V^{\otimes 0} = \Lambda^0 V := \mathbb{K}$).

We can also characterize the exterior power and the exterior algebra by means of a universal property.

⁵ $S(k)$ denotes the symmetric group on k elements

Definition 9.8 (Universal property of the exterior power). The $(k\text{-fold})$ exterior power of V , denoted $\Lambda^k V$, is a vector space together with an alternating multilinear map

$$\pi: V^{\times k} \rightarrow \Lambda^k V,$$

such that for any other alternating multilinear map

$$T: V^{\times k} \rightarrow W$$

into a vector space W there exists a unique linear map

$$\tilde{T}: \Lambda^k V \rightarrow W$$

making the following diagram commute

$$\begin{array}{ccc} V^{\times k} & \xrightarrow{T} & W \\ \pi \downarrow & \nearrow \tilde{T} & \\ \Lambda^k V & & \end{array}$$

Definition 9.9 (Universal property of the exterior algebra). The exterior algebra of V , denoted $\Lambda^* V$, is an algebra together with a linear map

$$\iota: V \rightarrow \Lambda^* V,$$

which satisfies $\iota(v)^2 = 0$ for all $v \in V$ and such that for any linear map

$$\psi: V \rightarrow A'$$

into an algebra A' with $\psi(v)^2 = 0$ for all $v \in V$ there exists a unique algebra homomorphism

$$L: \Lambda^* V \rightarrow A'$$

making the following diagram commute

$$\begin{array}{ccc} V & \xrightarrow{\iota} & \Lambda^* V \\ & \searrow \psi & \downarrow L \\ & & A' \end{array}$$

Remark 9.10. The universal property of the exterior power is analogous to the one of the tensor product, where we think of π as the product map $(v_1, \dots, v_k) \mapsto v_1 \wedge \dots \wedge v_k$. So alternating multilinear maps on $V^{\times k}$ are in 1 : 1 correspondence to linear maps on $\Lambda^k V$. In Definition 9.9 we think of ι as an inclusion and of L as extending the map ψ .

Lemma 9.11. *The construction of $\Lambda^k V \subseteq V^{\otimes k}$ at the beginning of the section together with the product map $\pi: V^{\times k} \rightarrow \Lambda^k V$, $(v_1, \dots, v_k) \mapsto v_1 \wedge \dots \wedge v_k$ satisfies the universal property in Definition 9.8.*

Proof. First of all it follows from the graded commutativity of the wedge product, that π is alternating. Uniqueness of $\tilde{T}$ is also clear. To show existence choose a basis $e_1, \dots, e_n \in V$ of V . As mentioned above this yields a basis of $\Lambda^k V$ and we define $\tilde{T}$ on this basis (necessarily) by

$$\tilde{T}(e_{i_1} \wedge \dots \wedge e_{i_k}) := T(e_{i_1}, \dots, e_{i_k})$$

for $1 \leq i_1 < \dots < i_k \leq n$. To see that

$$\begin{array}{ccc} V^{\times k} & \xrightarrow{T} & W \\ \pi \downarrow & \nearrow \tilde{T} & \\ \Lambda^k V & & \end{array}$$

does in fact commute it suffices to verify commutativity on basis vectors $(e_{i_1}, \dots, e_{i_k})$ of $V^{\times k}$ (where the tuple $(i_1, \dots, i_k)$ is not necessarily ordered!). But since T and π are both alternating we may assume after a suitable permutation of $(i_1, \dots, i_k)$, that this tuple is ordered, such that the statement follows from the definition of $\tilde{T}$. $\square$

Lemma 9.12. *The construction of Λ^*V at the beginning of the section together with the canonical inclusion map $V \hookrightarrow \Lambda^*V$ satisfies the universal property in Definition 9.9.*

Proof. Again uniqueness of L is clear. For the existence of L we define it on $\Lambda^k V$ (necessarily) by

$$L(v_1 \wedge \dots \wedge v_k) := \prod_{i=1}^k \psi'(v_i)$$

using the universal property of the exterior power shown in Lemma 9.11. $\square$

To conclude this section let us mention another commonly used construction of the exterior power and exterior algebra. To this end let

$$S^k(V) := \text{span}\{v_1 \otimes \dots \otimes v_k \mid \exists 1 \leq i, j \leq k, i \neq j \text{ with } v_i = v_j\} \subseteq V^{\otimes k}$$

Then it is not hard to show that $V^{\otimes k}/S^k(V)$ together with the canonical map

$$V^{\otimes k} \rightarrow V^{\otimes k} \rightarrow V^{\otimes k}/S^k(V)$$

also satisfies the universal property in Definition 9.8. Hence this is also a valid and commonly used construction of the exterior power. With this definition

$$\alpha \wedge \beta = [\alpha \otimes \beta].$$

Similarly we can also define the exterior algebra as a quotient of the tensor algebra

$$T(V) := \bigoplus_{k=0}^{\infty} V^{\otimes k},$$

by dividing out the two sided ideal $I \subseteq T(V)$ generated by elements of the form $v \otimes v \in T(V)$ for $v \in V$, i.e.

$$\Lambda^*V := T(V)/I.$$

Together with the canonical map $V \rightarrow \Lambda^*V$ this definition is also easily seen to satisfy the universal property in Definition 9.9.

10 Functional Analysis

10.1 L^p and Sobolev Spaces

The partial integration rule for C^1 -functions defined on open subsets of $\mathbb{R}^n$ is shown using Gauß's Theorem, which in turn motivates the definition of weakly differentiable functions for which the partial integration rule holds by definition. Interestingly we can also go the other way around and proof Gauß's Theorem from the partial integration rule, thus establishing the theorem for weakly differentiable functions as well.

Theorem 10.1 (Gauß's Theorem). *Let $U \subseteq \mathbb{R}^n$ be open and $\Omega \subseteq U$ a compact smooth manifold with boundary of codimension 0 with outer unit normal field $\nu: \partial\Omega \rightarrow \mathbb{R}^n$. Given a weakly differentiable continuous vector field $\xi \in L^1_{\text{loc}}(U, \mathbb{R}^n) \cap C^0(U, \mathbb{R}^n)$ with $\partial_i \xi \in L^1_{\text{loc}}(U, \mathbb{R}^n)$ for $1 \leq i \leq n$, the following holds*

$$\int_{\Omega} \operatorname{div}(\xi) \operatorname{vol}_{\Omega} = \int_{\partial\Omega} \langle \xi, \nu \rangle \operatorname{vol}_{\partial\Omega}.$$

Remark 10.2. By the Sobolev embedding Theorems, the conditions on ξ are for example satisfied if $\xi \in W^{k,p}(U, \mathbb{R}^n)$ for $pk > n$.

Proof of Theorem 10.1. The idea of the proof is to construct a particular sequence of cutoff functions

$$\phi_k: U \rightarrow \mathbb{R},$$

which will allow us to approximate both sides of the equation and at the same time lets us relate them by the partial integration rule for ξ . We begin by extending ν arbitrarily to a smooth vector field $\tilde{\nu}$ on all of U . Since $\partial\Omega$ is compact, there exists $\varepsilon > 0$, such that the integral curves of $-\tilde{\nu}$ starting at points in $\partial\Omega$ are defined on $(-\varepsilon, \varepsilon)$. Hence we can define the following bicollar neighbourhood of $\partial\Omega$ using the flow of $-\tilde{\nu}$:

$$\begin{aligned} \Psi: \partial\Omega \times (-\varepsilon, \varepsilon) &\rightarrow U \\ (x, t) &\mapsto \gamma_x(t), \end{aligned}$$

where $\gamma_x(t)$ is the integral curve of $-\tilde{\nu}$ with $\gamma_x(0) = x$. We will now use this bicollar neighbourhood to construct ϕ_k out of cutoff functions $\lambda_k \in C^\infty(\mathbb{R})$, with the following properties

- (i) $\operatorname{supp} \lambda_k \subseteq (-\frac{1}{k}, \frac{1}{k})$
- (ii) $\int_{-\infty}^{\infty} \lambda_k = 1$
- (iii) $\lambda_k(t) = \lambda_k(-t)$

So λ_k is a special kind of Dirac-sequence. The reason for this definition will become clear later. Define

$$\Lambda_k: \partial\Omega \times (-\varepsilon, \varepsilon) \rightarrow \mathbb{R}, (x, t) \mapsto \int_{-\infty}^t \lambda_k(s) ds.$$

Note that for k sufficiently large we have

$$\Lambda_k(x, t) = \begin{cases} 1, & t > \frac{\varepsilon}{3} \\ 0, & t < -\frac{\varepsilon}{3} \end{cases}$$

Moreover choosing ε sufficiently small we may assume, that $\Psi(\partial\Omega \times (0, \varepsilon)) \subseteq \operatorname{int}\Omega$ and $\Psi(\partial\Omega \times (-\varepsilon, 0)) \subseteq U \setminus \Omega$ (here we need compactness of $\partial\Omega$ again). This now allows us to define the desired cutoff functions (for large k) as

$$\phi_k(y) := \begin{cases} \Lambda_k(\Psi^{-1}(y)), & y \in \operatorname{im}\Psi \\ 1, & y \in \operatorname{int}\Omega \setminus \Psi\left(\partial\Omega \times \left[0, \frac{\varepsilon}{2}\right]\right) \\ 0, & y \in (U \setminus \Omega) \setminus \Psi\left(\partial\Omega \times \left[-\frac{\varepsilon}{2}, 0\right]\right) \end{cases}$$

By construction $\phi_k \in C_c^\infty(U)$, since Ω is compact and as $k \rightarrow \infty$

$$\phi_k(y) \rightarrow \chi_\Omega(y) := \begin{cases} 1, & y \in \Omega \\ 0, & y \notin \Omega \end{cases}$$

for almost all $y \in U$ (for $y \in \partial\Omega = \Psi(\partial\Omega \times \{0\})$ we have $\phi_k(y) = \frac{1}{2}$ for all k). By definition of weak differentiability we now get⁶

$$\int_U \operatorname{div}(\xi) \phi_k = - \sum_{i=1}^n \int_U \xi_i \partial_i \phi_k = - \int_U \langle \xi, \nabla \phi_k \rangle. \quad (11)$$

Since $\partial_i \xi_i \in L_{\text{loc}}^1(U)$ we can use the dominated convergence theorem to conclude that the left hand side converges to

$$\int_\Omega \operatorname{div}(\xi)$$

as $k \rightarrow \infty$. To show convergence of the right hand side to

$$\int_{\partial\Omega} \langle \xi, \nu \rangle$$

we will have to analyze this term more closely, which is where our very deliberate construction of ϕ_k comes into play. We begin by computing the gradient $\nabla \phi_k$. Let

$$\beta_k : U \rightarrow \mathbb{R}, y \mapsto \begin{cases} \lambda_k(t), & y = \Psi(x, t) \\ 0, & \text{otherwise} \end{cases}$$

Again for k sufficiently large this is a well defined smooth function. Moreover let

$$\pi : \operatorname{im}(\Psi) \rightarrow \mathbb{R}, \Psi(x, t) \mapsto t.$$

We now claim that

$$\nabla \phi_k = \beta_k \nabla \pi$$

where the right hand side is to be understood to vanish outside of $\operatorname{im}(\Psi)$, so clearly equality holds outside of $\operatorname{im}(\Psi)$. Note that Λ_k is the composition of the projection $\partial\Omega \times (-\varepsilon, \varepsilon) \rightarrow (-\varepsilon, \varepsilon)$ with the map

$$t \mapsto \int_{-\infty}^t \lambda_k(s) ds,$$

so for $y = \Psi(x, t)$ we get by the chain rule

$$d_y \phi_k = d_{(x,t)} \Lambda_k \circ d_y \Psi^{-1} = \lambda_k(t) d_y \pi.$$

Since $\lambda_k(t) = \beta_k(y)$ the assertion follows. Furthermore for $y \in \partial\Omega$ and $\tau \in T_y \partial\Omega$ we have

$$d_y \pi(\tau) = 0$$

and

$$d_y \pi(-\nu(y)) = d_y \pi(\dot{\gamma}_y(0)) = 1,$$

such that $\nabla \pi = -\nu$ on $\partial\Omega$. We can now compute the right hand side of (11) as

$$\int_U \langle \xi, \nabla \phi_k \rangle \operatorname{vol}_U = \int_{\operatorname{im} \Psi} \langle \xi, \nabla \pi \rangle \beta_k \operatorname{vol}_{\operatorname{im} \Psi} = \int_{\partial\Omega \times (-\varepsilon, \varepsilon)} (\langle \xi, \nabla \pi \rangle \circ \Psi) (\beta_k \circ \Psi) \Psi^* \operatorname{vol}_{\operatorname{im} \Psi} \quad (12)$$

Write

$$\Psi^* \operatorname{vol}_{\operatorname{im} \Psi} = \mu \operatorname{vol}_{\partial\Omega \times (-\varepsilon, \varepsilon)}$$

for a smooth function $\mu \in C^\infty(\partial\Omega \times (-\varepsilon, \varepsilon))$. By construction

$$d_{(y,0)} \Psi \left(\frac{\partial}{\partial t} \right) = -\nu(y)$$

and $d_{(y,0)} \Psi(\tau) = \tau$ for all $\tau \in T_y \partial\Omega$. It follows that $\mu(y, 0) = 1$ (if we choose the orientation on $\partial\Omega$ correctly).

⁶For better readability we will sometimes not write the volume forms $\operatorname{vol}_\Omega$ and $\operatorname{vol}_{\partial\Omega}$ in the integrals

Write

$$f := (\langle \xi, \nabla \pi \rangle \circ \Psi) \mu: \partial\Omega \times (-\varepsilon, \varepsilon) \rightarrow \mathbb{R}.$$

Then (12) becomes

$$\int_{\partial\Omega \times (-\varepsilon, \varepsilon)} f(\beta_k \circ \Psi) \text{vol}_{\partial\Omega \times (-\varepsilon, \varepsilon)} = \int_{-\varepsilon}^{\varepsilon} \int_{\partial\Omega} f(-, t) \lambda_k(t) \text{vol}_{\partial\Omega} dt \quad (13)$$

Let

$$\rho(t) := \int_{\partial\Omega} f(-, t) \text{vol}_{\partial\Omega}.$$

Since $\partial\Omega$ is compact and f is continuous (here our assumption $\xi \in C^0(U, \mathbb{R}^n)$ finally comes into play) it follows that ρ is continuous as well. Using this we can simplify (13):

$$\int_{-\varepsilon}^{\varepsilon} \lambda_k(t) \rho(t) dt = \int_{-\varepsilon}^{\varepsilon} \lambda_k(-t) \rho(t) dt \quad (14)$$

For large k this integral only depends on the values of ρ near 0, so using a cutoff function we may assume that ρ is a continuous bounded function defined on all of $\mathbb{R}$. Then finally we can write (14) as

$$\int_{-\infty}^{\infty} \lambda_k(-t) \rho(t) dt = (\rho * \lambda_k)(0),$$

where $\rho * \lambda_k$ denotes the convolution. Since λ_k is a Dirac-sequence it follows from [Sch12] Theorem 4.16. (ii), that

$$(\rho * \lambda_k)(0) \rightarrow \rho(0)$$

as $k \rightarrow \infty$. Hence we have shown, that the right hand side of (11) converges to

$$-\rho(0) = - \int_{\partial\Omega} f(-, 0) \text{vol}_{\partial\Omega}.$$

But $\mu(y, 0) = 1$ and $\nabla \pi(y) = -\nu(y)$ for $y \in \partial\Omega$, so by definition of f

$$f(y, 0) = -\langle \xi(y), \nu(y) \rangle,$$

from which the theorem follows. □

On two equivalent definitions of $L^p(U, \mathbb{R}^n)$

Let $U \subseteq \mathbb{R}^n$ be open. Starting with the space $L^p(U)$ of real valued L^p functions, there are two possible definitions for the space $L^p(U, \mathbb{R}^k)$ of $\mathbb{R}^k$ valued L^p functions. The first definition requires each component function to be L^p :

Definition 10.3. $L^p(U, \mathbb{R}^k) := \{f: U \rightarrow \mathbb{R}^k \text{ measurable} \mid f_i \in L^p(U) \text{ for all } 1 \leq i \leq k\}$

The second definition is more similar to the definition of $L^p(U)$:

Definition 10.4. $L^p(U, \mathbb{R}^k) := \{f: U \rightarrow \mathbb{R}^k \text{ measurable} \mid \|f\| \in L^p(U)\}$

These definitions are however equivalent (as one would certainly hope). To see this we compute for a measurable function $f: U \rightarrow \mathbb{R}^k$:

$$\|f_i\|_p^p = \int_U |f_i|^p d\lambda \leq \int_U \sqrt{|f_1|^2 + \dots + |f_k|^2}^p d\lambda = \int_U \|f\|^p d\lambda$$

and

$$\int_U \|f\|^p d\lambda \leq \int_U (|f_1| + \dots + |f_k|)^p d\lambda \leq \frac{1}{k} \sum_{i=1}^k \int_U |f_i|^p d\lambda,$$

where we used the fact that

$$a^2 + b^2 \leq (a + b)^2$$

for $a, b \geq 0$ in the first inequality and made use of Jensen's inequality in the second inequality. We may also define two norms on $L^p(U, \mathbb{R}^k)$ corresponding to the two definitions:

1. $\|f\|_{p,1} := \sum_{i=1}^k \|f_i\|_p$
2. $\|f\|_{p,2} := \left(\int_U \|f\|^p d\lambda \right)^{\frac{1}{p}}$

Again these norms turn out to be equivalent, which can be seen by a similar computation as above:

$$\|f\|_{p,1} \leq \sum_{i=1}^k \left(\int_U \|f\|^p d\lambda \right)^{\frac{1}{p}} = k \|f\|_{p,2}$$

and conversely

$$\begin{aligned} \|f\|_{p,2} &\leq \left(\left(\frac{1}{k} \right) \sum_{i=1}^k \int_U |f_i|^p d\lambda \right)^{\frac{1}{p}} = \left(\frac{1}{k} \right)^{\frac{1}{p}} \left(\sum_{i=1}^k \|f_i\|_p^p \right)^{\frac{1}{p}} \\ &\leq \left(\frac{1}{k} \right)^{\frac{1}{p}} \left(\sum_{i=1}^k \max_{1 \leq i \leq k} \|f_i\|_p^p \right)^{\frac{1}{p}} = \max_{1 \leq i \leq k} \|f_i\|_p \leq \sum_{i=1}^k \|f_i\|_p = \|f\|_{p,1}. \end{aligned}$$

Note that for $p = 2$ the second norm is the one induced by the L^2 -scalar product on $L^2(U, \mathbb{R}^k)$.

10.2 Complemented Subspaces

Given a vector space X and a subspace $X_0 \subseteq X$ we may want to decompose X as the direct sum of X_0 and some other subspace $X_1 \subseteq X$. More precisely this means that the canonical map

$$X_0 \oplus X_1 \rightarrow X$$

induced by the inclusions $X_i \hookrightarrow X$ should be an isomorphism. In the category of vector spaces this is of course always possible and we call X_1 and **algebraic complement** of X_0 . However the situation looks quite a bit different in the category of topological vector spaces, where such X_1 may not always exist. If it exists we call X_1 a **topological complement** of X_0 and the space X_0 **complemented**. The following Lemma gives a characterization for when an algebraic complement is also a topological complement (at least in the case of Banach spaces).

Lemma 10.5. *Let X be a Banach space and $X_0, X_1 \subseteq X$ subspaces, where X_1 is an algebraic complement of X_0 . Then X_1 is a topological complement of X_0 if and only if X_0 and X_1 are closed.*

Proof. Clearly $X_0, X_1 \subseteq X_0 \oplus X_1$ are closed, so if X is topologically isomorphic to $X_0 \oplus X_1$ they are also closed in X . On the contrary if X_0 and X_1 are closed, they are again Banach spaces and since we assume the canonical map

$$X_0 \oplus X_1 \rightarrow X$$

to be a bijection it follows from the open mapping theorem that this map is already a topological isomorphism. $\square$

Here are two simple sufficient conditions for the existence of topological complements. For a vector space X and a subspace $X_0 \subseteq X$ we define its **codimension** by

$$\text{codim}(X_0) := \dim(X/X_0).$$

Lemma 10.6. *Let X be a Banach space and $X_0 \subseteq X$ a subspace. The following two conditions are both sufficient for the existence of a topological complement of X_0 :*

- (i) $\dim(X_0) < \infty$
- (ii) $\text{codim}(X_0) < \infty$ and X_0 is closed

Proof. First assume (i). Choose a basis $e_1, \dots, e_n$ of X_0 and define linear maps $\phi_i: X_0 \rightarrow \mathbb{R}$ by

$$\phi_i(e_j) = \delta_{ij}.$$

Since X_0 is finite dimensional the ϕ_i are continuous. So using the Hahn-Banach theorem we can extend them to continuous linear functionals Φ_i defined on all of X . Now set

$$X_1 := \{x \in X \mid \Phi_1(x) = \dots = \Phi_n(x) = 0\}.$$

Clearly $X_1 \subseteq X$ is closed, so by Lemma 10.5 we only have to show that X_1 is an algebraic complement of X_0 . To this end let $x \in X$ be arbitrary and define

$$x_0 := \sum_{i=1}^n \Phi_i(x) e_i \in X_0.$$

Then $x_1 := x - x_0 \in X_1$ and $x = x_0 + x_1$. Finally observe that $X_0 \cap X_1 = 0$ from which the statement follows.

For (ii) we begin by choosing any algebraic complement X_1 of X_0 . Since the restriction of the projection

$$X \rightarrow X/X_0$$

to X_1 is an isomorphism, our assumption tells us that X_1 is finite dimensional and so in particular closed. Thus the statement follows again from Lemma 10.5. $\square$

10.3 The Arzelà-Ascoli Theorem

In this section we give a quite general formulation of the Arzelà-Ascoli theorem. The proof we give here is a very straightforward generalization of the one found in [Wer18]. Before stating the theorem let us recall two definitions. Given metric spaces (X, d_X) and (Y, d_Y) and a sequence $(f_n)_{n \in \mathbb{N}} \subseteq C(X, Y)$ of continuous functions we say that $(f_n)_{n \in \mathbb{N}}$ is **pointwise relatively compact** if for all $x \in X$ the set

$$\{f_n(x) \mid n \in \mathbb{N}\} \subseteq Y$$

is relatively compact, i.e. has compact closure. Moreover we call $(f_n)_{n \in \mathbb{N}}$ **uniformly equicontinuous** if for all $\varepsilon > 0$ there exists $\delta > 0$, such that

$$f_n(B_\delta(x)) \subseteq B_\varepsilon(f_n(x))$$

for all $x \in X$ and $n \in \mathbb{N}$.

Theorem 10.7 (Arzelà-Ascoli). *Let (X, d_X) be a compact metric space, (Y, d_Y) a complete metric space and $(f_n)_{n \in \mathbb{N}} \subseteq C(X, Y)$ a sequence of continuous functions, which is pointwise relatively compact and uniformly equicontinuous. Then there exists a subsequence that converges uniformly.*

Proof. We begin by showing that any compact metric space is separable. Since X is compact the open cover

$$\{B_{1/n}(x)\}_{x \in X}$$

of X admits a finite subcover for any $n \in \mathbb{N}$. So there exist finitely many points $x_{n,1}, \dots, x_{n,k(n)} \in X$, such that

$$B_{1/n}(x_{n,1}), \dots, B_{1/n}(x_{n,k(n)})$$

covers X . Then the countable set

$$\bigcup_{n \in \mathbb{N}} \{x_{n,1}, \dots, x_{n,k(n)}\}.$$

is easily seen to be dense.

With this fact at hand we can now actually proof the theorem. Let $x_1, x_2, \dots \in X$ be a dense sequence. Using a diagonalization argument we first construct a subsequence of $(f_n)_{n \in \mathbb{N}}$ which converges pointwise at all x_k . By assumption

$$\{f_n(x_1) \mid n \in \mathbb{N}\} \subseteq Y$$

is relatively compact, such that there exists a subsequence $(f_{k_n^1})_{n \in \mathbb{N}}$ of $(f_n)_{n \in \mathbb{N}}$ for which $(f_{k_n^1}(x_1))_{n \in \mathbb{N}}$ converges in Y . By the same argument we can find a subsequence $(f_{k_n^2})_{n \in \mathbb{N}}$ of $(f_{k_n^1})_{n \in \mathbb{N}}$ for which $(f_{k_n^2}(x_2))_{n \in \mathbb{N}}$ converges in Y and so on. Then the sequence

$$g_n := f_{k_n^n}$$

is a subsequence of $(f_n)_{n \in \mathbb{N}}$ for which $(g_n(x_k))_{n \in \mathbb{N}}$ converges in Y for any $k \in \mathbb{N}$.

We now claim that $(g_n)_{n \in \mathbb{N}}$ is already a Cauchy sequence in $C(X, Y)$ with respect to the metric of uniform convergence. Since Y is complete the same holds for $C(X, Y)$ thus proving the theorem. To show the claim let $\varepsilon > 0$ be arbitrary. By uniform equicontinuity we can find $\delta > 0$, such that

$$g_n(B_\delta(x)) \subseteq B_\varepsilon(g_n(x))$$

for all $x \in X$ and $n \in \mathbb{N}$. Using the fact that $x_1, x_2, \dots$ is dense in the compact space X we can find $N \in \mathbb{N}$, such that

$$B_\delta(x_1), \dots, B_\delta(x_N)$$

cover X . Finally choose $L \in \mathbb{N}$, such that

$$d_Y(g_n(x_k), g_m(x_k)) < \varepsilon$$

for all $n, m \geq L$ and all $k = 1, \dots, N$. Then for arbitrary $x \in X$ there exists some $1 \leq k \leq N$ with $x \in B_\delta(x_k)$. So for $n, m \geq L$ we compute

$$d_Y(g_n(x), g_m(x)) \leq d_Y(g_n(x), g_n(x_k)) + d_Y(g_n(x_k), g_m(x_k)) + d_Y(g_m(x_k), g_m(x)) < 3\varepsilon$$

The claim follows. $\square$

Theorem 10.7 is usually stated differently in the case $Y = \mathbb{R}^k$ by replacing the condition of pointwise relative compactness with the stronger condition of **uniform boundedness**, i.e. there exists some constant $C > 0$, such that

$$\max_{x \in X} |f_n(x)| < C$$

for all $n \in \mathbb{N}$. Since this is usually easier to check in practice let us note this version of the theorem down as well.

Theorem 10.8 ($\mathbb{R}^k$ -valued Arzelà-Ascoli). *Let (X, d_X) be a compact metric space and*

$$f_n: X \rightarrow \mathbb{R}^k$$

a sequence of continuous functions, which is uniformly bounded and uniformly equicontinuous. Then there exists a subsequence that converges uniformly.

As an application of Theorem 10.7 we show that the dual of a compact operator is again compact.

Lemma 10.9. *Let $K: X \rightarrow Y$ be a compact operator between normed spaces. Then its dual $K^*: Y^* \rightarrow X^*$ is compact as well.*

Proof. Let $(\psi_n)_{n \in \mathbb{N}} \subseteq Y^*$ be a sequence bounded in the operator norm by 1. We have to show that

$$\phi_n := K^* \psi_n$$

admits a convergent subsequence. Since

$$C := \overline{K(B_1(0))}$$

is compact (by definition of compact operators) we can try and apply the Arzelà-Ascoli theorem to the sequence

$$f_n := \psi_n|_C.$$

For uniform boundedness note that since C is compact it must be contained in some large ball of radius R , such that for $y \in C$

$$|f_n(y)| = R \left| \psi_n\left(\frac{y}{R}\right) \right| \leq R \|\psi_n\|_{\text{op}} \leq R.$$

To see uniform equicontinuity note that for $y, z \in C$

$$|f_n(y) - f_n(z)| = |\psi(y - z)| \leq \|y - z\|.$$

So by the Arzelà-Ascoli theorem we may assume that $(f_n)_{n \in \mathbb{N}}$ converges uniformly (at least after restriction to a subsequence). We now claim that $(\phi_n)_{n \in \mathbb{N}}$ is Cauchy with respect to the operator norm. For $x \in B_1(0)$ we compute

$$|\phi_n(x) - \phi_m(x)| = |f_n(Kx) - f_m(Kx)| \leq \max |f_n - f_m|.$$

Since $(f_n)_{n \in \mathbb{N}}$ is Cauchy the statement follows. $\square$

11 Whitney Topologies

11.1 Approximation Results

Let M, N be smooth manifolds with or without boundary. Consider the set $C^\infty(M, N)$ of smooth maps. On this set one can define the so called Whitney topologies, which provide a precise language to make sense of such statements as "slightly perturbing f yields a map transversal to a given submanifold". In the following we will prove a number of useful approximation results of this kind concerning immersions, one-to-one immersions, embeddings and transversal maps.

Let us begin by recalling the definition of the Whitney topologies. First we define a topology on the space $C^k(M, N)$ of C^k maps from M to N . We will be using the definition from [Hir12] using local charts. For the discussion of an equivalent definition see section 11.2. Suppose we are given the following data

- A family of charts $\Phi = (U_\alpha, \phi_\alpha)_{\alpha \in J}$ for M , such that $(U_\alpha)_{\alpha \in J}$ is locally finite, that is each point $x \in M$ possesses a neighbourhood that intersects only finitely many U_α
- A family of charts $\Psi = (V_\alpha, \psi_\alpha)_{\alpha \in J}$ for N
- A family $K = (K_\alpha)_{\alpha \in J}$ of compact sets $K_\alpha \subseteq U_\alpha$
- A family of positive numbers $\varepsilon = (\varepsilon_\alpha)_{\alpha \in J}$

Then if $f \in C^k(M, N)$ with $f(K_\alpha) \subseteq V_\alpha$ for each $\alpha \in J$, we define a basic open neighbourhood

$$\mathcal{N}^k(f, \Phi, \Psi, K, \varepsilon)$$

as the set of functions $g \in C^k(M, N)$, such that for all $\alpha \in J$, $g(K_\alpha) \subseteq V_\alpha$ and the coordinate representations $f_\alpha := \psi_\alpha^{-1} \circ f \circ \phi_\alpha$ and $g_\alpha := \psi_\alpha^{-1} \circ g \circ \phi_\alpha$ defined on some open neighbourhood of $\phi_\alpha^{-1}(K_\alpha)$ satisfy

$$\|f_\alpha - g_\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} := \sum_{|\beta| \leq k} \max_{x \in \phi_\alpha^{-1}(K_\alpha)} \|\partial^\beta f_\alpha(x) - \partial^\beta g_\alpha(x)\| < \varepsilon_\alpha.$$

The sum is taken over multiindices $\beta \in \mathbb{N}_0^n$ (where $n = \dim(M)$). The collection of all such basic open neighbourhoods form the basis for a topology on $C^k(M, N)$ called the Whitney C^k -topology (it is easy to see that the intersection of two such neighbourhoods is again of the same form). The Whitney C^∞ -topology on $C^\infty(M, N)$ is then simply defined as the topology induced by all inclusion maps $C^\infty(M, N) \rightarrow C^k(M, N)$.

A first interesting approximation result is the following.

Theorem 11.1. *Let M and N be manifolds with boundary. Then $C^k(M, N)$ is dense in $C^0(M, N)$, $0 \leq k \leq \infty$.*

For a proof see [Hir12]. Moreover many interesting classes of smooth maps are open in the Whitney topologies:

Theorem 11.2. *Let M and N be manifolds with boundary. Then the following sets are open*

- (i) *The set of immersions $\text{Imm}^k(M, N) \subseteq C^k(M, N)$, $k \geq 1$,*
- (ii) *The set of submersions $\text{Sub}^k(M, N) \subseteq C^k(M, N)$, $k \geq 1$,*
- (iii) *The set of embeddings $\text{Emb}^k(M, N) \subseteq C^k(M, N)$, $k \geq 1$,*
- (iv) *The set of proper maps $\text{Prop}^k(M, N) \subseteq C^k(M, N)$, $k \geq 0$.*

Again this is shown in [Hir12]. We now state the announced approximation results. The proofs will be given further below. For simplicity we assume all maps to be smooth from this point on ⁷. By Theorem 11.1 this is not a serious restriction.

⁷The reason for this is to avoid discussions about the necessary regularity conditions required by Sard's Theorem for C^k -functions, $k < \infty$.

Theorem 11.3 (Immersion). *Let M^n and N^p be manifolds with boundary, $p \geq 2n$. Then the set of immersions $\text{Imm}(M, N) \subseteq C^\infty(M, N)$ is dense. Moreover if $f: M \rightarrow N$ is already an immersion on some closed subset $A \subseteq M$ there exists an immersion $g: M \rightarrow N$ with $g|_A = f|_A$ arbitrarily close to f .*

Here the term “arbitrarily close” is to be understood with respect to the Whitney C^∞ -topology on $C^\infty(M, N)$. Increasing the dimension of N by one we even obtain the following.

Theorem 11.4 (One-to-one Immersions). *Let M^n and N^p be manifolds with boundary, $p \geq 2n+1$. Then the set of one-to-one immersions is dense in $C^\infty(M, N)$. Moreover if $f: M \rightarrow N$ is already a one-to-one immersion on some neighbourhood $U \subseteq M$ of a closed subset $A \subseteq M$ there exists a one-to-one immersion $g: M \rightarrow N$ with $g|_A = f|_A$ arbitrarily close to f .*

In general of course a one-to-one immersion need not be an embedding if M is not compact and in fact the set of embeddings is not dense in $C^\infty(M, N)$ no matter the dimension of N . See [Hir12] on page 55 for a counterexample. However one-to-one immersions which are also proper are embeddings. Since the set of proper maps is open by Theorem 11.2 we obtain:

Corollary 11.5 (Embeddings). *Let M^n and N^p be manifolds with boundary, $p \geq 2n+1$. Then the set of proper embeddings is dense in $\text{Prop}(M, N)$. Moreover if $f: M \rightarrow N$ is a proper map which is a one-to-one immersion on some neighbourhood $U \subseteq M$ of a closed subset $A \subseteq M$ there exists a proper embedding $g: M \rightarrow N$ with $g|_A = f|_A$ arbitrarily close to f .*

In Theorem 11.4 and Corollary 11.5 the existence of such a neighbourhood $U \subseteq M$ is automatically guaranteed if A is compact by virtue of the following Lemma.

Lemma 11.6. *Let M and N be manifolds with boundary and $f: M \rightarrow N$ a smooth map, which is a one-to-one immersion on a compact set $A \subseteq M$. Then f is already a one-to-one immersion on some neighbourhood $U \subseteq M$ of A .*

Proof. Suppose this were not the case. Then we can use compactness A to construct sequences $(a_n)_{n \in \mathbb{N}}$ and $(a'_n)_{n \in \mathbb{N}}$ in M with $a_n \neq a'_n$ and $f(a_n) = f(a'_n)$, which converge to points z and z' in A respectively. This implies $f(z) = f(z')$ and since f is one-to-one on A we conclude $z = z'$. However, since f is an immersion at z it must be one-to-one on a neighbourhood of z contradicting $a_n \neq a'_n$. $\square$

Finally we have a statement to approximate any smooth map $f: M \rightarrow N$ by one which is transversal to a given submanifold. We denote the restriction of f to the boundary ∂M of M by $\partial f: \partial M \rightarrow N$.

Theorem 11.7 (Transversality). *Let M and N be manifolds with boundary and $S \subseteq N$ a submanifold with boundary. Then the set of maps $g: M \rightarrow N$ with $g \pitchfork S$ and $\partial g \pitchfork S$ is dense in $C^\infty(M, N)$. Moreover if $f: M \rightarrow N$ already satisfies $f \pitchfork S$ for all points in some closed subset $A \subseteq M$ and $\partial f \pitchfork S$ for all points in $A \cap \partial M$, there exists a map $g: M \rightarrow N$ with $g \pitchfork S$, $\partial g \pitchfork S$ and $g|_A = f|_A$ arbitrarily close to f .*

Remark 11.8. The proof of Theorem 11.7 also shows the following version of this Theorem, which may be useful. If $f: M \rightarrow N$ satisfies $f \pitchfork S$ at all points in A but not necessarily $\partial f \pitchfork S$ at all points in $A \cap \partial M$, there still exists $g: M \rightarrow N$ arbitrarily close to f with $g \pitchfork S$ and $g|_A = f|_A$ but not necessarily with $\partial g \pitchfork S$.

In practice one is often interested in the following problem: Given a map $f: M \rightarrow N$ can we find a map $g: M \rightarrow N$ homotopic to f with certain desired properties? All of the above theorems can be applied in this context by virtue of the following Proposition, which furthermore allows us to impose conditions on the homotopy itself.

Proposition 11.9. *Let M and N be manifolds with boundary, $\mathcal{U} \subseteq C^\infty(M, N)$ be some open subset and $f \in \mathcal{U}$. Then for any $g \in \mathcal{U}$ sufficiently close to f there exists a homotopy $h_t: M \rightarrow N$, $0 \leq t \leq 1$, from f to g with $h_t \in \mathcal{U}$ for all $0 \leq t \leq 1$. Moreover if $f|_A = g|_A$ for some closed subset $A \subseteq M$ we can choose this homotopy rel A .*

Together all of these results comprise a very efficient machinery for approaching problems in differential topology. As an example we proof the following, which - up to the precise restrictions on the dimension - is a generalisation of Theorem 8.8.

Corollary 11.10. *Let M^n and N^p be compact manifolds with boundary of dimensions $p \geq 2n + 3$. Then two embeddings $f, g: M \rightarrow N$ are isotopic if and only if they are homotopic.*

Proof. By Theorem 11.7 we can find $f': M \rightarrow N$ arbitrarily close to f , such that f' is transversal to $g(M)$, which for these relative dimensions is equivalent to $f'(M) \cap g(M) = \emptyset$. By Theorem 11.2 we may assume f' to still be an embedding and by Proposition 11.9 we may assume that f' is homotopic to f via some homotopy h_t which is an embedding for each time $t \in [0, 1]$. In other words f' and f are isotopic. So without loss of generality we can assume that $f' = f$. Let $H: M \times \mathbb{R} \rightarrow N$ be some smooth homotopy with $H(-, 0) = f$ and $H(-, 1) = g$, which exists by assumption. Then H is a one-to-one immersions on the compact subset $A := M \times 0 \cup M \times 1 \subseteq M \times \mathbb{R}$ since we already established that $f(M) \cap g(M) = \emptyset$. Thus H is also a one-to-one immersion on some neighbourhood of A by Lemma 11.6. Since $2 \dim(M \times [0, 1]) + 1 = 2n + 3 \leq p$ we can apply Theorem 11.4 to replace H by some other homotopy $H': M \times \mathbb{R} \rightarrow N$, which is a one-to-one immersion and agrees with H on A . In particular $H'(-, t): M \rightarrow N$ is an embedding for each $t \in [0, 1]$ since M is compact. Thus H' is an isotopy with $H'(-, 0) = f$ and $H'(-, 1) = g$. $\square$

Proofs of the Approximation Theorems

In this section we will proof Theorem 11.3, 11.4, 11.7 as well as Proposition 11.9. The proofs of Theorem 11.3 and 11.4 are generalisations of the proofs of Theorem 2.5 and Lemma 2.5 in [Ada93] respectively.

In the following we will necessarily need to compute some C^k -norms. The following technical Lemma takes care of all of these computations. We denote by $\mathbb{H}^n$ the n -dimensional half space.

Lemma 11.11. *Suppose we are given open sets $U, U' \subseteq \mathbb{H}^n, W, W' \subseteq \mathbb{H}^m$, compact subsets $K \subseteq U, L \subseteq W$, a smooth function $\lambda: U \rightarrow \mathbb{R}$, diffeomorphisms $\phi: U' \rightarrow U$ and $\psi: W \rightarrow W'$ as well as constants $\varepsilon > 0$ and $C > 0$. Then there exists $\delta > 0$ with the following property. Let $r, s, \theta: U_K \rightarrow W$ be smooth functions defined on some neighbourhood $U_K \subseteq U$ of K , which can depend on the functions r, s, θ . Suppose that s satisfies $s(K) \subseteq L$ and $\|s\|_{C^k(K)} \leq C$. Then if*

$$\|r - s\|_{C^k(K)} < \delta \text{ and } \|\theta\|_{C^k(K)} < \delta$$

the following holds.

- (i) $\|\psi \circ r - \psi \circ s\|_{C^k(K)} < \varepsilon$,
- (ii) $\|\theta \circ \phi\|_{C^k(\phi^{-1}(K))} < \varepsilon$,
- (iii) $\|\lambda \theta\|_{C^k(K)} < \varepsilon$.

Proof. We start with (i). Let $\alpha \in \mathbb{N}_0^n$ be some multiindex of order $|\alpha| = l \leq k$. A computation shows that $\partial^\alpha(\psi \circ r)$ can be expressed as a sum of the form

$$\partial^\alpha(\psi \circ r) = \sum_{|\beta| \leq l} P_\beta^\alpha((\partial^\gamma r_j)_{|\gamma| \leq l, j=1, \dots, m})(\partial^\beta \psi) \circ r$$

for polynomials $P_\beta^\alpha((\partial^\gamma r_j)_{|\gamma| \leq l, j=1, \dots, m})$ in the partial derivatives $\partial^\gamma r_j$ of the component functions r_j of r . These Polynomials

$$P_\beta^\alpha: \mathbb{R}^{N(l, m)} \rightarrow \mathbb{R}$$

only depend on the multiindices α, β and m , where $N(l, m) \in \mathbb{N}$ denotes the cardinality of the set $\{(\gamma, j) \in \mathbb{N}_0^n \times \{1, \dots, m\} \mid |\gamma| \leq l\}$. Similiarly

$$\partial^\alpha(\psi \circ s) = \sum_{|\beta| \leq l} P_\beta^\alpha((\partial^\gamma s_j)_{|\gamma| \leq l, j=1, \dots, m})(\partial^\beta \psi) \circ s$$

for the same polynomials P_β^α . For simplicity write

$$\begin{aligned} R_\beta^\alpha &:= P_\beta^\alpha((\partial^\gamma r_j)_{|\gamma| \leq l, j=1, \dots, m}), \\ S_\beta^\alpha &:= P_\beta^\alpha((\partial^\gamma s_j)_{|\gamma| \leq l, j=1, \dots, m}). \end{aligned}$$

So in order for $\|\partial^\alpha(\psi \circ r) - \partial^\alpha(\psi \circ s)\|_{C^0(K)}$ to be small we have to choose δ , such that

$$\|R_\beta^\alpha(\partial^\beta\psi) \circ r - S_\beta^\alpha(\partial^\beta\psi) \circ s\|_{C^0(K)}$$

is small for each β . Using the triangle inequality we compute

$$\begin{aligned} & \|R_\beta^\alpha(\partial^\beta\psi) \circ r - S_\beta^\alpha(\partial^\beta\psi) \circ s\|_{C^0(K)} \\ & \leq \|R_\beta^\alpha(\partial^\beta\psi) \circ r - R_\beta^\alpha(\partial^\beta\psi) \circ s\|_{C^0(K)} + \|(S_\beta^\alpha - R_\beta^\alpha)(\partial^\beta\psi) \circ s\|_{C^0(K)} \\ & \leq \|R_\beta^\alpha\|_{C^0(K)}\|(\partial^\beta\psi) \circ r - (\partial^\beta\psi) \circ s\|_{C^0(K)} + \|S_\beta^\alpha - R_\beta^\alpha\|_{C^0(K)}\|(\partial^\beta\psi) \circ s\|_{C^0(K)} \end{aligned} \quad (15)$$

Choosing $\delta < 1$ we obtain from $\|s\|_{C^k(K)} \leq C$ that also $\|r\|_{C^k(K)} \leq C + 1$. So in particular $|\partial^\gamma r_j| \leq C + 1$ on K and so if we write $I := [-(C + 1), C + 1]$ we conclude that

$$\|R_\beta^\alpha\|_{C^0(K)} \leq \|P_\beta^\alpha\|_{C^0(I^{N(l,m)})}$$

Note that the right hand side is some constant not depending on r and s . We now claim that if we choose $\delta > 0$ sufficiently small then

$$\|(\partial^\beta\psi) \circ r - (\partial^\beta\psi) \circ s\|_{C^0(K)}$$

can be made arbitrarily small. To see this we fix some compact neighbourhood $\tilde{L} \subseteq W$ of L . Then there exists some small radius $\rho > 0$ depending only on L and $\tilde{L}$, such that each ball of radius ρ around a point in L is contained in $\tilde{L}$. Thus if we choose $\delta < \rho$ we can conclude from $s(K) \subseteq L$ that $r(K) \subseteq \tilde{L}$. But by compactness $\partial^\beta\psi$ is uniformly continuous on $\tilde{L}$ from which the claim follows. Thus we can guarantee that the first term in (15) is arbitrarily small for sufficiently small $\delta > 0$, which does not depend on r, s . For the second term we argue as follows. First of all we obtain from $s(K) \subseteq L$ that

$$\|(\partial^\beta\psi) \circ s\|_{C^0(K)} \leq \|\partial^\beta\psi\|_{C^0(L)}.$$

Again the right hand side is some constant not depending on r, s . Thus we have to show that

$$\|S_\beta^\alpha - R_\beta^\alpha\|_{C^0(K)}$$

can be made arbitrarily small for sufficiently small δ . To see this note that $(\partial^\gamma r_j)_{|\gamma| \leq l, j=1, \dots, m}$ restricted to K only takes values in the compact set I^n defined above. The same holds for $(\partial^\gamma s_j)_{|\gamma| \leq l, j=1, \dots, m}$. Hence the statement follows from the fact that P_β^α is uniformly continuous on the compact set I^n . This shows that δ can be chosen such that (i) holds.

For (ii) note that we can again write

$$\partial^\alpha(\theta \circ \phi) = \sum_{|\beta| \leq l} T_\beta^\alpha((\partial^\gamma \phi_j)_{|\gamma| \leq l, j=1, \dots, n})(\partial^\beta \theta) \circ \phi$$

for any multiindex $\alpha \in \mathbb{N}_0^n$ of order $|\alpha| = l \leq k$, this time for potentially different polynomials T_β^α . Again we abbreviate $\Phi_\beta^\alpha := T_\beta^\alpha((\partial^\gamma \phi_j)_{|\gamma| \leq l, j=1, \dots, n})$. We now have to choose $\delta > 0$, such that

$$\|\Phi_\beta^\alpha(\partial^\beta \theta) \circ \phi\|_{C^0(\phi^{-1}(K))} \leq \|\Phi_\beta^\alpha\|_{C^0(\phi^{-1}(K))}\|(\partial^\beta \theta) \circ \phi\|_{C^0(\phi^{-1}(K))}$$

is small for each β . But $\|(\partial^\beta \theta) \circ \phi\|_{C^0(\phi^{-1}(K))} = \|\partial^\beta \theta\|_{C^0(K)} < \delta$ and $\|\Phi_\beta^\alpha\|_{C^0(\phi^{-1}(K))}$ is just some constant not depending on θ .

Finally for (iii) it is easy to compute that

$$\partial^\alpha(\lambda \theta) = \sum_{|\beta| + |\gamma| = l} \partial^\beta \lambda \partial^\gamma \theta$$

for any multiindex $\alpha \in \mathbb{N}_0^n$ of order $|\alpha| = l \leq k$. Since

$$\|\partial^\beta \lambda \partial^\gamma \theta\|_{C^0(K)} \leq \|\partial^\beta \lambda\|_{C^0(K)} \|\partial^\gamma \theta\|_{C^0(K)} < \delta \|\partial^\beta \lambda\|_{C^0(K)}$$

and $\|\partial^\beta \lambda\|_{C^0(K)}$ is just some constant not depending on θ the statement follows. $\square$

We now begin with the proof of Theorem 11.3 for which we will need the following Lemma, which is essentially a local version of the theorem we aim to proof.

Lemma 11.12. *Let $f: U \rightarrow \mathbb{R}^p$, $p \geq 2n$, be a smooth function defined on an open subset $U \subseteq \mathbb{H}^n$. Then for almost all $A \in \mathbb{R}^{p \times n}$ the map $g: U \rightarrow \mathbb{R}^p$ defined by*

$$g(x) := f(x) + Ax$$

is an immersion.

Proof. First we argue that we may assume $\partial U = \emptyset$. If this is not the case we can use a partition of unity to extend f to a smooth function $f': U' \rightarrow \mathbb{R}^p$ defined on an open subset $U' \subseteq \mathbb{R}^n$ containing U . Then apply the Lemma to f' and restrict to U again. Thus we can assume $\partial U = \emptyset$. Define

$$F: U \times S^{n-1} \rightarrow U \times S^{n-1} \times \mathbb{R}^p, (x, v) \mapsto (x, v, D_x f(v)).$$

Clearly F is an embedding and thus its image $S := \text{im}(F)$ is a $2n - 1$ dimensional submanifold of $U \times S^{n-1} \times \mathbb{R}^p$. We aim to find $A \in \mathbb{R}^{p \times n}$, such that $D_x f + A$ is an injective map $\mathbb{R}^n \rightarrow \mathbb{R}^p$ for all $x \in U$. Phrased differently, $D_x f(v) \neq Av$ should hold for all $v \in S^{n-1}$ and $x \in U$. This in turn is equivalent to the requirement that the map

$$\lambda_A: U \times S^{n-1} \rightarrow U \times S^{n-1} \times \mathbb{R}^p, (x, v) \mapsto (x, v, -Av)$$

is transversal to S , since $(2n - 1) + (2n - 1) < 2n - 1 + p$. To find such A consider the map

$$\lambda: U \times S^{n-1} \times \mathbb{R}^{p \times n} \rightarrow U \times S^{n-1} \times \mathbb{R}^p, (x, v, A) \mapsto (x, v, -Av).$$

A simple computation shows that λ is a submersion, so the statement follows from the parametric transversality theorem. $\square$

With this Lemma at hand we can now proof Theorem 11.3.

Proof of Theorem 11.3. By definition of the topology on $C^\infty(M, N)$ we have to show the following. For any neighbourhood of the form $\mathcal{N}^k = \mathcal{N}^k(f, \Phi, \Psi, K, \varepsilon) \subseteq C^k(M, N)$, there exists an immersion $g \in \mathcal{N} := \mathcal{N}^k \cap C^\infty(M, N)$ which agrees with f on the subset A . For the data $\Phi, \Psi, K, \varepsilon$ used to define the neighbourhood $\mathcal{N}^k$ we will be using the same notation as at the beginning of this section.

Let $U \subseteq M$ be an open neighbourhood of A on which f is an immersion. Choose a locally finite cover of charts $(O_i, o_i)_{i \in \mathbb{Z}}$ of M , compact subsets $B_i, D_i \subseteq O_i$ which still cover M , charts $(W_i, w_i)_{i \in \mathbb{Z}}$ of N as well as cut-off functions $\lambda_i: M \rightarrow [0, 1]$, such that

- (i) $\lambda_i = 1$ in a neighbourhood of B_i ,
- (ii) D_i contains the support of λ_i in its interior,
- (iii) $f(D_i) \subseteq W_i$,
- (iv) $D_i \subseteq U$ for $i \leq 0$ and $D_i \subseteq M - A$ for $i > 0$.

The existence of the data described above follows easily from paracompactness of M by using the following basic fact: For two given open covers of M one can always find a third open cover which is a locally finite refinement of both.

We will construct g as the limit of an inductively defined sequence $f_n: M \rightarrow N$, $n \geq 0$, with $f_0 := f$, satisfying the following properties.

- 1) f_n is an immersion on $C_n := \bigcup_{-\infty \leq i \leq n} B_i$,
- 2) $f_n = f_{n-1}$ outside D_n ,
- 3) $f_n(D_i) \subseteq W_i$ for all $i \in \mathbb{Z}$,
- 4) $f_n(K_\alpha) \subseteq V_\alpha$ for all $\alpha \in J$ and the coordinate representations $f_n^\alpha := \psi_\alpha^{-1} \circ f_n \circ \phi_\alpha$ of f_n satisfy

$$\|f_n^\alpha - f_{n-1}^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} < \frac{\varepsilon_\alpha}{2^n}.$$

It is clear that $f_0 = f$ satisfies these properties. Let us first assume $\partial N = \emptyset$. We will construct f_n by applying Lemma 11.12 to f_{n-1} expressed in the coordinates of the charts w_n and o_n for suitable matrices $A_n \in \mathbb{R}^{p \times n}$. Define

$$f_n(x) := \begin{cases} w_n(w_n^{-1}(f_{n-1}(x)) + \lambda_n(x)A_n y), & x = o_n(y) \in D_n \\ f_{n-1}(x), & x \notin D_n \end{cases}$$

For sufficiently small A_n this is well defined, that is $w_n^{-1}(f_{n-1}(x)) + \lambda_n(x)A_n y$ is contained in the domain of w_n . This is due to compactness of $f_{n-1}(D_n) \subseteq W_n$ and the fact that W_n is an open subset of $\mathbb{R}^p$, as we assumed $\partial N = \emptyset$. Smoothness of f_n follows from (ii). By definition $f_n(D_n) \subseteq W_n$ and similarly to well definedness we can choose A_n , such that $f_n(D_i) \subseteq W_i$ at least for finitely many $i \in \mathbb{Z}$. This however suffices to achieve the same for all $i \in \mathbb{Z}$ simultaneously. Indeed, note that by local finiteness of $(O_i)_{i \in \mathbb{Z}}$ and compactness of D_n only finitely many O_i , and so in particular only finitely many D_i , intersect D_n . But $f_{n-1} = f_n$ outside D_n and $f_{n-1}(D_i) \subseteq W_i$ already holds for all $i \in \mathbb{Z}$. Thus 3) holds for sufficiently small A_n . An analogous argument allows us to ensure that $f_n(K_\alpha) \subseteq V_\alpha$ for all $\alpha \in J$ by using local finiteness of $(U_\alpha)_{\alpha \in J}$ instead. Furthermore 2) holds by definition.

Moreover we can apply Lemma 11.12 to choose A_n , such that f_n is an immersion on an open neighbourhood of B_n on which $\lambda_n = 1$, which exists by (i). In particular it is an immersion on B_n . We can furthermore guarantee that f_n remains an immersion on C_{n-1} by choosing A_n sufficiently small. To see this note that C_{n-1} is closed by local finiteness of $(O_i)_{i \in \mathbb{Z}}$ and thus $D_n \cap C_{n-1}$ is compact. The claim then follows essentially from the fact that the set of $p \times n$ matrices of full rank is open in $\mathbb{R}^{p \times n}$. Thus 1) is also established.

It only remains to show the second part of 4). For $x \in \phi_\alpha^{-1}(K_\alpha \cap D_n)$ let

$$\begin{aligned} \hat{f}_{n-1}^\alpha(x) &:= w_n^{-1}(f_{n-1}(\phi_\alpha(x))), \\ \hat{A}_n^\alpha(x) &:= \lambda_n(\phi_\alpha(x)) A_n o_n^{-1}(\phi_\alpha(x)), \\ \tau_n^\alpha(x) &:= \psi_\alpha^{-1}(w_n(x)). \end{aligned}$$

In fact all of these expressions may be defined on an open neighbourhood of $\phi_\alpha^{-1}(K_\alpha \cap D_n)$. Since $f_n^\alpha = f_{n-1}^\alpha$ outside $\phi_\alpha^{-1}(K_\alpha \cap D_n)$ we have

$$\|f_n^\alpha - f_{n-1}^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} = \|\tau_n^\alpha \circ (\hat{f}_{n-1}^\alpha + \hat{A}_n^\alpha) - \tau_n^\alpha \circ \hat{f}_{n-1}^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha \cap D_n))}$$

Note that we can make $\hat{A}_n^\alpha$ and all of its partial derivatives arbitrarily small by choosing A_n sufficiently small (this is essentially Lemma 11.11 (iii)). Thus we can apply Lemma 11.11 (i) to

- $r = \hat{f}_{n-1}^\alpha + \hat{A}_n^\alpha$,
- $s = \hat{f}_{n-1}^\alpha$,
- $\psi = \tau_n^\alpha$,
- $K = \phi_\alpha^{-1}(K_\alpha \cap D_n)$
- $L = s(K)$,
- $C = \|s\|_{C^k(K)}$

to conclude that the above sum can be made arbitrarily small for sufficiently small A_n . Again a local finiteness argument shows that $\|f_n^\alpha - f_{n-1}^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))}$ is non-zero for only finitely many $\alpha \in J$. This shows 4).

We now define

$$g := \lim_{n \rightarrow \infty} f_n.$$

This is a well defined smooth map, since local finiteness of $(O_i)_{i \in \mathbb{Z}}$ implies that an arbitrary point $x \in M$ has a neighbourhood $U(x)$ which intersects only finitely many D_i . Thus for sufficiently large $n > 0$ we conclude from 2) that

$$f_n = f_{n+1} = f_{n+2} = \dots$$

on $U(x)$. It then follows from 1) that g is an immersion, since the B_i cover M . Moreover property (iv) of the D_i makes sure that g agrees with f on A . Finally another local finiteness argument shows that for a given $\alpha \in J$ there exists some sufficiently large $n > 0$, such that $g = f_n$ in a neighbourhood of K_α . So by 4) we obtain $g(K_\alpha) \subseteq V_\alpha$ and the coordinate representations $g^\alpha := \psi_\alpha^{-1} \circ g \circ \phi_\alpha$ and $f^\alpha := \psi_\alpha^{-1} \circ f \circ \phi_\alpha$ satisfy

$$\begin{aligned} \|g^\alpha - f^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} &= \|f_n^\alpha - f_0^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} \\ &\leq \sum_{i=0}^{n-1} \|f_{i+1}^\alpha - f_i^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} \\ &< \sum_{i=0}^{n-1} \frac{\varepsilon_\alpha}{2^{i+1}} \\ &< \varepsilon_\alpha. \end{aligned}$$

Hence $g \in \mathcal{N}$ as we wanted to show.

This completes the proof for $\partial N = \emptyset$. Next we explain the necessary changes if $\partial N \neq \emptyset$. In this case the definition of f_n becomes an issue since

$$w_n^{-1}(f_{n-1}(x)) + \lambda_n(x)A_n y$$

may not be contained in the domain of w_n anymore if w_n is a map around the boundary of N . We can however resolve this by slightly modifying the definition. In the following let

$$\mathbb{H}^p := \{(y_1, \dots, y_p) \in \mathbb{R}^p \mid y_1 \leq 0\}$$

be our model for the half space and denote by $e_1 = (1, 0, \dots, 0) \in \mathbb{R}^p$ the first standard basis vector. Now define

$$f_n(x) := \begin{cases} w_n(w_n^{-1}(f_{n-1}(x)) + \lambda_n(x)A_n y - \lambda_n(x)\delta_n e_1), & x = o_n(y) \in D_n \\ f_{n-1}(x), & x \notin D_n \end{cases}$$

for some suitable scalar $\delta_n > 0$, which we define next. We choose δ_n , such that the first component of the vector

$$A_n y - \delta_n e_1$$

is non-positive for all $y \in o_n^{-1}(D_n)$. By compactness of D_n such δ_n always exists and can be chosen arbitrarily small for sufficiently small A_n . With this modification we can choose A_n and δ_n small enough, such that

$$w_n^{-1}(f_{n-1}(x)) + \lambda_n(x)A_n y - \lambda_n(x)\delta_n e_1$$

is again contained in the domain of w_n . Moreover the added term $\lambda_n(x)\delta_n e_1$ does not change the fact that f_n is an immersion on B_n for almost all A_n , since this term is constant a neighbourhood of B_n . The rest of the proof then goes through unchanged if in addition to A_n we also choose δ_n sufficiently small in each step. $\square$

Next we will proof Theorem 11.4 about one-to-one immersions.

Proof of Theorem 11.4. First we may assume, that f is already an immersion. To see this choose some closed neighbourhood $U_A \subseteq U$ of A . By Theorem 11.3, which we just proofed, we can choose an immersion arbitrarily close to f , which agrees with f on the closed set U_A and so in particular this immersion is still one-to-one in an neighbourhood of A . So let us assume, that f is an immersion.

Using the same notation as in the proof of Theorem 11.3 let $\mathcal{N} := \mathcal{N}^k(f, \Phi, \Psi, K, \varepsilon) \cap C^\infty(M, N)$ be some open neighbourhood of f . We choose the following data: A locally finite cover $(D_i)_{i \in \mathbb{Z}}$ of M of compact sets $D_i \subseteq M$, open sets $B_i \subseteq M$ which still cover M , charts $(W_i, w_i)_{i \in \mathbb{Z}}$ of N as well as cut-off functions $\lambda_i: M \rightarrow [0, 1]$, such that

- (i) $\lambda_i^{-1}((0, 1]) = B_i$,
- (ii) D_i contains the support of λ_i , that is $\overline{B_i}$, in its interior,
- (iii) $f(D_i) \subseteq W_i$,

(iv) $D_i \subseteq U$ for $i \leq 0$ and $D_i \subseteq M - A$ for $i > 0$,

(v) $f|_{B_i}$ is injective.

Again the existence of this data follows from paracompactness of M . For (v) recall that immersions are locally embeddings. We now inductively define a sequence $f_n: M \rightarrow N$, $n \geq 0$, with $f_0 := f$, satisfying the following properties.

- 1) f_n is an immersion and is injective on $C_n := \bigcup_{-\infty \leq i \leq n} B_i$,
- 2) $f_n = f_{n-1}$ outside D_n ,
- 3) $f_n(D_i) \subseteq W_i$ for all $i \in \mathbb{Z}$,
- 4) $f_n(K_\alpha) \subseteq V_\alpha$ for all $\alpha \in J$ and the coordinate representations $f_n^\alpha := \psi_\alpha^{-1} \circ f_n \circ \phi_\alpha$ of f_n satisfy

$$\|f_n^\alpha - f_{n-1}^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} < \frac{\varepsilon_\alpha}{2^n}.$$

Here properties 2)- 4) are the same as in the proof of Theorem 11.3. Again it is clear that $f_0 = f$ satisfies these properties. At first we want to assume that both M and N have empty boundary. We define f_n using f_{n-1} by

$$f_n(x) := \begin{cases} w_n(w_n^{-1}(f_{n-1}(x)) + \lambda_n(x)b_n), & x \in D_n, \\ f_{n-1}(x), & x \notin D_n \end{cases}$$

for some suitable vector $b_n \in \mathbb{R}^p$, which we choose next. Firstly it follows just as in the proof of Theorem 11.3 that for sufficiently small b_n , f_n is a well defined smooth map satisfying 2), 3) and 4). Furthermore we can choose b_n small enough such that f_n remains an immersion. This follows from compactness of D_n by again using the fact that the set of full rank $p \times n$ matrices is open in $\mathbb{R}^{p \times n}$. For the second part of 1) we claim that for almost all $b_n \in \mathbb{R}^p$ for which f_n is defined it holds that

$$f_n(x) = f_n(y) \text{ if and only if } f_{n-1}(x) = f_{n-1}(y) \text{ and } \lambda_n(x) = \lambda_n(y). \quad (\star)$$

The “if” part is clear. For the converse let us examine what $f_n(x) = f_n(y)$ implies. Let $Q_n := f_{n-1}^{-1}(W_n)$. Note that we may rewrite the definition of f_n as

$$f_n(x) := \begin{cases} w_n(w_n^{-1}(f_{n-1}(x)) + \lambda_n(x)b_n), & x \in Q_n, \\ f_{n-1}(x), & x \notin Q_n. \end{cases}$$

We now consider three cases. First of all if $x, y \notin Q_n$ then clearly $f_{n-1}(x) = f_{n-1}(y)$ and $\lambda_n(x) = \lambda_n(y) = 0$. Next assume $x \in Q_n$ and $y \notin Q_n$. Then $f_n(x) \in W_n$ and $f_n(y) = f_{n-1}(y) \notin W_n$, so this case can not happen. Now the final and only interesting case is when $x, y \in Q_n$. If moreover $\lambda_n(x) = \lambda_n(y)$ it follows immediately that $f_{n-1}(x) = f_{n-1}(y)$, so assume $\lambda_n(x) \neq \lambda_n(y)$. Rearranging terms it then follows from $f_n(x) = f_n(y)$ that necessarily

$$b_n = \frac{w_n^{-1}(f_{n-1}(x)) - w_n^{-1}(f_{n-1}(y))}{\lambda_n(y) - \lambda_n(x)}.$$

We can avoid this by choosing b_n as follows. Consider the open subset

$$G_n := \{(x, y) \in Q_n \times Q_n \mid \lambda_n(x) \neq \lambda_n(y)\} \subseteq Q_n \times Q_n$$

and define the smooth map

$$\gamma_n: G_n \rightarrow \mathbb{R}^p, (x, y) \mapsto \frac{w_n^{-1}(f_{n-1}(x)) - w_n^{-1}(f_{n-1}(y))}{\lambda_n(y) - \lambda_n(x)}.$$

Since $\dim(G_n) = 2 \dim(M) < p$ the image of γ_n is a set of measure zero. So $(\star)$ holds for all b_n not contained in this set. Let us now choose such b_n to define f_n . Using $(\star)$ it is now easy to show that f_n is injective on C_n . To this end assume there exist $x, y \in C_n$ with $f_n(x) = f_n(y)$ and $x \neq y$. Applying $(\star)$ inductively shows

$$f_{i-1}(x) = f_{i-1}(y) \text{ and } \lambda_i(x) = \lambda_i(y)$$

for all $1 \leq i \leq n$. The first equation implies in particular that $f(x) = f(y)$, such that x and y cannot belong to the same B_i as $f|_{B_i}$ is injective by property (v). Moreover if $x \in B_i$ for some $1 \leq i \leq n$ we obtain from the second equation together with property (i) that also $y \in B_i$, which contradicts what has just been shown. So both x and y have to belong to the set $C_0 = \bigcup_{-\infty \leq i \leq 0} B_i$, which is contained in U by property (iv). But f is assumed to be injective on U contradicting $x \neq y$. This shows 1).

As in the proof of Theorem 11.3 it now follows that

$$g := \lim_{n \rightarrow \infty} f_n$$

is a well-defined smooth immersion, which agrees with f on A and $g \in \mathcal{N}$. To see that g is also injective let $x, y \in M$ with $g(x) = g(y)$. Since the B_i cover M there exists some $n_0 > 0$ with $x, y \in C_{n_0}$. Furthermore there exists $n_1 \geq n_0$ with $g(x) = f_{n_1}(x)$ and $g(y) = f_{n_1}(y)$. From 1) we then obtain $x = y$. This completes the proof of Theorem 11.4 in the case $\partial N = \emptyset = \partial M$.

We now explain how to generalize the proof to the case, where M and N are allowed to have boundary. The first problem arises in the definition of f_n , since $f_{n-1}(x) + \lambda_n(x)b_n$ may not be contained in the domain of the chart w_n anymore. We can however easily resolve this by choosing b_n in the half space $\mathbb{H}^p \subseteq \mathbb{R}^p$ instead. The second problem is that $Q_n \times Q_n$ may not be a smooth manifold anymore (only one with corners). However we can still argue that the image of γ_n has measure zero. To this end note that both $\lambda_n|_{Q_n}$ and $\phi_n := w_n^{-1} \circ f_{n-1}|_{Q_n}$ are smooth maps with values in some euclidian space. So using a partition of unity they can be extended to smooth maps

$$\lambda'_n: Q'_n \rightarrow \mathbb{R}, \quad \phi'_n: Q'_n \rightarrow \mathbb{R}^p$$

defined on some boundaryless manifold Q'_n containing Q_n as a codimension 0 submanifold (we can for example choose Q'_n as the double of Q_n). Then the image of

$$\gamma'_n: G'_n \rightarrow \mathbb{R}^p, \quad (x, y) \mapsto \frac{\phi'_n(x) - \phi'_n(y)}{\lambda'_n(y) - \lambda'_n(x)}$$

defined on

$$G'_n := \{(x, y) \in Q'_n \times Q'_n \mid \lambda'_n(x) \neq \lambda'_n(y)\}$$

has measure zero. Since the image of γ_n is contained in the image of γ'_n , the claim follows. The rest of the proof goes through unchanged. $\square$

Next we proof Theorem 11.7 about transversality. As for the proof of Theorem 11.3 we start with a local result.

Lemma 11.13. *Let M be a manifold with boundary, $f: M \rightarrow \mathbb{H}^p$ a smooth function, $S \subseteq \mathbb{H}^p$ a submanifold with boundary and $\lambda: M \rightarrow (0, \infty)$ a positive function. Then for almost all vectors $b \in \mathbb{H}^p$ the map $f_b: M \rightarrow \mathbb{H}^p$, $x \mapsto f(x) + \lambda(x)b$ satisfies*

$$f_b \pitchfork S \text{ and } \partial f_b \pitchfork S.$$

Proof. Let $B := \text{int}(\mathbb{H}^p)$ be the interior of the half space. Then the map

$$F: M \times B \rightarrow \mathbb{H}^p, \quad (x, b) \mapsto f(x) + \lambda(x)b$$

is clearly a submersion. We now proof a version of the parametric transversality theorem which holds in this case. Using a partition of unity we can extend F to some map $F': M' \times B \rightarrow \mathbb{R}^p$, where M' is some boundaryless manifold containing M as a codimension 0 submanifold. Since F' is a submersion on $M \times B$ we can find some open neighbourhood $U \subseteq M' \times B$ of $M \times B$ on which F' is a submersion. Let

$$G := F'|_U: U \rightarrow \mathbb{R}^p$$

denote this submersion. U is a boundaryless manifold so by Lemma 1.5 the preimage $W := G^{-1}(S) \subseteq U$ is a submanifold (with boundary). Now consider the map

$$\pi: W \hookrightarrow M' \times B \xrightarrow{\text{proj.}} B,$$

where $W \hookrightarrow M' \times B$ is the inclusion. We claim that $f_b \pitchfork S$ for all regular values $b \in B$. So let $b \in B$ be such a regular value and let $x \in f_b^{-1}(S) \subseteq M$. We have to show that any vector $w \in \mathbb{R}^p$ can be written as

$$w = v + d_x f_b(y) \quad (16)$$

for some $v \in T_{f_b(x)}S = T_{F(x,b)}S$ and $y \in T_x M$. Since F is a submersion we can find $(y_1, z_1) \in T_x M \oplus T_b B$ with

$$w = d_{(x,b)} F(y_1, z_1). \quad (17)$$

Moreover as b is a regular value of π and $(x, b) \in W$ we can find $(y_2, z_2) \in T_{(x,b)}W \subseteq T_x M' \oplus T_b B = T_x M \oplus T_b B$ with

$$z_1 = d_{(x,b)} \pi(y_2, z_2).$$

Since $d_{(x,b)} \pi$ is just the restriction of the projection $T_x M \oplus T_b B \rightarrow T_b B$ it follows that $z_1 = z_2$. So by linearity we can write (17)

$$w = d_{(x,b)} F(y_1 - y_2, 0) + d_{(x,b)} F(y_2, z_1).$$

The first term is just $d_x f_b(y_1 - y_2)$. The second term may also be written as $d_{(x,b)} G(y_2, z_1)$ and since G restricts to a smooth map $G|_W: W \rightarrow S$ and $(y_2, z_1) \in T_{(x,b)}W$ we conclude

$$d_{(x,b)} G(y_2, z_1) \in T_{G(x,b)}S = T_{f_b(x)}S.$$

Hence (16) follows by setting $y := y_1 - y_2$ and $v := d_{(x,b)} G(y_2, z_1)$. So by Sard's Theorem $f_b \pitchfork S$ holds for almost all $b \in B$. The same argument applied to the submersion ∂F shows that $\partial f_b \pitchfork S$ for almost all $b \in B$. In fact this can be seen as a special case of what has just been shown, since ∂M is a manifold without boundary already, so we don't even have to worry about extending ∂F first. $\square$

Proof of Theorem 11.7. Again using the same notation as in the proof of Theorem 11.3 let $\mathcal{N} := \mathcal{N}^k(f, \Phi, \Psi, K, \varepsilon) \cap C^\infty(M, N)$ be some open neighbourhood of f . Let $U \subseteq M$ be some open neighbourhood of A , such that $f \pitchfork S$ on U and $\partial f \pitchfork S$ on $U \cap \partial M$. Such a neighbourhood exists due to the following fact: If f is transversal to S at some point $x \in M$, then it is also transversal to S in a whole neighbourhood of x . Applying this to both f and ∂f one can easily construct such U . We now choose the following data: A locally finite cover $(D_i)_{i \in \mathbb{Z}}$ of M of compact sets $D_i \subseteq M$, open sets $B_i \subseteq M$ which still cover M , charts $(W_i, w_i)_{i \in \mathbb{Z}}$ of N as well as cut-off functions $\lambda_i: M \rightarrow [0, 1]$, such that

- (i) $\lambda_i^{-1}((0, 1]) = B_i$,
- (ii) D_i contains the support of λ_i in its interior,
- (iii) $f(D_i) \subseteq W_i$,
- (iv) $D_i \subseteq U$ for $i \leq 0$ and $D_i \subseteq M - A$ for $i > 0$.

We now inductively define a sequence $f_n: M \rightarrow N$, $n \geq 0$, with $f_0 := f$, satisfying the following properties.

- 1) $f_n \pitchfork S$ at all points in $C_n := \bigcup_{-\infty \leq i \leq n} B_i$ and $\partial f_n \pitchfork S$ at all points in $\partial C_n = C_n \cap \partial M$,
- 2) $f_n = f_{n-1}$ outside D_n ,
- 3) $f_n(D_i) \subseteq W_i$ for all $i \in \mathbb{Z}$,
- 4) $f_n(K_\alpha) \subseteq V_\alpha$ for all $\alpha \in J$ and the coordinate representations $f_n^\alpha := \psi_\alpha^{-1} \circ f_n \circ \phi_\alpha$ of f_n satisfy

$$\|f_n^\alpha - f_{n-1}^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} < \frac{\varepsilon_\alpha}{2^n}.$$

For this theorem we proof the general case where both M and N may have non-empty boundary right away. Clearly $f_0 = f$ satisfies these properties and we define f_n using f_{n-1} by

$$f_n(x) := \begin{cases} w_n(w_n^{-1}(f_{n-1}(x)) + \lambda_n^2(x)b_n), & x \in D_n, \\ f_{n-1}(x), & x \notin D_n \end{cases}$$

for some suitable vector $b_n \in \mathbb{H}^p$, which we choose next. Again it follows as before that for sufficiently small b_n , f_n is a well defined smooth map satisfying 2), 3) and 4). Using Lemma 11.13 we can moreover choose b_n , such that $f_n \pitchfork S$ on B_n and $\partial f_n \pitchfork S$ on ∂B_n , since $\lambda_n^2 > 0$ on B_n . From the chain rule we obtain $d_x \lambda_n^2 = 0$ for all points $x \in M - B_n$, such that

$$d_x f_n = d_x f_{n-1}.$$

Thus also $d_x \partial f_n = d_x \partial f_{n-1}$ for $x \in \partial M - \partial B_n$. Hence f_n is transversal to S at all points $x \in M - B_n$ where f_{n-1} is transversal to S and the same relationship holds for ∂f_n and ∂f_{n-1} at all $x \in \partial M - \partial B_n$. In particular this transversality condition for f_n still holds for points in $C_{n-1} - B_n$ and for ∂f_n it still holds for points in $\partial C_{n-1} - \partial B_n$. From this 1) follows.

As before it now follows that

$$g := \lim_{n \rightarrow \infty} f_n$$

is a well-defined smooth map, which agrees with f on A and $g \in \mathcal{N}$. Moreover both g and ∂g are transversal to S by 1) since the B_i cover M . $\square$

Finally we proof Proposition 11.9. In the proof we will repeatedly make use of the following technical lemma. The precise formulas in this lemma may look strange at first but it will become clear in the proof of Proposition 11.9 where they come from.

Lemma 11.14. *Suppose we are given open sets $U, U' \subseteq \mathbb{H}^n, W, W' \subseteq \mathbb{H}^m, V \subseteq W$ compact sets $K \subseteq U, L \subseteq V$, a smooth function $\lambda: U' \rightarrow [0, 1]$, diffeomorphisms $\phi: U' \rightarrow U$ and $\psi: W \rightarrow W'$ as well as constants $\varepsilon > 0$ and $C > 0$. Then there exists $\delta > 0$ with the following property. Let $p, q: U_K \rightarrow W$ be smooth functions defined on some neighbourhood $U_K \subseteq U$ of K , which can depend on the functions p, q . Suppose that q satisfies $q(K) \subseteq L$ and $\|q\|_{C^k(K)} \leq C$. Then if*

$$\|p - q\|_{C^k(K)} < \delta$$

we have

$$(i) \ ((1 - \lambda)p + \lambda q)(K) \subseteq V \text{ and}$$

$$\|\psi((1 - \lambda)(p \circ \phi) + \lambda(q \circ \phi)) - \psi \circ p \circ \phi\|_{C^k(\phi^{-1}(K))} < \varepsilon.$$

$$(ii) \ ((1 - \lambda)(\psi \circ p) + \lambda(\psi \circ q))(K) \subseteq \psi(V) \text{ and}$$

$$\|\psi^{-1}((1 - \lambda)(\psi \circ p) + \lambda(\psi \circ q)) - q\|_{C^k(K)} < \varepsilon.$$

Proof. We first choose δ , such that $((1 - \lambda)p + \lambda q)(K) \subseteq V$ holds. By compactness of $L \subseteq V$ there exists some radius $r > 0$ depending only on L and V with the property that any ball of radius r around a point in L is contained in V . So choosing $\delta < r$ it follows from $q(K) \subseteq L$ and $0 \leq \lambda \leq 1$, that $(1 - \lambda)p(x) + \lambda q(x) \in V$ for any point $x \in K$. In fact the same argument shows that if we fix some compact neighbourhood $\tilde{L} \subseteq W$ of L we can choose δ sufficiently small, such that $p(K) \subseteq \tilde{L}$ is automatically satisfied. We will make use of this fact later in the proof.

We will now repeatedly apply Lemma 11.11 to make the desired norm in (i) small. First of all we can choose δ to make

$$\|p \circ \phi - q \circ \phi\|_{C^k(\phi^{-1}(K))} = \|(p - q) \circ \phi\|_{C^k(\phi^{-1}(K))}$$

arbitrarily small by applying part (ii) of Lemma 11.11, where $p - q$ plays the role of θ . Applying part (iii) of Lemma 11.11 we can thus also make

$$\|\lambda(p \circ \phi - q \circ \phi)\|_{C^k(\phi^{-1}(K))}$$

arbitrarily small. We now want to apply part (i) of Lemma 11.11 to

$$\begin{aligned} r &= p \circ \phi + \lambda(p \circ \phi - q \circ \phi) \\ s &= p \circ \phi \end{aligned}$$

to conclude that

$$\|\psi(p \circ \phi + \lambda(p \circ \phi - q \circ \phi)) - \psi \circ p \circ \phi\|_{C^k(\phi^{-1}(K))},$$

can be made arbitrarily small (this is the norm we are interested in just slightly rewritten). To apply the Lemma the assumption $(p \circ \phi)(\phi^{-1}(K)) = p(K) \subseteq \tilde{L}$ for some compact set $\tilde{L} \subseteq W$, which is fixed prior, is still missing. But as mentioned before we can choose $\tilde{L}$ as any fixed compact neighbourhood of L and if we then choose δ sufficiently small we automatically only consider functions p with $p(K) \subseteq \tilde{L}$. Hence we can apply the Lemma. The resulting δ now may of course depend on $\tilde{L}$ but not on p or q , which is what we are interested in.

Part (ii) of the Lemma can be shown analogously. In fact this part is slightly easier because the argument involving $\tilde{L}$ given above is not even needed here. $\square$

We can now proof Proposition 11.9.

Proof of Proposition 11.9. Choose a locally finite cover of charts $(O_i, o_i)_{i \in \mathbb{N}}$ of M , compact subsets $D_i \subseteq O_i$, subsets $B_i \subseteq D_i$ which still cover M , charts $(W_i, w_i)_{i \in \mathbb{N}}$ of N as well as cut-off functions $\lambda_i: M \rightarrow [0, 1]$, such that

- (i) $\lambda_i = 1$ in a neighbourhood of B_i ,
- (ii) D_i contains the support of λ_i in its interior,
- (iii) $f(D_i) \subseteq W_i$,
- (iv) $w_i^{-1}(W_i) \subseteq \mathbb{H}^p$ is convex.

Let us introduce some more notation. Consider the following data:

- The locally finite cover of charts $\mathbf{O} := (O_i, o_i)_{i \in \mathbb{N}}$ of M ,
- The family of charts $\mathbf{W} := (W_i, w_i)_{i \in \mathbb{N}}$ of N ,
- The family of compact sets $\mathbf{D} := (D_i)_{i \in \mathbb{N}}$,
- Some sequence $\Delta = (\delta_i)_{i \in \mathbb{N}}$ of positive numbers.

Then we denote by

$$\mathcal{V}(\Delta) := \mathcal{N}^k(f, \mathbf{O}, \mathbf{W}, \mathbf{D}, \Delta) \cap C^\infty(M, N)$$

the neighbourhood of f determined by this data. Moreover we shall assume that $\mathcal{U}$ is of the form

$$\mathcal{U} = \mathcal{N}^k(f, \Phi, \Psi, K, \varepsilon) \cap C^\infty(M, N),$$

using the same notation as in the other proofs. Now the goal at hand is to find a sufficiently small neighbourhood $\mathcal{V} \subseteq \mathcal{U}$ of f , such that the Proposition holds for any $g \in \mathcal{V}$. To construct this neighbourhood we inductively define numbers $\delta_n \in (0, 1)$, $n \geq 0$, with the following property. For $\Delta_n := (\delta_1, \dots, \delta_n, 1, 1, \dots)$ and $g \in \mathcal{V}_n := \mathcal{V}(\Delta_n)$ we can recursively define homotopies h_t^k , $t \in [0, 1]$, $0 \leq k \leq n$ as follows. For $k = 0$ set $h_t^0 := f$. To construct h_t^k out of h_t^{k-1} choose a smooth function $\alpha: [0, 1] \rightarrow [0, 1]$ with $\alpha = 0$ on $[0, \frac{1}{4})$ and $\alpha = 1$ on $(\frac{3}{4}, 1]$. On D_k we define h_t^k by a straight line homotopy from h_1^{k-1} to g in the coordinates of the chart w_k and then use the cut-off function λ_k to keep it constant outside of D_k . More precisely that is

$$h_t^k(x) := \begin{cases} w_k((1 - \alpha(t)\lambda_k(x))w_k^{-1}(h_1^{k-1}(x)) + \alpha(t)\lambda_k(x)w_k^{-1}(g(x))) & , x \in D_k \\ h_1^{k-1} & , x \notin D_k \end{cases}$$

These homotopies are well defined and satisfy the following properties for all $0 \leq k \leq n$.

- 1) $h_1^k = g$ on $C_k := \bigcup_{i=1}^k B_i$,
- 2) $h_t^k = h_1^{k-1}$ outside of D_k for all $t \in [0, 1]$,
- 3) $h_t^k(D_i) \subseteq W_i$ for all $t \in [0, 1]$,
- 4) $h_t^k(x) = f(x)$ for all $x \in A$ and $t \in [0, 1]$,
- 5) $h_t^k = h_1^{k-1}$ for $t \in [0, \frac{1}{4})$ and $h_t^k = h_1^k$ for $t \in (\frac{3}{4}, 1]$.

6) $h_t^k(K_\alpha) \subseteq V_\alpha$ for all $\alpha \in J$ and the coordinate expressions $h_t^{t,\alpha} := \psi_\alpha^{-1} \circ h_t^k \circ \phi_\alpha$ satisfy

$$\|h_t^{k,\alpha} - h_1^{k-1,\alpha}\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} < \frac{\varepsilon_\alpha}{2^k}$$

for all $t \in [0, 1]$.

The base case $n = 0$ is clear. So suppose we have already constructed $\delta_1, \dots, \delta_{n-1}$. Then for any $g \in \mathcal{V}_{n-1}$, h_t^n is well-defined since $h_1^{n-1}(D_n) \subseteq W_n$, $g(D_n) \subseteq W_n$ and $w_n^{-1}(W_n)$ is convex, such that

$$(1 - \alpha(t)\lambda_n(x))w_n^{-1}(h_1^{n-1}(x)) + \alpha(t)\lambda_n(x)w_n^{-1}(g(x))$$

is still contained in the domain of w_n . Properties 1), 2), 4) and 5) also clearly hold for $k = n$. For 3) and 6) to hold further constraints must be imposed on g . To this end we now construct $\delta_n \in (0, 1)$ such that for any $g \in \mathcal{V}_n$ both claims are fulfilled for $k = n$ under the additional assumption that

$$\|w_n^{-1} \circ h_1^{n-1} \circ o_n - w_n^{-1} \circ g \circ o_n\|_{C^k(o_n^{-1}(D_n))} \quad (18)$$

is sufficiently small. We achieve this by applying Lemma 11.14. Let us begin with 3). For some fixed $i \in \mathbb{N}$ we apply the first part of Lemma 11.14 (i) to

- $p = w_n^{-1} \circ h_1^{n-1} \circ o_n$,
- $q = w_n^{-1} \circ g \circ o_n$,
- $\lambda = \alpha(t)\lambda_n \circ o_n$ for some fixed $t \in [0, 1]$,
- $K = o_n^{-1}(D_n \cap D_i)$,
- $V = w_n^{-1}(W_n \cap W_i)$.

Then we can write

$$(w_n^{-1} \circ h_t^n \circ o_n)(x) = (1 - \lambda(x))p(x) + \lambda(x)q(x)$$

for $x \in K$ and if (18) is sufficiently small then in particular $\|p - q\|_{C^k(K)}$ can be made arbitrarily small. To apply the Lemma we still need to choose the compact set $L \subseteq V$, such that q always satisfies $q(K) \subseteq L$ no matter which precise g we used to define q . We fix L as some compact neighbourhood of $(w_n^{-1} \circ f \circ o_n)(K) \subseteq V$. Then we can choose δ_n so small that $q(K) \subseteq L$ for any $g \in \mathcal{V}_n$. Now Lemma 11.14 tells us that $((1 - \lambda)p + \lambda q)(K) \subseteq V$ for (18) sufficiently small, which is equivalent to $h_t^n(D_n \cap D_i) \subseteq W_i$. Of course we can then also achieve this for finitely many $i \in \mathbb{N}$ if (18) and δ_n are sufficiently small. But by a local finiteness argument only finitely many D_i intersect D_n and since $h_t^{n-1}(D_i) \subseteq W_i$ already holds for all $i \in \mathbb{N}$ and $h_t^n = h_1^{n-1}$ outside D_n we conclude that we can also achieve $h_t^n(D_i) \subseteq W_i$ for all $i \in \mathbb{N}$.

For 6) we apply Lemma 11.14 to the same p, q, λ and

- $\phi = o_n^{-1} \circ \phi_\alpha$,
- $\psi = \psi_\alpha^{-1} \circ w_n$,
- $K = o_n^{-1}(D_n \cap K_\alpha)$,
- $V = w_n^{-1}(W_n \cap V_\alpha)$,

with the obvious domains and codomains. An analogous argument as above shows that $h_t^n(K_\alpha) \subseteq V_\alpha$ holds for all $\alpha \in J$ if (18) and δ_n are sufficiently small. Furthermore we can write

$$h_t^{n,\alpha}(x) = \psi((1 - \lambda(x))p(\phi(x)) + \lambda(x)q(\phi(x)))$$

and $h_1^{n-1,\alpha}(x) = (\psi \circ p \circ \phi)(x)$ for $x \in \phi_\alpha^{-1}(D_n \cap K_\alpha) = \phi^{-1}(K)$. So Lemma 11.14 (i) tells us that

$$\|h_t^{n,\alpha} - h_1^{n-1,\alpha}\|_{C^k(\phi_\alpha^{-1}(D_n \cap K_\alpha))} < \frac{\varepsilon_\alpha}{2^n}$$

for (18) and δ_n sufficiently small. But again since $h_t^n = h_1^{n-1}$ outside D_n this is already equivalent to

$$\|h_t^{n,\alpha} - h_1^{n-1,\alpha}\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} < \frac{\varepsilon_\alpha}{2^n}.$$

Thus properties 3) and 6) are established if (18) becomes sufficiently small. Similiarly to the argument above we now choose δ_n even smaller, such that (18) is sufficiently small for any $g \in \mathcal{V}_n$ under the additional assumption that

$$\|w_n^{-1} \circ h_1^{n-2} \circ o_n - w_n^{-1} \circ g \circ o_n\|_{C^k(o_n^{-1}(D_n))} \quad (19)$$

is sufficiently small. For this we again apply Lemma 11.14, this time part (ii), to

- $p = w_n^{-1} \circ h_1^{n-2} \circ o_n$,
- $q = w_n^{-1} \circ g \circ o_n$,
- $\lambda = \lambda_{n-1} \circ o_n$,
- $K = o_n^{-1}(D_n \cap D_{n-1})$,
- $V = w_n^{-1}(W_n \cap W_{n-1})$,
- $\psi = w_{n-1}^{-1} \circ w_n$.

Again if δ_n is sufficiently small we obtain $q(K) \subseteq L$ for any $g \in \mathcal{V}_n$, where $L \subseteq V$ is some fixed compact neighbourhood of $(w_n^{-1} \circ f \circ o_n)(K) \subseteq V$. Write

$$(w_n^{-1} \circ h_1^{n-1} \circ o_n)(x) = \begin{cases} \psi^{-1}((1 - \lambda(x))(\psi \circ p)(x) + \lambda(x)(\psi \circ q)(x)) & , x \in K \\ (w_n^{-1} \circ h_1^{n-2} \circ o_n)(x) & , x \in o_n^{-1}(D_n) - K \end{cases}$$

As before we see that if (19) is sufficiently small then in particular $\|p - q\|_{C^k(K)}$ is sufficiently small. Thus we can conclude from Lemma 11.14 that also (18) is sufficiently small. Proceeding inductively we can construct δ_n , such that (19) and hence (18) becomes arbitrarily small for any $g \in \mathcal{V}_n$ as long as

$$\|w_n^{-1} \circ h_1^0 \circ o_n - w_n^{-1} \circ g \circ o_n\|_{C^k(o_n^{-1}(D_n))} \quad (20)$$

is also sufficiently small. But since $h_1^0 = f$ we have that (20) $< \delta_n$ by definition of the neighbourhood $\mathcal{V}_n$. So we can simply choose δ_n , such that (20) is as small as desired for all $g \in \mathcal{V}_n$. This completes the construction of the δ_n 's.

Now we actually define the homotopy asserted by the propostion, which we aim to proof. Let $\Delta := (\delta_1, \delta_2, \delta_3, \dots)$ for the $\delta_n \in (0, 1)$ constructed above and set

$$\mathcal{V} := \mathcal{V}(\Delta) = \bigcap_{n=0}^{\infty} \mathcal{V}_n.$$

Then for any $g \in \mathcal{V}$ the homotopies h_t^k are well defined for all $k \geq 0$ and satisfy properties 1) - 6) for all $k \geq 0$. For better readability in what follows let us write $h^k(-, t) := h_t^k$. Moreover let $a_n := \sum_{k=1}^n 2^{-k}$ and $I_n := [a_n, a_{n+1}]$. For $t \in I_n$ the homotopy h_t is now defined by h_t^n with the time parameter t rescaled to the interval I_n , that is

$$h_t(x) := \begin{cases} h^n(x, 2^{n+1}(t - a_n)) & , t \in I_n \\ g(x) & , t = 1. \end{cases}$$

This is well-defined for $t \in I_{n-1} \cap I_n$ since $h_1^{n-1} = h_0^n$. We claim moreover that h_t defines a smooth homotopy. For $t_0 \neq 1$, t_0 is either an interior point or a boundary point of some Interval I_n . If it is an interior point, then $h_t = h^n(-, 2^{n+1}(t - a_n))$ for t in some neighbourhood of t_0 . If it is a boundary point, say $t_0 = a_{n+1}$, then $h_t = h_1^n$ in some neighbourhood of t_0 by 5). If $t_0 = 1$ we obtain from 1) using a local finiteness argument that for any $x_0 \in M$ there exists some neighbourhood $U(x_0) \subseteq M$ of x_0 , such that $h_t(x) = h_t^n(x) = g(x)$ for n sufficiently large, $x \in U(x_0)$ and t in some neighbourhood of t_0 . This shows smoothness.

It is then clear from the definition that $h_0 = f$ and $h_1 = g$. Furthermore $h_t(x) = f(x)$ for all $x \in A$ by 4). Finally we have to show that $h_t \in \mathcal{U}$ for all $t \in [0, 1]$. Again using 1) as well as compactness of K_α it follows that $g = h_t^n$ in a neighbourhood of K_α for sufficiently large n . So since $h_t^n(K_\alpha) \subseteq V_\alpha$ holds for all $\alpha \in J$ and $t \in [0, 1]$ we also obtain $h_t(K_\alpha) \subseteq V_\alpha$ for all $\alpha \in J$ and

$t \in [0, 1]$. Now given any $t < 1$ we can write $h_t = h_s^n$ for some $n \geq 0$ and $s = 2^{n+1}(t - a_n)$. Write $h_t^\alpha := \psi_\alpha^{-1} \circ h_t \circ \phi_\alpha$ and $f^\alpha := \psi_\alpha^{-1} \circ f \circ \phi_\alpha$. Using 6) we compute

$$\begin{aligned} \|h_t^\alpha - f^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} &= \|h_s^{n,\alpha} - f^\alpha\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} \\ &\leq \|h_s^{n,\alpha} - h_1^{n-1,\alpha}\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} + \sum_{i=1}^{n-1} \|h_1^{i,\alpha} - h_1^{i-1,\alpha}\|_{C^k(\phi_\alpha^{-1}(K_\alpha))} \\ &< \frac{\varepsilon_\alpha}{2^n} + \sum_{i=1}^{n-1} \frac{\varepsilon_\alpha}{2^i} \\ &< \varepsilon_\alpha \end{aligned}$$

For $t = 1$ the same follows by again using that $g = h_t^n$ in a neighbourhood of K_α for sufficiently large n . This shows $h_t \in \mathcal{U}$ for all $t \in [0, 1]$ and completes the proof. $\square$

11.2 An equivalent definition of the Whitney Topologies

In this section we want to show that the Whitney C^k -topologies on $C^k(X, Y)$ as defined at the beginning of section 11.1 agree with the definition given in [GG12] based on jet bundles.

For the moment, let us call the topology defined in section 11.1 the “local chart topology” and the topology defined in [GG12] using jet bundles the “jet topology”. First note that in both cases the topology on $C^\infty(X, Y)$ is defined as the topology induced by all inclusion maps $C^\infty(X, Y) \rightarrow C^k(X, Y)$, so in the following we assume k to be finite. Before showing equivalence we introduce some notation used in the proof. Suppose we are given the data $(U_i, \phi_i), (V_i, \psi_i), K_i, \varepsilon_i$ for an open neighbourhood $\mathcal{N}^k(f, \Phi, \Psi, K, \varepsilon)$ as explained in section 11.1. We abbreviate

$$\begin{aligned} \mathcal{N}_i &:= \mathcal{N}^k(f, (U_i, \phi_i), (V_i, \psi_i), K_i, \varepsilon_i), \\ \mathcal{N} &:= \mathcal{N}^k(f, \Phi, \Psi, K, \varepsilon) \end{aligned}$$

and $\tilde{U}_i := \phi_i^{-1}(U_i)$, $\tilde{K}_i := \phi_i^{-1}(K_i)$, $\tilde{V}_i := \psi_i^{-1}(V_i)$, $\tilde{f} := \psi_i^{-1} f \phi_i$. Define

$$\tilde{W}_i := \{(x, y, (v_\alpha)_\alpha) \in \tilde{U}_i \times \tilde{V}_i \times \mathbb{R}^{d(n,k)} \mid \|\tilde{f}(x) - y\| < \varepsilon_i, \|\partial^\alpha \tilde{f}(x) - v_\alpha\| < \varepsilon_i \text{ for all } \alpha\},$$

where $\alpha \in \mathbb{N}_0^n$ ranges over all multiindices with $1 \leq |\alpha| \leq k$, $v_\alpha \in \mathbb{R}^n$ and $d(n, k)$ is the total number of such multiindices as defined in the proof of Proposition 11.19. The charts ϕ_i and ψ_i define a chart

$$\tau_i: \tilde{U}_i \times \tilde{V}_i \times \mathbb{R}^{d(n,k)} \xrightarrow{\cong} J^k(U_i, V_i) \subseteq J^k(X, Y)$$

which we use to define

$$W_i := \tau(\tilde{W}_i).$$

Clearly $W_i \subseteq J^k(X, Y)$ is open. Moreover W_i is constructed in such a way that

$$g \in \mathcal{N}_i \Leftrightarrow j^k g(K_i) \subseteq W_i. \quad (21)$$

We now show that any open set in the local chart topology is also open in the jet topology. To this end consider

$$F_i := W_i \cup \alpha^{-1}(X \setminus K_i),$$

where $\alpha: J^k(X, Y) \rightarrow X$ is the source map. Then from (21) we get

$$g \in \mathcal{N}_i \Leftrightarrow j^k g(X) \subseteq F_i \Leftrightarrow g \in M(F_i)$$

Since

$$\mathcal{N} = \bigcap_{i \in I} \mathcal{N}_i$$

we can show that $\mathcal{N} \subseteq C^k(X, Y)$ is open in the jet topology if we can show that

$$F := \bigcap_{i \in I} F_i \subseteq J^k(X, Y)$$

is open, because $\bigcap_{i \in I} M(F_i) = M(F)$. For this we will use the local finiteness of the sets U_i . Let $\sigma \in W$ be a jet. Then around its source $\alpha(\sigma) \in X$ we can find an open neighbourhood

$$A \subseteq X,$$

such that A only intersects finitely many (possibly none) of the U_i , say $U_{i_1}, \dots, U_{i_N}$. Then

$$B := W_{i_1} \cap \dots \cap W_{i_N} \cap \alpha^{-1}(A) \subseteq J^k(X, Y)$$

is open and for any $\sigma' \in B$ we have

$$\alpha(\sigma') \in A \subseteq X \setminus K_i$$

for all $i \neq i_1, \dots, i_N$, thus $\sigma \in F_i$ for all $i \in I$ and hence F is open.

Now we show that for an open set $T \subseteq J^k(X, Y)$ the set $M(T) \subseteq C^k(X, Y)$ is also open in the local chart topology. For $f \in M(T)$ choose charts (ϕ_i, U_i) of X , (ψ_i, V_i) of Y and compact sets $K_i \subseteq U_i$, such that the U_i are a locally finite open cover of X , the sets K_i still cover X and $f(K_i) \subseteq V_i$ for all $i \in I$. We have to find positive numbers $\varepsilon_i > 0$, such that the basic open neighbourhood $\mathcal{N}$ defined using this data is contained in $M(T)$. Suppose that for each $i \in I$ we can find $\varepsilon_i > 0$, such that

$$\tilde{W}_i \cap (\tilde{K}_i \times \tilde{V}_i \times \mathbb{R}^{d(n,k)}) \subseteq \tau_i^{-1}(T) \cap (\tilde{K}_i \times \tilde{V}_i \times \mathbb{R}^{d(n,k)}). \quad (22)$$

For any $g \in \mathcal{N}$ we have

$$(\tau^{-1} \circ j^k g)(K_i) \subseteq \tilde{W}_i \cap (\tilde{K}_i \times \tilde{V}_i \times \mathbb{R}^{d(n,k)})$$

and thus we get from (22)

$$j^k g(K_i) \subseteq T.$$

This holds for all $i \in I$ and since the K_i cover X we conclude that

$$j^k g(X) \subseteq T.$$

Thus $\mathcal{N} \subseteq M(T)$. To show the existence of these ε_i we invoke the following Lemma from point set topology.

Lemma 11.15. *Let D be a compact topological space, (Z, d) a metric space and $h: D \rightarrow Z$ continuous. Then for any open neighbourhood $S \subseteq D \times Z$ of the graph $\Gamma_h \subseteq D \times Z$ of h , there exists $\varepsilon > 0$, such that the ε -tube*

$$\{(x, z) \in D \times Z \mid d(h(x), z) < \varepsilon\}$$

is contained in S .

Proof. See Lemma 6.3. □

Applying this Lemma to

- $D = \tilde{K}_i$,
- $Z = \tilde{V}_i \times \mathbb{R}^{d(n,k)}$,
- $h = (\tilde{f}, (\partial^\alpha \tilde{f})_\alpha)$,
- $S = \tau_i^{-1}(T) \cap (\tilde{K}_i \times \tilde{V}_i \times \mathbb{R}^{d(n,k)})$

with the correct norm on $\mathbb{R}^{d(n,k)}$ yields the desired ε_i .

11.3 C^k -convergence

Let X, Y be smooth manifolds. In this section we discuss the concept of C^k -convergence for sequences of functions $(f_n) \subseteq C^k(X, Y)$ with respect to the Whitney topologies. We will be using the definition of the Whitney topologies based on jet bundles. The notation and terminology used will be the same as in [GG12]. First we remark that X or Y is also allowed to have boundary. The definition of the jet bundles $J^k(X, Y)$ and the Whitney topologies works exactly the same⁸. Let us begin by understanding the C^0 -topology.

Lemma 11.16. *Let X, Y be smooth manifolds with X compact. Then the Whitney C^0 -topology (as described in [GG12]) and the exponential topology (as described in [Ste23b]) on $C^0(X, Y)$ agree.*

Proof. We identify $J^0(X, Y) \cong X \times Y$. Then by definition the sets of the form

$$M(W) := \{f \in C^0(X, Y) \mid j^0 f(X) = \text{Graph}(f) \subseteq W\}$$

for $W \subseteq X \times Y$ open are a basis of the C^0 -topology on $C^0(X, Y)$. To see that this topology agrees with the exponential topology we show that it satisfies the correct universal property. So let S be any topological space. We have to show that a map

$$F: S \rightarrow C^0(X, Y)$$

is continuous if and only if

$$\tilde{F}: S \times X \rightarrow Y, (s, x) \mapsto F(s)(x)$$

is continuous. First assume F to be continuous. Let $V \subseteq Y$ be open and $(s_0, x_0) \in \tilde{F}^{-1}(V)$. Define $f_0 := F(s_0) \in C^0(X, Y)$, then

$$U := f_0^{-1}(V)$$

is open in X . and $x_0 \in U$. Choose some smaller open neighbourhood $U_0 \subseteq U$ of x_0 with $\overline{U_0} \subseteq U$ and set

$$W := U \times V \cup (X \setminus \overline{U_0}) \times Y \subseteq X \times Y.$$

By assumption $S_0 := F^{-1}(M(W)) \subseteq S$ is open and we have constructed W , such that $s_0 \in S_0$. But then

$$S_0 \times U_0 \subseteq S \times X$$

is an open neighbourhood of (s_0, x_0) and $S_0 \times U_0 \subseteq \tilde{F}^{-1}(V)$.

Now assume that $\tilde{F}$ is continuous. Let $W \subseteq X \times Y$ be open and $s_0 \in F^{-1}(M(W))$. Again define $f_0 := F(s_0) \in C^0(X, Y)$. Then $\text{Graph}(f_0) \subseteq W$ and around each $(x, f_0(x)) \in \text{Graph}(f_0)$ we can find a product neighbourhood $U_x \times V_x \subseteq W$, where $U_x \subseteq X$ is an open neighbourhood of x and $V_x \subseteq Y$ is an open neighbourhood of $f_0(x)$. After possibly shrinking U_x we may further assume that $\overline{U_x} \subseteq f_0^{-1}(V_x)$. Since X is compact we can cover $\text{Graph}(f_0)$ with finitely many such neighbourhoods $U_1 \times V_1, \dots, U_n \times V_n$. Now for each V_i we have that

$$\{s_0\} \times \overline{U_i} \subseteq \tilde{F}^{-1}(V_i).$$

But $\tilde{F}^{-1}(V_i) \subseteq S \times X$ is open by assumption and $\overline{U_i}$ is compact, so we can find open neighbourhoods $S_i \subseteq S$ of s_0 with

$$S_i \times \overline{U_i} \subseteq \tilde{F}^{-1}(V_i).$$

Set $S_0 := S_1 \cap \dots \cap S_n$. Then $S_0 \subseteq S$ is an open neighbourhood of s_0 and for any $(s, x) \in S_0 \times X$ there exists U_i with $x \in U_i$, hence

$$F(s)(x) = \tilde{F}(s, x) \in V_i.$$

It follows that $\text{Graph}(F(s)) \subseteq \bigcup_{i=1}^n U_i \times V_i \subseteq W$ for all $s \in S_0$ and thus $S_0 \subseteq F^{-1}(M(W))$. $\square$

⁸If both have boundary, $J^k(X, Y)$ will be a manifold with corners, so we might want to avoid that. However $J^k(X, Y)$ still carries a well-defined topology and I think this is all we need and the proofs in this section all go through the same. In [Hir12] at the end of Chapter 2.4 on page 63 the author briefly describes the definition of $J^k(X, Y)$ in this case.

Remark 11.17. If X is not compact the C^0 -topology will in general be finer than the exponential topology. To see this recall that a subbase for the exponential topology on $C^0(X, Y)$ is given by sets of the form

$$F(K, V) := \{f \in C^0(X, Y) \mid f(K) \subseteq V\}$$

for compact sets $K \subseteq X$ and open sets $V \subseteq Y$. Set

$$W := X \times V \cup (X \setminus K) \times Y.$$

Then $W \subseteq X \times Y$ is open and

$$F(K, V) = M(W).$$

Before giving an example where the C^0 -topology is strictly finer, we fix some terminology to differentiate between the two topologies on $C^0(X, Y)$. We shall use the term C_{exp}^0 -convergence to refer to convergence with respect to the exponential topology and reserve the term C^0 -convergence for convergence with respect to the Whitney C^0 -topology. Now consider the following example. Define a sequence

$$f_n := \text{const}_{\frac{1}{n}} : \mathbb{R} \rightarrow \mathbb{R}, \quad x \mapsto \frac{1}{n}$$

of constant continuous functions. We claim that

$$f_n \xrightarrow{C_{\text{exp}}^0} \text{const}_0.$$

To see this let $F(K, V) \subseteq C^0(\mathbb{R}, \mathbb{R})$ be any open neighbourhood of const_0 of the form described above. Then $0 \in V$ and so for n sufficiently large $\frac{1}{n} \in V$, which is equivalent to $f_n \in F(K, V)$. Since the sets of the form $F(K, V)$ are a subbase for the exponential topology, this shows the claim. However the sequence (f_n) does not C^0 converge to const_0 , since this would necessarily require for f_n to equal 0 outside a compact subset for sufficiently large n (see [GG12]).

Next we understand convergence in the C^k -topology. Before stating the main proposition we shall introduce another type of convergence which is better suited for non-compact domains, since in the non-compact case usual C^k -convergence is often too strong, because (as already statetd above)

$$f_n \xrightarrow{C^k} f$$

would already imply that f_n and f agree outside a compact subset (again see [GG12]).

Definition 11.18. Let X, Y be smooth manifolds and $f, f_n \in C^k(X, Y)$. We say that

$$f_n \xrightarrow{C_{\text{loc}}^k} f$$

if for every compact codimension 0-submanifold $K \subseteq X$ we have

$$f_n|_K \xrightarrow{C^k} f|_K.$$

Note that for compact domains C_{loc}^k -convergence and C^k -convergence coincide, so we are actually interested in understanding C_{loc}^k -convergence. The following two propositions give a useful characterization of C_{loc}^k -convergence to work with in practice.

Proposition 11.19. Let $D \subseteq \mathbb{R}^n$ be a compact codimension 0 submanifold, $V \subseteq \mathbb{R}^m$ open and $f, f_n \in C^k(D, V)$. Then

$$f_n \xrightarrow{C^k} f$$

if and only if for every multiindex $\alpha \in \mathbb{N}_0^n$ with $0 \leq |\alpha| \leq k$

$$\frac{\partial^{|\alpha|} f_n}{\partial x^\alpha} \xrightarrow{\text{uniformly}} \frac{\partial^{|\alpha|} f}{\partial x^\alpha}.$$

Proposition 11.20. Let X, Y be smooth manifolds and $f, f_n \in C^k(X, Y)$. Then the following statements are equivalent.

$$(i) \quad f_n \xrightarrow{C_{\text{loc}}^k} f,$$

$$(ii) \ j^k f_n \xrightarrow{C_{\text{loc}}^0} j^k f,$$

(iii) For all charts $\phi: U \rightarrow U' \subseteq X$ and $\psi: V \rightarrow V' \subseteq Y$ with $f(U'), f_n(U') \subseteq V'$ (for n sufficiently large) we have

$$\psi^{-1} \circ f_n \circ \phi \xrightarrow{C_{\text{loc}}^k} \psi^{-1} \circ f \circ \phi,$$

(iv) For every $p \in X$ there exist charts $\phi: U \rightarrow U' \subseteq X$ around p and $\psi: V \rightarrow V' \subseteq Y$ around $f(p)$ with $f(U'), f_n(U') \subseteq V'$ (for n sufficiently large) such that

$$\psi^{-1} \circ f_n \circ \phi \xrightarrow{C_{\text{loc}}^k} \psi^{-1} \circ f \circ \phi.$$

Before proving the propositions we recall a few facts about the exponential topology not mentioned in [Ste23b], which will be used in the proofs.

Lemma 11.21. *Let X, Y, Z be topological spaces and $A \subseteq X, B \subseteq Y$ subspaces. Then the following holds (assuming the exponential topology exists on each space)*

(i) *The canonical map $C^0(X, B) \hookrightarrow C^0(X, Y)$ is an embedding*

(ii) *The restriction map $C^0(X, Y) \rightarrow C^0(A, Y)$ is continuous*

(iii) *The canonical bijection $C^0(X, Y) \times C^0(X, Z) \rightarrow C^0(X, Y \times Z)$ is a homeomorphism*

Proof. For (i) we show that the exponential topology on $C^0(X, B)$ agrees with the topology induced by the map $C^0(X, B) \hookrightarrow C^0(X, Y)$ by showing the corresponding universal property. So let

$$\begin{array}{ccc} & C^0(X, Y) & \\ & \uparrow F & \\ Z & \xrightarrow{f} & C^0(X, B) \end{array}$$

commute with F being continuous. We have to show that f is continuous, but by definition of the exponential topology this is equivalent to

$$Z \times X \rightarrow B, (z, x) \mapsto f(z)(x)$$

being continuous and continuity of F is equivalent to continuity of

$$Z \times X \rightarrow Y, (z, x) \mapsto F(z)(x),$$

so (i) follows. For (ii) note that the restriction map is continuous if and only if

$$C^0(X, Y) \times A \rightarrow Y, (f, x) \mapsto f|_A(x)$$

is continuous. But this is just the restriction of the evaluation map which is continuous by [Ste23b]. Similarly the map in (iii) is continuous if and only if

$$C^0(X, Y) \times C^0(X, Z) \times X \rightarrow Y \times Z, (f, g, x) \mapsto (f(x), g(x))$$

is. But again the continuity follows from the continuity of the evaluation map. $\square$

Proof of Proposition 11.19. In [GG12] it was shown that for compact domains C^k -convergence is equivalent to the k -jets converging uniformly with respect to any metric on the Jet-bundle compatible with its topology, that is

$$f_n \xrightarrow{C^k} f \Leftrightarrow j^k f_n \xrightarrow{\text{uniformly}} j^k f$$

Note that for compact domains uniform convergence and C_{exp}^0 -convergence agree (see [Ste23b]). Now let $d(n, k) = \frac{(n+k)!}{n!k!}$ be the total number of all partial derivatives ∂^α of order $1 \leq |\alpha| \leq k$ and let

$$\tau: D \times V \times \mathbb{R}^{d(n, k)} \xrightarrow{\cong} J^k(D, V)$$

be the canonical diffeomorphism. Then

$$j^k f_n \xrightarrow{C_{\text{exp}}^0} j^k f \Leftrightarrow j^k f_n \circ \tau \xrightarrow{C_{\text{exp}}^0} j^k f \circ \tau.$$

But a sequence of maps with values in a product C_{exp}^0 -converges if and only if each component C_{exp}^0 -converges. Since the components of $j^k f_n \circ \tau$ and $j^k f \circ \tau$ are precisely the partial derivatives of f_n and f up to order k the statement follows (after again applying the equivalence of uniform and C_{exp}^0 -convergence on compact domains). $\square$

Proof of Proposition 11.20. Let $K \subseteq X$ be any compact codimension 0 submanifold. It then has been shown in [GG12] that

$$f_n|_K \xrightarrow{C^k} f|_K$$

is equivalent to

$$j^k f_n|_K \xrightarrow{\text{uniformly}} j^k f|_K$$

with respect to any metric on $J^k(X, Y)$ compatible with the topology. But again uniform convergence on compact domains is equivalent to C_{exp}^0 -convergence, which in turn is equivalent to C^0 -convergence by Lemma 11.16. This shows the equivalence of (i) and (ii).

For (ii) $\Rightarrow$ (iii) let ϕ, ψ be as in (iii) and let $D \subseteq U$ be any compact codimension 0 submanifold. Then by Lemma 11.16 we again observe that C^0 -convergence of $j^k f_n$ on $\phi(D)$ is equivalent to C_{exp}^0 -convergence and since the exponential topology behaves nicely with respect to restriction in the codomain (see Lemma 11.21) we get that the restricted maps

$$j^k f_n|_{\phi(D)}: \phi(D) \rightarrow J^k(U', V')$$

C_{exp}^0 -converge to

$$j^k f|_{\phi(D)}: \phi(D) \rightarrow J^k(U', V'),$$

where $J^k(U', V') \subseteq J^k(X, Y)$ is considered an open subset. Under the diffeomorphisms

$$\begin{aligned} D &\xrightarrow{\cong} \phi(D), \\ U \times V \times \mathbb{R}^{d(n,k)} &\xrightarrow{\cong} J^k(U', V') \end{aligned}$$

induced by ϕ and ψ the maps $j^k f_n|_{\phi(D)}, j^k f|_{\phi(D)}$ correspond to maps

$$g_n, g: D \rightarrow U \times V \times \mathbb{R}^{d(n,k)}$$

and $g_n \xrightarrow{C_{\text{exp}}^0} g$. But a sequence of maps with values in a product C_{exp}^0 -converges if and only if each component converges (again see Lemma 11.21). Since the components of g_n are id_U and the partial derivatives of $\psi^{-1} \circ f_n \circ \phi$ of order less than k (and similarly for g) we conclude that

$$\left. \frac{\partial^\alpha (\psi^{-1} \circ f_n \circ \phi)}{\partial x^\alpha} \right|_D \xrightarrow{C_{\text{exp}}^0} \left. \frac{\partial^\alpha (\psi^{-1} \circ f \circ \phi)}{\partial x^\alpha} \right|_D$$

for all $\alpha \in \mathbb{N}_0^n$ with $0 \leq |\alpha| \leq k$. Now (iii) follows from Proposition 11.19 (and the equivalence of uniform and C_{exp}^0 -convergence on compact domain).

Clearly (iii) $\Rightarrow$ (iv), so let's assume (iv). Using the same notation and argument as above but reversed we see that

$$(\psi^{-1} \circ f_n \circ \phi)|_D \xrightarrow{C^k} (\psi^{-1} \circ f \circ \phi)|_D$$

if and only if

$$j^k f_n|_{\phi(D)} \xrightarrow{C_{\text{exp}}^0} j^k f|_{\phi(D)},$$

where we consider $j^k f_n|_{\phi(D)}$ and $j^k f|_{\phi(D)}$ as maps

$$\phi(D) \rightarrow J^k(X, Y).$$

Now choose any metric on $J^k(X, Y)$ compatible with the topology. Then C_{exp}^0 -convergence of $j^k f_n|_{\phi(D)}$ is equivalent to uniform convergence with respect to that metric. Thus we have shown

that around each point $p \in X$ there exists some neighbourhood $\phi(D)$ of p , such that $j^k f_n$ converges uniformly to $j^k f$ on $\phi(D)$. But any compact codimension 0 submanifold $K \subseteq X$ is covered by finitely many such neighbourhoods and it is easily seen that this already implies uniform convergence of $j^k f_n$ to $j^k f$ on all of K . Then again using the equivalence of uniform convergence and C_{exp}^0 -convergence on compact domains as well as Lemma 11.16 we conclude that

$$j^k f_n \xrightarrow{C_{\text{loc}}^0} j^k f.$$

□

12 Miscellaneous

12.1 Smoothness on arbitrary subsets

Let M, N be manifolds and $A \subseteq M$ an arbitrary subset. In [Lee11] the author defines what it means for a map

$$f: A \rightarrow N$$

to be smooth, namely

We call f smooth if around each $x \in A$ there exists a smooth extension $\tilde{f}: U \rightarrow N$ of f to some open neighbourhood $U \subseteq M$ of x (see [Lee11, p. 45]).

However this definition has a fatal flaw: if A happens to be a submanifold of M , a map $f: A \rightarrow N$ might be smooth in the usual sense (i.e. as a map between smooth manifolds) but not smooth in the sense of Lee's definition. Consider for example

Counterexample 12.1.

$$M := \mathbb{R}, N := [0, 1], A := [0, 1], \\ f: A \rightarrow N, t \mapsto t.$$

Then clearly f is smooth as a map between smooth manifolds, but around $0, 1 \in A$ it does not admit a smooth extension onto an open neighbourhood $U \subseteq M$, since the image of such an extension would fail to be contained within N .

We can however correct this flaw by defining:

Definition 12.2 (Extension-smooth). Let M^n, N^m be manifolds with or without boundary, $A \subseteq M$ an arbitrary subset and $f: A \rightarrow N$ a map. We call f *extension-smooth* if around each $x \in A$ there exist charts

$$h: U \rightarrow M \text{ around } x \\ k: V \rightarrow N \text{ around } f(x),$$

such that

$$g := k^{-1} \circ f \circ h|_{h^{-1}(A)}$$

admits a smooth extension around $h^{-1}(x)$ in the sense that there exist open subsets $U_0 \subseteq \mathbb{R}^n$ and $V_0 \subseteq \mathbb{R}^m$ with $h^{-1}(x) \in U_0$ and $k^{-1}(f(x)) \in V_0$ and a smooth map

$$\tilde{g}: U_0 \rightarrow V_0$$

with $g|_{U_0 \cap h^{-1}(A)} = \tilde{g}|_{U_0 \cap h^{-1}(A)}$.

We now claim the following.

Lemma 12.3. *If $\partial N = \emptyset$ then Lee's definition and Definition 12.2 are equivalent. In general, if f is smooth in the sense of Lee's definition, then it is also extension-smooth but the converse may fail.*

Lemma 12.4. *If A is a submanifold of M , then $f: A \rightarrow N$ is smooth as a map of smooth manifolds if and only if it is extension-smooth.*

In particular the extension Lemma in [Lee11] (Lemma 2.26.) still holds and the proof of the Whitney approximation theorems in chapter 6 work the same (here the extension Lemma is used). But with our new definition Problem 6-7 in [Lee11] now is an actual counterexample. (with Lee's definition F would not be smooth on the closed subset A but only smooth as a map between manifolds).

Another way to properly define smoothness of $f: A \rightarrow N$ would be to change the codomain in Lee's definition to be the double of N . This should be equivalent to Definition 12.2 and is arguably more elegant at the cost of being less elementary.

Smooth Homotopies

This allows us to define what a smooth homotopy is:

Definition 12.5. Let M, N be manifolds with boundary. A homotopy

$$H: M \times [0, 1] \rightarrow N,$$

is called smooth, if H is extension-smooth (see Definition 12.2), where we consider $M \times [0, 1]$ as a subset of the manifold $M \times \mathbb{R}$.

In [Lee11] the author defines a smooth homotopy as a map $H: M \times [0, 1] \rightarrow N$, which can be extended to a smooth map $\bar{H}: M \times \mathbb{R} \rightarrow N$. As we have seen in Counterexample 12.1 this is a stronger requirement compared to Definition 12.5, as less homotopies are considered to be smooth by Lee's definition. However the homotopy classes of smooth maps are the same. More precisely this is formulated in the following lemma.

Lemma 12.6. *Let M, N be manifolds with or without boundary and $f, g: M \rightarrow N$ smooth maps. Then f and g are smoothly homotopic in the sense of Definition 12.5 if and only if there exists a homotopy $H: M \times [0, 1] \rightarrow N$ between f and g , which can be extended to a smooth map $\bar{H}: M \times \mathbb{R} \rightarrow N$.*

Proof. Let $G: M \times [0, 1] \rightarrow N$ be a smooth homotopy from f to g in the sense of Definition 12.5. Choose a smooth function $\phi: \mathbb{R} \rightarrow [0, 1]$ with

$$\begin{aligned}\phi(t) &= 0 \text{ for } t \leq 0 \\ \phi(t) &= 1 \text{ for } t \geq 1\end{aligned}$$

Then

$$\Phi: M \times [0, 1] \rightarrow M \times [0, 1], (x, t) \mapsto (x, \phi(t))$$

is extension-smooth (it can be smoothly extended to $M \times \mathbb{R}$ and this implies that it is smooth in the sense of Definition 12.2, as Lee's definition is stronger by Lemma 12.3). Furthermore, it is easily verified that the composition of extension-smooth maps on arbitray subsets is extension-smooth, such that

$$H := G \circ \Phi: M \times [0, 1] \rightarrow N$$

is extension-smooth. Then H is a homotopy from f to g and we can extend H to the smooth map

$$\bar{H}: M \times \mathbb{R} \rightarrow N, (x, t) \mapsto G(x, \phi(t)).$$

The converse follows directly from Lemma 12.3. □

12.2 Integrating vector valued functions

Let V be a finite dimensional real vector space and $(X, \mathcal{A}, \mu)$ a measure space. Given a function $f: X \rightarrow V$ we would like to define its integral. First of all any finite dimensional vector space carries a canonical topology (the topology induced by any norm), hence we can consider the Borel σ -algebra on V and get a well defined notion of what it means for f to be measurable. The definition of the integral can now be reduced to the case $V = \mathbb{R}^n$:

Definition 12.7. Let $f: (X, \mathcal{A}, \mu) \rightarrow V$ be a measurable function. Choose a linear isomorphism $A: V \rightarrow \mathbb{R}^n$. We call f integrable if $Af: X \rightarrow \mathbb{R}^n$ is integrable and in this case we set

$$\int_X f d\mu := A^{-1} \int_X Af d\mu$$

Linearity of the integral implies that this is well defined, i.e. independent of the choice of A . Moreover, this definition describes how to compute such an integral in practice: First choose any basis of V and express f in that basis. Then integrate each component function. The result gives the coordinates of $\int_X f d\mu$ in the chosen basis.

Linearity of the integral of $\mathbb{R}^n$ -valued functions also generalizes immediately:

Lemma 12.8. *Let $f: (X, \mathcal{A}, \mu) \rightarrow V$ be an integrable function and $T: V \rightarrow V$ linear, then*

$$T \int_X f d\mu = \int_X T f d\mu$$

Proof. Choose an isomorphism $A: V \rightarrow \mathbb{R}^n$ then by definition

$$T \int_X f d\mu = T A^{-1} \int_X Af d\mu = A^{-1} (ATA^{-1}) \int_X Af d\mu = A^{-1} \int_X AT f d\mu$$

□

12.3 Miscellaneous Statements about Manifolds

To show smoothness of a function which is defined implicitly, the following theorem is often useful.

Theorem 12.9 (Implicit function theorem). *Let M, N, S be smooth manifolds (without boundary) with $\dim(N) = \dim(S)$ and $f: M \times N \rightarrow S$ a smooth function. Moreover let $p_0 \in M, q_0 \in N$ and $s_0 = f(p_0, q_0) \in S$. Suppose that the map*

$$N \rightarrow S, q \mapsto f(p_0, q)$$

is regular at q_0 , then around p_0 the equation

$$f(p, q) = s_0$$

can be solved implicitly for q , i.e. there exists an open neighbourhood $U \subseteq M$ of p_0 and a smooth function $g: U \rightarrow N$, such that

$$f(p, g(p)) = s_0$$

for all $p \in U$.

Proof. By the regular value theorem there exists some open neighbourhood $V \subseteq M \times N$ of (p_0, q_0) , such that

$$P := f^{-1}(s_0) \cap V \subseteq M \times N$$

is a submanifold. Let $\pi_M: M \times N \rightarrow M$ be the projection. Suppose we have shown that $\pi_M|_P$ is a local diffeomorphism around (p_0, q_0) , then we can find a local inverse

$$\hat{g}: U \rightarrow P$$

defined on some open neighbourhood $U \subseteq M$ of p_0 . Since $\pi_M \circ \hat{g} = \text{id}_U$ we get that $\hat{g}$ is of the form

$$\hat{g}(p) = (p, g(p))$$

for some smooth map $g: U \rightarrow N$ and thus

$$f(p, g(p)) = s_0.$$

To see that $\pi_M|_P$ is a local diffeomorphism around (p_0, q_0) it suffices to show that

$$d_{(p_0, q_0)}(\pi_M|_P): T_{(p_0, q_0)}P \rightarrow T_{p_0}M$$

is injective since $\dim(P) = \dim(M)$. The kernel of $d_{(p_0, q_0)}\pi_M: T_{p_0}M \oplus T_{q_0}N \rightarrow T_{p_0}M$ is precisely $T_{q_0}N$ and

$$T_{(p_0, q_0)}P = \ker d_{(p_0, q_0)}f,$$

so

$$\ker d_{(p_0, q_0)}(\pi_M|_P) = T_{q_0}N \cap \ker d_{(p_0, q_0)}f.$$

But given $v \in T_{q_0}N \cap \ker d_{(p_0, q_0)}f$ we get that

$$0 = d_{(p_0, q_0)}f(v) = d_{q_0}(q \mapsto f(p_0, q))(v)$$

and since we assumed $q \mapsto f(p_0, q)$ to be regular at q_0 we conclude that $v = 0$. □

13 Solutions to selected exercises

Exercise 13.1 (Own exercise). Let M be the Moebius line bundle. Consider the map $f: S^1 \rightarrow S^1$, $z \mapsto z^2$. Then the pullback bundle $E := f^*M$ is trivial.

Intuitively, we can see this as follows: The fiber over $e^{\pi i}$ of E is just the fiber over $e^{2\pi i}$ of M , which has already made a half-twist, i.e. the fiber of E over 1 can be imagined as a straight line and then begins twisting upon increasing the angle. When the angle $\theta = \pi$ is reached a half twist has been performed. But upon further increasing the angle, another half twist is performed as we "walk" around the moebius bundle a second time. So E is basically a double twisted moebius bundle, which is trivial.

Proof. Proof of Exercise 13.1 We define $M := \mathbb{R} \times \mathbb{R} / \sim$, where the equivalence relation is given by

$$(x, t) \sim (x + k, (-1)^k t)$$

and the projection $p: M \rightarrow S^1$ is defined by $p([(x, t)]) := e^{2\pi i x}$. By identifying S^1 with $[0, 1] / \sim_1$ we can write f as

$$f([\theta]) = e^{2\pi i \theta},$$

such that E is a line bundle over $[0, 1] / \sim$. To see that E is trivial we define a global frame:

$$\sigma: [0, 1] / \sim \rightarrow E, [\theta] \mapsto ([\theta], [(2\theta, 1)])$$

Note that $[(0, 1)] = [(2, 1)]$ in M , such that σ is a well-defined section. Furthermore, it is easily seen that this section is nowhere vanishing. To see that σ is smooth we could write σ in the local trivialisations of E . Alternatively we can argue as follows: Consider

$$s: [0, 1] \rightarrow M, \theta \mapsto [(2\theta, 1)]$$

and let $\pi: [0, 1] \rightarrow [0, 1] / \sim$ be the projection map. Then $s = \sigma \circ \pi$. Clearly s is smooth and it is easily verified that π is a local diffeomorphism, such that smoothness of σ follows. This completes the proof. $\square$

Exercise 13.2 (Exercise 10-14 from [Lee11]). Let M be a smooth manifold and $S \subseteq M$ a submanifold. Then TS is a subbundle of $TM|_S$.

Proof. Let $\sigma_1, \dots, \sigma_k$ be a local frame for TS . Then we can also consider the σ_i as local sections of $TM|_S$ via the inclusion $TS \hookrightarrow TM|_S$. If we can show that these are still smooth sections we are done as this shows that TS (as a set) is a subbundle of $TM|_S$ and the induced smoothness structure is the same as the usual one on TS (see the construction of the smoothness structure on the subbundle in Lemma 10.32 in [Lee11] to see that this indeed yields the same structures). But smoothness of σ_i as local sections of $TM|_S$ follows directly from the fact that the inclusion $TS \hookrightarrow TM|_S$ is just the restriction of the differential of the inclusion $\iota: S \hookrightarrow M$ to the submanifold $TM|_S \subseteq TM$ in the codomain (the fact that $TM|_S$ is a submanifold of TM follows for example from Lemma ??). $\square$

Exercise 13.3 (Mentioned in [Bre13] in chapter 'Conventions for CW complexes'). Let (X, x_0) and (Y, y_0) be compact pointed topological spaces, then the smash product $X \wedge Y$ is homeomorphic to the one point compactification of $Z := (X - x_0) \times (Y - y_0)$. In particular

$$S^n \wedge S^m \cong S^{n+m}.$$

Proof sketch. Let Z^+ be the one point compactification of Z . Define $f: X \times Y \rightarrow Z^+$ by

$$f(x, y) := \begin{cases} (x, y) & , x \neq x_0 \text{ and } y \neq y_0 \\ \infty & , \text{otherwise} \end{cases}$$

It is then straightforward to show that f is continuous and the following diagram is a push-out:

$$\begin{array}{ccc} X \vee Y & \longrightarrow & X \times Y \\ \downarrow & & \downarrow f \\ * & \longrightarrow & Z^+ \end{array}$$

So by the universal property of push-outs we get $Z^+ \cong X \wedge Y$. The statement about the smash product of spheres then just follows from $S^n - * \cong \mathbb{R}^n$ and that the one point compactification of $\mathbb{R}^n$ is S^n . $\square$

14 Alternative Proofs

This is a collection of alternative proofs for various statements from books, lecture notes etc.

Theorem 14.1 (Theorem 10.34 from [Lee11]). *Let E and E' be smooth vector bundles over a manifold M , and let $F: E \rightarrow E'$ be a bundle homomorphism. Then $\ker F$ and $\operatorname{im} F$ are smooth subbundles of E and E' , respectively, if and only if F has constant rank.*

Proof. We present an alternative proof to show that $\ker F$ is a smooth subbundle. Let $\operatorname{rank} F = k$. Choose a local frame

$$\sigma_1, \dots, \sigma_n: U \rightarrow E,$$

such that $F \circ \sigma_1, \dots, F \circ \sigma_k$ are linearly independent sections, i.e. a local frame for $\operatorname{im} F$. So for $j > k$ we can write

$$F \circ \sigma_j = \sum_{i=1}^k \tau_{ij} F \circ \sigma_i$$

for smooth functions

$$\tau_{ij}: U \rightarrow \mathbb{R}$$

Define

$$\sigma'_j := \sigma_j - \sum_{i=1}^k \tau_{ij} \sigma_i$$

Then $F \circ \sigma'_j = 0$, thus $\operatorname{im} \sigma'_j \subseteq \ker F$. Furthermore, $\sigma'_{k+1}, \dots, \sigma'_n$ are linearly independent, since $\sigma_1, \dots, \sigma_n$ is a local frame. This shows the statement. $\square$

Proposition 14.2 (Proposition 5.17 from [Lee18]). *The musical isomorphisms commute with the (total) covariant derivative: if F is any smooth tensor field with a contravariant i -th index position and $\flat$ represents the operation of lowering the i -th index, then*

$$\nabla_X(F^\flat) = (\nabla_X F)^\flat \quad (23)$$

for all vector fields X . In particular

$$\nabla(F^\flat) = (\nabla F)^\flat. \quad (24)$$

Similarly, if G has covariant i -th position and $\sharp$ denotes raising the i -th index, then

$$\nabla_X(F^\sharp) = (\nabla_X F)^\sharp, \quad \nabla(F^\sharp) = (\nabla F)^\sharp. \quad (25)$$

In the book [Lee18], the author uses the identity $F^\flat = \operatorname{tr}(F \otimes g)$, which is however only correct in the case where F is a vector field, since then

$$F^\flat = \langle F, - \rangle = \operatorname{tr}(F \otimes g)$$

by definition of the trace operator. In general this argument fails, as can be seen by writing each side in a local frame. Thus we present an alternative proof.

Proof. First let $F = Z$ be a vector field. Instead of using the identity mentioned above we can also argue as follows: Write $\alpha := \langle Z, - \rangle$. Then for an arbitrary vector field Y we compute:

$$(\nabla_X \alpha)(Y) = X \cdot \alpha(Y) - \alpha(\nabla_X Y) = \langle \nabla_X Z, Y \rangle + \langle Z, \nabla_X Y \rangle - \langle Z, \nabla_X Y \rangle = \langle \nabla_X Z, Y \rangle$$

hence

$$\nabla_X(Z^\flat) = \nabla_X \langle Z, - \rangle = \langle \nabla_X Z, - \rangle = (\nabla_X Z)^\flat.$$

For general F we may assume (by linearity of the $\flat$ operator)

$$F = Z_1 \otimes \dots \otimes Z_k \otimes \varepsilon^1 \otimes \dots \otimes \varepsilon^l$$

for some vector fields Z_j and covector fields ε^j . But then

$$F^\flat = Z_1 \otimes \dots \otimes \langle Z_i, - \rangle \otimes \dots \otimes Z_k \otimes \varepsilon^1 \otimes \dots \otimes \varepsilon^l$$

so (23) follows from the product rule and what has been shown above. Then (24) follows from (23) and the definition of the $\flat$ operator as well as the total covariant derivative. For (25) we argue as in [Lee18]: note that $\sharp$ and $\flat$ are inverses to each other (since they operate on the same index position), so by substituting $F = G^\sharp$ in (23) and (24) and applying $\sharp$ to both sides, the statement follows. $\square$

References

- [Ada93] Masahisa Adachi. *Embeddings and immersions*, volume 124. American Mathematical Soc., 1993.
- [Bre13] Glen E Bredon. *Topology and geometry*. Springer Science & Business Media, 2013.
- [Ebe12] Johannes Ebert. A lecture course on cobordism theory, 2012. Lecture notes.
- [GG12] Martin Golubitsky and Victor Guillemin. *Stable mappings and their singularities*, volume 14. Springer Science & Business Media, 2012.
- [GP74] Victor Guillemin and Alan Pollack. *Differential Topology*. 1974.
- [Hir12] Morris W Hirsch. *Differential topology*, volume 33. Springer Science & Business Media, 2012.
- [Kon26] Emilia Konrad. Globale analysis, 2026. Lecture notes for course "Globale Analysis" in WS 2025/26.
- [Lee11] John M. Lee. *Introduction to Topological Manifolds*, volume 202. Springer, New York, 2011.
- [Lee18] John M. Lee. *Introduction to Riemannian manifolds*, volume 2. Springer, 2018.
- [Mil63] John W. Milnor. *Morse Theory*. Princeton University Press, 1963.
- [Mil65] John Milnor. *Lectures on the h-cobordism Theorem*. University Press of Virginia, 1965.
- [Mit11] Stephen A. Mitchell. Notes on principal bundles and classifying spaces. 2011.
- [Rah17] Adel Rahman. Chern-weil theory. 2017.
- [Sch12] Bernd Schmidt. Analysis 3, 2012. Lecture notes for course "Analysis 3" in WS 2011/12.
- [Ste23a] Wolfgang Steimle. Einführung in die geometrie, 2023. Lecture notes for course "Einführung in die Geometrie".
- [Ste23b] Wolfgang Steimle. Topologie, 2023. Lecture notes for course "Einführung in die Topologie", LaTeX version.
- [Ste25] Wolfgang Steimle. Algebraische topologie, 2025. Lecture notes for course "Algebraische Topologie", LaTeX version.
- [tD08] Tammo tom Dieck. *Algebraic topology*. 2008.
- [Vin24] Johannes Vinnemeier. Homology and perfect morse functions for manifolds with boundary and their gluings. 2024. Bachelor Thesis.
- [Wer18] Dirk Werner. *Funktionalanalysis*. Springer, 8. edition, 2018.