

Contents

1	On manifolds with boundary	2
1.1	Regular values and transversality	2
1.2	Morse Theory	4
1.3	Integral curves and flows	7
2	Miscellaneous	12
2.1	Smoothness on arbitrary subsets	12
2.2	Fiber Bundles	14
2.2.1	Smooth bundles	20
2.2.2	Whitney Sums	21
	References	22

This is a collection of statements I have proven myself and considered to be useful in some cases. I will often only present proof sketches instead of fully detailed proofs. Furthermore these notes contain a variety of counterexamples.

1 On manifolds with boundary

In this section we discuss how known theorems/lemmas fail for manifolds with non-empty boundary and how they can be generalized.

1.1 Regular values and transversality

It is known that for the regular value theorem, and more generally statements about transversality, to still hold for manifolds with boundary one needs to impose additional conditions on the boundary behaviour of a smooth function $f: M \rightarrow N$. Here is another lemma, that fits into the same category. We call $q \in N$ a **boundary value** of f if $q \in f(\partial M)$.

Lemma 1.1. *Let M be a manifold with boundary and $f: M \rightarrow \mathbb{R}$ a smooth function. Furthermore, let $a \in \mathbb{R}$ be a regular value of f , that is not a boundary value. Then $M_a := f^{-1}(-\infty, a]$ is a submanifold with boundary $\partial M_a = f^{-1}(a)$.*

Proof sketch. For $p \in f^{-1}(a)$ we have $p \in \text{int}(M)$ by assumption. Since $\text{int}(M)$ is a boundaryless manifold the local submersion theorem holds, i.e. f looks like the orthogonal projection $\mathbb{R}^n \rightarrow \mathbb{R}$ around p . Then the proof goes the same as in the boundaryless case. $\square$

Counterexample 1.2. To see that Lemma 1.1 fails without the assumption that a is not a boundary value consider $M := S^1 \times [0, 1]$ and $f: M \rightarrow \mathbb{R}, (z, t) \mapsto t$. Then f is everywhere regular but $f^{-1}(-\infty, 0]$ does not satisfy the statements in Lemma 1.1.

Let M, N be manifolds with boundary and $f: M \rightarrow N$ smooth. We define the **boundary map** ∂f of f by

$$\partial f := f|_{\partial M}.$$

We now generalize two theorems from [GP74] about transversal maps by allowing the codomain to have boundary as well.

Theorem 1.3 (Transversality Theorem). *Let M, N be manifolds with boundary, $f: M \rightarrow N$ smooth and $X \subseteq N$ a submanifold without boundary. If*

$$f \pitchfork X \text{ and } \partial f \pitchfork X$$

then $X' := f^{-1}(X)$ is a submanifold with boundary

$$\partial X' = X' \cap \partial M$$

and

$$\text{codim}(X') = \text{codim}(X).$$

Proof. The case $\partial N = \emptyset$ is shown in [GP74] (p. 60). If $\partial N \neq \emptyset$ consider the inclusion

$$\iota: N \rightarrow D(N),$$

where $D(N)$ denotes the double of N and define $F := \iota \circ f$. Then $Y := \iota(X)$ is a submanifold of $D(N)$ and $X' = F^{-1}(Y)$. Furthermore, since ι is an embedding (and $\dim(N) = \dim(D(N))$) it follows that

$$F \pitchfork Y \text{ and } \partial F \pitchfork Y,$$

such that the statement follows from the boundaryless case. $\square$

Theorem 1.4 (Parametric Transversality Theorem). *Let M, N be manifolds with boundary and $X \subseteq N$ a submanifold without boundary. Furthermore, let S be a manifold without boundary and $F: M \times S \rightarrow N$ a smooth map with*

$$F \pitchfork X \text{ and } \partial F \pitchfork X.$$

Then for almost every $s \in S$ the map $f_s: M \rightarrow N$ defined by $f_s(p) := F(p, s)$ satisfies

$$f_s \pitchfork X \text{ and } \partial f_s \pitchfork X.$$

Proof. By Theorem 1.3 we know that

$$X' := F^{-1}(X)$$

is a submanifold (with boundary) of $M \times S$. Consider the projection $\pi: M \times S \rightarrow S$. We can then show, that for a regular value $s \in S$ of the map $\pi|_{X'}$ the maps f_s and ∂f_s are transversal to X , which is just some linear algebra. The proof is exactly the same as in the case $\partial N = \emptyset$, see [GP74] (p. 68). Then the statement follows from Sard's Theorem. $\square$

Theorem 1.5 (Transversality Homotopy Theorem). *Let M, N be manifolds with boundary, $f: M \rightarrow N$ smooth and $X \subseteq N$ a submanifold without boundary. Then f is smoothly homotopic to a map $g: M \rightarrow N$ with*

$$g \pitchfork X \text{ and } \partial g \pitchfork X.$$

Proof. First we may assume that $\text{im}(f) \subseteq \text{int}(N)$, since f is always smoothly homotopic to such a map (the inclusion $\text{int}(N) \hookrightarrow N$ is a smooth homotopy equivalence). Define

$$Y := X \cap \text{int}(N)$$

and

$$g: M \rightarrow \text{int}(N), p \mapsto f(p).$$

Using the Transversality Homotopy Theorem for the case $\partial N = \emptyset$ (see [GP74] p. 70) we get that g is smoothly homotopic to a map $g': M \rightarrow \text{int}(N)$, such that

$$g' \pitchfork Y \text{ and } \partial g' \pitchfork Y.$$

Now define $f' := \iota \circ g$, where $\iota: \text{int}(N) \hookrightarrow N$ is the inclusion. Since f' does not intersect X at boundary points, i.e. $\text{im}(f') \cap X \subseteq \text{int}(N)$, the transversality properties of g' directly imply

$$f' \pitchfork X \text{ and } \partial f' \pitchfork X.$$

□

1.2 Morse Theory

For a smooth function $f: M \rightarrow \mathbb{R}$ and a regular value $a \in \mathbb{R}$ that is not a boundary value we define the **sublevel set** of a to be

$$M_a := f^{-1}(-\infty, a].$$

Lemma 1.1 guarantees, that M_a is a (sub-)manifold with boundary.

Lemma 1.6. *Let M be a compact manifold with boundary and $f: M \rightarrow \mathbb{R}$ a Morse function. Furthermore let $a, b \in \mathbb{R}$ be regular values of f , such that $[a, b]$ does not contain any critical values nor any boundary values. Then*

$$M_a \cong M_b.$$

Proof. Let $K := f^{-1}[a, b]$ and let $U := \{p \in \text{int}(M) \mid p \text{ is a regular point of } f\}$. Then $U \subseteq M$ is open and $K \subseteq U$. Choose a cutoff function $\phi: M \rightarrow \mathbb{R}$ with

- $\phi|_K = 1$
- $\text{supp}(\phi) \subseteq U$
- $0 \leq \phi \leq 1$

Now choose a pseudo gradient¹ X of f and define the smooth vector field

$$Y := \phi \frac{-1}{X \cdot f} X,$$

which we define to be 0 outside of U . Since Y vanishes in a neighbourhood around ∂M (or more generally since Y is tangent to ∂M , see [Lee11] Theorem 9.34) we can consider the flow of Y :

$$\theta: M \times \mathbb{R} \rightarrow M.$$

Since M is compact (and Y tangent to ∂M) each integral curve is defined for all time. Let $\theta_t := \theta(-, t): M \rightarrow M$. We claim that

$$\theta_{b-a}(M_b) \subseteq M_a, \tag{1}$$

$$\theta_{a-b}(M_a) \subseteq M_b. \tag{2}$$

¹Here we agree to the convention that $X \cdot f < 0$ outside of critical points.

Let $p \in f^{-1}[a, b]$ and let $\gamma_p: \mathbb{R} \rightarrow M$ be the integral curve of Y starting at p . Consider

$$h: \mathbb{R} \rightarrow \mathbb{R}, t \mapsto f(\gamma_p(t)).$$

Then for $t \in \mathbb{R}$ with $h(t) \in [a, b]$ we have $\phi(\gamma_p(t)) = 1$ and thus

$$h'(t) = \dot{\gamma}_p(t) \cdot f = Y_{\gamma_p(t)} \cdot f = -1.$$

So h is a smooth function with $c := h(0) = f(p) \in [a, b]$ and $h'(t) = -1$ as long as $h(t) \in [a, b]$. We claim that $h(c - a) = a$. To see this let

$$s := \sup\{t \in [0, c - a] \mid h([0, t]) \subseteq [a, b]\}.$$

Assume $s < c - a$. For continuity reasons we get that $h([0, s]) \subseteq [a, b]$ and thus $h'(t) = -1$ for $t \in [0, s]$. Then $h(t) = c - t$ for $t \in [0, s]$. But then $h(s) \in (a, b]$ and h is strictly decreasing around $t = s$ such that there exists $\epsilon > 0$ with $h([0, s + \epsilon]) \subseteq [a, b]$, contradicting the definition of s . So $s = c - a$ and we get that $h(t) = c - t$ for $t \in [0, c - a]$. Thus $h(c - a) = a$.

As $h'(t) \leq 0$ for all $t \in \mathbb{R}$ we conclude

$$f(\theta_{b-a}(p)) = h(b - a) \leq h(c - a) = a,$$

thus $\theta_{b-a}(p) \in M_a$. For $p \in M_b \setminus f^{-1}[a, b] = \text{int} M_a$ we also have

$$\theta_{b-a}(p) \in M_a$$

as f decreases along gradient flow lines of Y . This shows (1). We argue similiarly for (2). Let $p \in M_a$ and let $\gamma_p: \mathbb{R} \rightarrow M$ be the integral curve of Y starting at p . Consider

$$h: \mathbb{R} \rightarrow \mathbb{R}, t \mapsto f(\gamma_p(-t)).$$

Then $h'(t) = \phi(\gamma_p(-t)) \leq 1$. Assume that $h(b - a) = f(\theta_{a-b}(p)) > b$. Then by the mean value theorem there exists some $\xi \in [0, b - a]$ with

$$h'(\xi) = \frac{h(b - a) - h(0)}{b - a} \geq \frac{h(b - a) - a}{b - a} > 1,$$

which is a contradiction. So $f(\theta_{a-b}(p)) \leq b$, showing (2). Thus θ_{b-a} restricts to a diffeomorphism $M_b \xrightarrow{\cong} M_a$ with inverse θ_{a-b} . $\square$

Counterexample 1.7. Example 1.2 also shows, that we need to impose conditions on the boundary values in Lemma 1.6. To see that we also need that condition for Lemma 1.9 we may similiarly define f as the height function on the disjoint union of two cylinders which are slightly shifted in height. Similiar examples show, that Theorem 1.10 fails without the assumption on boundary values.

Remark 1.8. We can weaken the assumptions of Lemma 1.6. First of all f does not need to be a Morse function. We can then still construct a vector field X , such that $X \cdot f < 0$ outside of critical points. Secondly, we may replace the

assumption that M is compact by instead requiring $f^{-1}[a, b]$ to be compact, which is for example the case if f is proper. We then only have to modify the proof in the definition of U . Instead we define U to be a *compact* neighbourhood of $K := f^{-1}[a, b]$, i.e. $K \subseteq \text{int}U$, which does not contain any critical points nor boundary points. For example we may define U as a union of finitely many (here we need compactness of $f^{-1}[a, b]$) compact sets K_i not containing critical points nor boundary points such that $\text{int}K_i$ still cover K .

Similiarly we may proof the following, for which we can also weaken the assumptions as described in the remark above.

Lemma 1.9. *Let M be a compact manifold with boundary and $f: M \rightarrow \mathbb{R}$ a Morse function. Furthermore let $a, b \in \mathbb{R}$ be regular values of f , such that $[a, b]$ does not contain any critical values nor any boundary values. Then M_a is a (not necessarily smooth) deformation retract of M_b .*

Proof. We consider the flow θ as in Lemma 1.6 and define a retraction

$$r: M_b \rightarrow M_a, p \mapsto \begin{cases} \theta(f(p) - a, p), & f(p) \geq a \\ p, & f(p) \leq a \end{cases}$$

As we have shown above $\theta(f(p) - a, p) \in f^{-1}(a)$. It follows that r is continous and well-defined. To see that r is a deformation retract consider

$$H: M_b \times [0, 1] \rightarrow M_b, (p, t) \mapsto \begin{cases} \theta(t(f(p) - a), p), & f(p) \geq a \\ p, & f(p) \leq a \end{cases}$$

Then $H(-, 0) = \text{id}_{M_b}$ and $H(-, 1) = r$. □

We would now also like to generalize the result, that the topology of M changes by attaching a λ -cell when crossing a critical point c_0 of index λ to compact manifolds with boundary. This statement can now be shown using the two lemmas above by requiring $f^{-1}[a, b]$ to only contain the critical point c_0 and no boundary points. The proof now works exactly the same as for the boundaryless case. See [Mil63]. Let us state the theorem for manifolds with boundary:

Theorem 1.10. *Let M be a compact manifold with boundary and $f: M \rightarrow \mathbb{R}$ a Morse function. Let $c_0 \in M$ be a critical point of f of index λ and $c := f(c_0)$ as well as $\epsilon > 0$, such that $f^{-1}[c - \epsilon, c + \epsilon]$ contains no boundary points and no critical points besides c . Then for ϵ sufficiently small $M_{c+\epsilon}$ has the homotopy type of $M_{c-\epsilon}$ with a λ -cell attached. More precisely, there exists an attaching map*

$$\phi: S^{\lambda-1} = \partial D^\lambda \rightarrow \partial M_{c-\epsilon}$$

and a homotopy equivalence

$$r: M_{c+\epsilon} \rightarrow M_{c-\epsilon} \cup_\phi D^\lambda$$

with homotopy inverse

$$\iota: M_{c-\epsilon} \cup_{\phi} D^{\lambda} \rightarrow M_{c+\epsilon},$$

such that $r \circ \iota = id_{M_{c-\epsilon} \cup_{\phi} D^{\lambda}}$.

Proof Sketch. Following [Mil63] we first construct $F: M \rightarrow \mathbb{R}$, such that

$$F^{-1}(-\infty, c + \epsilon] = f^{-1}(-\infty, c + \epsilon] = M_{c+\epsilon}.$$

Furthermore $F^{-1}[c - \epsilon, c + \epsilon]$ does not contain any critical points or boundary points, such that $F^{-1}(-\infty, c - \epsilon]$ is a deformation retract of $F^{-1}(-\infty, c + \epsilon]$ by Lemma 1.9. We then show that $M_{c-\epsilon} \cup_{\phi} D^{\lambda}$ is a deformation retract of $F^{-1}(-\infty, c - \epsilon]$ (or more precisely the image of $M_{c-\epsilon} \cup_{\phi} D^{\lambda}$ under an embedding is a deformation retract). This then shows that $M_{c-\epsilon} \cup_{\phi} D^{\lambda}$ is a deformation retract of $M_{c+\epsilon}$. $\square$

However this does not yield a cell decomposition of M as we do not know how the topology changes when crossing boundary points. Instead we can use the fact that any compact without boundary admits a cell-decomposition directly, to proof the case when M has non-empty boundary.

Theorem 1.11 (Cell-decompositions of Manifolds). *Let M be compact manifold with or without boundary. Then M has the homotopy type of a CW-complex.*

Proof. The case when M has no boundary is shown in [Mil63]. If $\partial M \neq \emptyset$ we can argue as follows: consider the double $D(M)$ of M . Then $D(M)$ is a compact manifold without boundary. Now we construct a Morse function on $D(M)$. Choose a Morse function $f: M \rightarrow \mathbb{R}$ with $f \leq 0$ and $f(p) = 0$ if and only if $p \in \partial M$, such that f has no critical points on the boundary. Such Morse functions always exist, see [Vin24]. We can then glue f and $-f$ together to get a Morse function F on $D(M)$, again see [Vin24]. Then $M \cong F^{-1}(-\infty, 0]$ and 0 is a regular value of F , allowing us to refer back to the boundaryless case, which tells us that every sublevel set M_a for a regular value $a \in \mathbb{R}$ has the homotopy type of a CW-complex. $\square$

Remark 1.12. We may also formulate Theorem 1.10 and 1.11 for handle bodies instead of CW-complexes. The proof is essentially the same but instead of construction a deformation retract $F^{-1}(-\infty, c - \epsilon] \rightarrow M_{c-\epsilon} \cup_{\phi} D^{\lambda}$ one constructs a homeomorphism $F^{-1}(-\infty, c - \epsilon] \rightarrow M_{c-\epsilon} \cup_{\phi} D^{\lambda} \times D^{n-\lambda}$.

1.3 Integral curves and flows

For manifolds with boundary we essentially define integral curves in the same way as for boundaryless manifolds, but allow the domain to be any non-trivial interval (i.e. not a single point) containing 0. In particular this interval may only be half-open or closed. In general, the existence of integral curves on manifolds with boundary fails as illustrated by the following example:

Counterexample 1.13. Consider the lower halfspace $M := \mathbb{R}_{y \leq 0}^2$ and the vector field $X := \frac{\partial}{\partial x} + 2x \frac{\partial}{\partial y}$. Then for $p = (0, 0)$ the integral curve is only defined for $t = 0$, since on $\mathbb{R}^2$ it would be given by $\gamma(t) = (t, t^2)$ but $t^2 > 0$ for $t \neq 0$.




Figure 1: The vector field X . The integral curves are parabolas.

We do however still get a uniqueness result *if* the integral curve exists. Furthermore, if the vector field is sufficiently well-behaved we can guarantee the existence.

Lemma 1.14 (Uniqueness and existence of Integral curves). *Let M be a manifold with boundary, X a smooth vector field on M and $p \in M$. If there exists an integral curve starting at p , then there exists a unique maximal integral curve $\gamma: I \rightarrow M$ starting at p , where one of the four following cases holds for $a \leq 0 \leq b$, $a < b$:*

- (i) $I = (a, b)$
- (ii) $I = [a, b)$ and $\gamma(a) \in \partial M$
- (iii) $I = (a, b]$ and $\gamma(b) \in \partial M$
- (iv) $I = [a, b]$ and $\gamma(a), \gamma(b) \in \partial M$

Furthermore, if X is either transversal to the boundary or everywhere tangent, the unique maximal integral curve starting at p always exists.

Proof sketch of Lemma 1.14. Let's begin by showing uniqueness. The proof is essentially the same as in the boundaryless case (see for example [Lee11]): We show that two integral curves $\gamma: I \rightarrow M$ and $\gamma': I' \rightarrow M$ starting at p agree on their common domain. This then allows us to define the maximal integral curve on some interval I and every interval containing 0 is of the form (i)–(iv)². The only difference is that I and I' may not be open intervals. So we have to show that uniqueness still applies. We claim: Let $U \subseteq \mathbb{R}^n$ be open, $F: U \rightarrow \mathbb{R}^n$ a smooth vector field, I an interval of the form (i)–(iv) and $\gamma, \gamma': I \rightarrow U$ integral curves both starting at $p \in U$. Then $\gamma = \gamma'$. If 0 is an interior point of I we can apply the usual uniqueness theorem for γ, γ' restricted to $\text{int}(I)$. By continuity they must then also agree at the endpoint. So it remains to show uniqueness if 0 is a boundary point of I (say for example $I = [0, b)$). But by the existence

theorem for integral curves we can find an integral curve $\eta: (-\epsilon, \epsilon) \rightarrow U$ starting at p and define

$$\beta: (-\epsilon, b) \rightarrow U, t \mapsto \begin{cases} \eta(t), t \leq 0 \\ \gamma(t), t \geq 0 \end{cases}$$

Since $\eta(0) = \gamma(0)$ and $\dot{\eta}(0) = F(p) = \dot{\gamma}(0)$ we get that β is at least C^1 . Similarly we define

$$\beta': (-\epsilon, b) \rightarrow U, t \mapsto \begin{cases} \eta(t), t \leq 0 \\ \gamma'(t), t \geq 0 \end{cases}$$

By the uniqueness theorem we have

$$\beta = \beta'$$

and our claim follows. The case $I = (a, 0]$ goes analogous and for $I = [0, b]$ or $I = [a, 0]$ it is either trivial (if $a = b = 0$) or we consider $I \setminus \{b\}$ resp. $I \setminus \{a\}$ and apply the above argument. Equality at $t = a, t = b$ respectively then again follows by continuity.

To see that $\gamma(t) \in \partial M$ for $t \in \{a, b\}$ a boundary point of I we note that for $p = \gamma(t) \in \text{int}(M)$ we can define the integral curve starting at p on some open interval $(-\epsilon, \epsilon)$ by the existence theorem for integral curves (Picard-Lindelöf). But then t is not a boundary point of I since I is maximal.

Next we show the existence if X is transversal to ∂M . For $p \in \text{int}(M)$ this follows directly from the Picard-Lindelöf Theorem as in the boundaryless case. If $p \in \partial M$, we may argue similarly: Consider a chart $h: \mathbb{R}_{x_n \geq 0}^n \supseteq U \rightarrow M$ around $p = h(0)$ and let Y be the vector field X in this chart, i.e. $D_q h(Y_q) = X_{h(q)}$. Then by definition of smoothness we can extend Y to a smooth vector field Y' on some open neighbourhood $U' \subseteq \mathbb{R}^n$ around $0 = h^{-1}(p)$. Applying Picard-Lindelöf we get the existence of an integral curve $\gamma: (-\epsilon, \epsilon) \rightarrow U'$ of Y' starting at 0. Write

$$Y' = \sum_i F_i \frac{\partial}{\partial x_i}$$

and

$$\gamma = (\gamma_1, \dots, \gamma_n).$$

Assume that X is inwards pointing along the boundary. After restricting to a smaller domain we may assume $F_n \geq 0$. Then

$$\gamma'_n(t) = F_n(\gamma(t)) \geq 0,$$

such that γ_n is increasing. In particular $\gamma_n(t) \geq 0$ for $t \in [0, \epsilon)$, hence $\gamma(t) \in U$ for $t \in [0, \epsilon)$. Then

$$\eta := h \circ \gamma: [0, \epsilon) \rightarrow M$$

is the desired integral curve.

For the case that X is everywhere tangent to ∂M see [Lee11] (Theorem 9.34).

□

On compact manifolds we can say more about the maximal Interval I :

Lemma 1.15 (Existence of Integral curves on compact manifolds). *If M is compact in Lemma 1.14 then the maximal interval I has one of the following forms:*

- (i) $I = (-\infty, \infty)$
- (ii) $I = [a, \infty)$ and $\gamma(a) \in \partial M$
- (iii) $I = (-\infty, b]$ and $\gamma(b) \in \partial M$
- (iv) $I = [a, b]$ and $\gamma(a), \gamma(b) \in \partial M$

Proof. Let $\gamma: I \rightarrow M$ be the unique maximal integral curve guaranteed by Lemma 1.14. We will show that any open boundary point of I must be $\pm\infty$. For simplicity let us only consider $I = [a, b)$. The other cases are completely analogous. By compactness we can find a sequence $0 < t_n \nearrow b$ such that $\gamma(t_n)$ converges to a point $q \in M$.

Case 1: $q \in \text{int}(M)$. By Picard-Lindelöf we can find an open neighbourhood $U \subseteq M$ around q and $\epsilon > 0$, such that for every point $x \in U$ the integral curve starting at x is defined for $t \in (-\epsilon, \epsilon)$. For some fixed m sufficiently large we have $q_0 := \gamma(t_m) \in U$. Assume that $b - t_m < \frac{\epsilon}{2}$. But then we can extend γ to $[a, b + \frac{\epsilon}{2})$ as follows. Consider

$$\gamma': [a - t_m, b - t_m) \rightarrow M, \quad t \mapsto \gamma(t + t_m).$$

Then γ' is a integral curve starting at q_0 . Let $\alpha: (-\epsilon, \epsilon) \rightarrow M$ be another integral curve starting at q_0 . Then α and γ' agree on their common domain by the arguments in Lemma 1.14 and thus we can put them together to define an integral curve

$$\beta: J \rightarrow M$$

for $J := [a - t_m, b - t_m) \cup (-\epsilon, \epsilon) \supseteq [a - t_m, \epsilon)$. But then we can define

$$\gamma'': [a, \epsilon + t_m) \rightarrow M, \quad t \mapsto \beta(t - t_m),$$

which is an integral curve starting at $\gamma(0) = p$. However

$$b + \frac{\epsilon}{2} < t_m + \epsilon$$

and so $[a, b + \frac{\epsilon}{2}) \subseteq [a, \epsilon + t_m)$, contradicting maximality of γ .

²If we put together integral curves defined on open intervals that agree on their common domain, it is clear that this results in a smooth curve, since around each t this curve is given by one of the smooth curves which we put together. If we put together integral curves γ_1, γ_2 defined on something like $I_1 = (a, b]$ and $I_2 = [b, c)$ we have to be more careful around the boundary point b . But the same argument as we applied to β in the proof of Lemma 1.14 shows that putting γ_1 and γ_2 together results in a C^1 -curve. We can then apply the usual uniqueness theorem to see that this curve is already given by a C^∞ -curve.

Case 2: $q \in \partial M$. We argue similarly as above. In a local chart around q we can again apply Picard-Lindelöf (smoothness of X guarantees by definition the existence of a smooth extension of X to an open neighbourhood in $\mathbb{R}^n$). We then again take $q_0 := \gamma(t_m)$ sufficiently close to q , choose an integral curve α starting at q , which we can use to define an integral curve α' starting at q_0 defined on some open interval $(-\epsilon, \epsilon)$ by translation in time as above. Also we define γ' as above. The same argument as above then shows that we can extend γ' (in the local chart) to an integral curve β on $J := [a - t_m, b - t_m] \cup (-\epsilon, \epsilon) \supseteq [a - t_m, b - t_m]$. In particular we have $\beta([a - t_m, b - t_m]) \subseteq M$. Translating in time again yields an integral curve $\gamma'' : [a, b] \rightarrow M$ contradicting maximality. $\square$

Remark 1.16 (A note on flows for manifolds with boundary). In general the concept of a flow is not well behaved if M has boundary, as integral curves may not even exist (see Counterexample 1.13). In [Lee11] the author shows, that if X is tangent to ∂M we still get a well-behaved flow, i.e. it is defined on some open subset $\mathcal{O} \subseteq M \times \mathbb{R}$. I think that if we instead assume, that X is transversal to ∂M , we still get a "good" flow domain in the sense that we can define the flow on some subset $\mathcal{D} \subseteq M \times \mathbb{R}$, which is a codimension 0 submanifold with boundary and $I_p := \{t \in \mathbb{R} \mid (p, t) \in \mathcal{D}\}$ is an interval around 0 for each $p \in M$.

The statement in the above remark should follow from the following lemma, which allows us to define submanifold charts for $\mathcal{D}$.

Lemma 1.17. *Let M be a manifold with boundary and X a vector field of M , which is inward pointing along ∂M . Let*

$$M_\partial := \{p \in M \mid \text{im} \gamma_p \cap \partial M \neq \emptyset\}$$

be all points in M whose trajectory contains boundary points (this is at most one for each $p \in M$). For $p \in M_\partial$ the integral curve γ_p starting at p is defined on some interval $[-\tau(p), a)$ for some $\tau(p) \geq 0$ and $a > -\tau(p)$. Then the following statements hold true

- (i) M_∂ is an open subset of M
- (ii) $\tau : M_\partial \rightarrow \mathbb{R}_{\geq 0}, p \mapsto \tau(p)$ is smooth

Proof sketch. The idea is to consider the maximal flow

$$\theta : \mathcal{D} \rightarrow M, (p, t) \mapsto \gamma_p(t)$$

on some open subset $\mathcal{D} \subseteq \partial M \times [0, \infty)$ and show that this is an embedding. Then

$$M_\partial = \text{im} \theta$$

is open and

$$\tau = \pi \circ \theta^{-1} : M_\partial \rightarrow [0, \infty)$$

for $\pi : \partial M \times [0, \infty) \rightarrow [0, \infty)$ is the canonical projection map. $\square$

2 Miscellaneous

2.1 Smoothness on arbitrary subsets

Let M, N be manifolds and $A \subseteq M$ an arbitrary subset. In [Lee11] the author defines what it means for a map

$$f: A \rightarrow N$$

to be smooth, namely

We call f smooth if around each $x \in A$ there exists a smooth extension $\tilde{f}: U \rightarrow N$ of f to some open neighbourhood $U \subseteq M$ of x (see [Lee11, p. 45]).

However this definition has a fatal flaw: if A happens to be a submanifold of M , a map $f: A \rightarrow N$ might be smooth in the usual sense (i.e. as a map between smooth manifolds) but not smooth in the sense of Lee's definition. Consider for example

Counterexample 2.1.

$$M := \mathbb{R}, N := [0, 1], A := [0, 1], \\ f: A \rightarrow N, t \mapsto t.$$

Then clearly f is smooth as a map between smooth manifolds, but around $0, 1 \in A$ it does not admit a smooth extension onto an open neighbourhood $U \subseteq M$, since the image of such an extension would fail to be contained within N .

We can however correct this flaw by defining:

Definition 2.2 (Extension-smooth). Let M^n, N^m be manifolds with or without boundary, $A \subseteq M$ an arbitrary subset and $f: A \rightarrow N$ a map. We call f *extension-smooth* if around each $x \in A$ there exist charts

$$h: U \rightarrow M \text{ around } x \\ k: V \rightarrow N \text{ around } f(x),$$

such that

$$g := k^{-1} \circ f \circ h|_{h^{-1}(A)}$$

admits a smooth extension around $h^{-1}(x)$ in the sense that there exist open subsets $U_0 \subseteq \mathbb{R}^n$ and $V_0 \subseteq \mathbb{R}^m$ with $h^{-1}(x) \in U_0$ and $k^{-1}(f(x)) \in V_0$ and a smooth map

$$\tilde{g}: U_0 \rightarrow V_0$$

with $g|_{U_0 \cap h^{-1}(A)} = \tilde{g}|_{U_0 \cap h^{-1}(A)}$.

We now claim the following.

Lemma 2.3. *If $\partial N = \emptyset$ then Lee's definition and Definition 2.2 are equivalent. In general, if f is smooth in the sense of Lee's definition, then it is also extension-smooth but the converse may fail.*

Lemma 2.4. *If A is a submanifold of M , then $f: A \rightarrow N$ is smooth as a map of smooth manifolds if and only if it is extension-smooth.*

In particular the extension Lemma in [Lee11] (Lemma 2.26.) still holds and the proof of the Whitney approximation theorems in chapter 6 work the same (here the extension Lemma is used). But with our new definition Problem 6-7 in [Lee11] now is an actual counterexample. (with Lee's definition F would not be smooth on the closed subset A but only smooth as a map between manifolds).

Another way to properly define smoothness of $f: A \rightarrow N$ would be to change the codomain in Lee's definition to be the double of N . This should be equivalent to Definition 2.2 and is arguably more elegant at the cost of being less elementary.

Smooth Homotopies

This allows to define what a smooth homotopy is:

Definition 2.5. Let M, N be manifolds with boundary. A homotopy

$$H: M \times [0, 1] \rightarrow N,$$

is called smooth, if H is extension-smooth (see Definition 2.2), where we consider $M \times [0, 1]$ as a subset of the manifold $M \times \mathbb{R}$.

In [Lee11] the author defines a smooth homotopy as a map $H: M \times [0, 1] \rightarrow N$, which can be extended to a smooth map $\bar{H}: M \times \mathbb{R} \rightarrow N$. As we have seen in Counterexample 2.1 this is a stronger requirement compared to Definition 2.5, as less homotopies are considered to be smooth by Lee's definition. However the homotopy classes of smooth maps are the same. More precisely this is formulated in the following lemma.

Lemma 2.6. *Let M, N be manifolds with or without boundary and $f, g: M \rightarrow N$ smooth maps. Then f and g are smoothly homotopic in the sense of Definition 2.5 if and only if there exists a homotopy $H: M \times [0, 1] \rightarrow N$ between f and g , which can be extended to a smooth map $\bar{H}: M \times \mathbb{R} \rightarrow N$.*

Proof. Let $G: M \times [0, 1] \rightarrow N$ be a smooth homotopy from f to g in the sense of Definition 2.5. Choose a smooth function $\phi: \mathbb{R} \rightarrow [0, 1]$ with

$$\begin{aligned}\phi(t) &= 0 \text{ for } t \leq 0 \\ \phi(t) &= 1 \text{ for } t \geq 1\end{aligned}$$

Then

$$\Phi: M \times [0, 1] \rightarrow M \times [0, 1], (x, t) \mapsto (x, \phi(t))$$

is extension-smooth (it can be smoothly extended to $M \times \mathbb{R}$ and this implies that it is smooth in the sense of Definition 2.2, as Lee's definition is stronger by Lemma 2.3). Furthermore, it is easily verified that the composition of extension-smooth maps on arbitrary subsets is extension-smooth, such that

$$H := G \circ \Phi: M \times [0, 1] \rightarrow N$$

is extension-smooth. Then H is a homotopy from f to g and we can extend H to the smooth map

$$\overline{H}: M \times \mathbb{R} \rightarrow N, (x, t) \mapsto G(x, \phi(t)).$$

The converse follows directly from Lemma 2.3. □

2.2 Fiber Bundles

For convenience, let us state the definition of a fiber bundle.

Definition 2.7 (Fiber Bundle). Let G be a topological group acting effectively on a space F . A fiber bundle E over B with fiber F and structure group G consists of a surjective continuous map $p: E \rightarrow B$ and a collection of maps

$$(\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha))_\alpha,$$

called a trivialising atlas, for open sets $U_\alpha \subseteq B$, such that

- (i) The maps ϕ_α are homeomorphisms and the following diagrams commute:

$$\begin{array}{ccc} U_\alpha \times F & \xrightarrow{\phi_\alpha} & p^{-1}(U_\alpha) \\ & \searrow \pi \quad \swarrow p & \\ & U_\alpha & \end{array}$$

where π is the usual projection onto the first factor.

- (ii) For any two maps ϕ_α, ϕ_β there exists a continuous map

$$\tau_{\alpha, \beta}: U_\alpha \cap U_\beta \rightarrow G,$$

such that

$$\phi_\alpha^{-1} \circ \phi_\beta: (U_\alpha \cap U_\beta) \times F \rightarrow (U_\alpha \cap U_\beta) \times F$$

is given by $(b, f) \mapsto (b, \tau_{\alpha, \beta} \cdot f)$.

- (iii) $(U_\alpha)_\alpha$ is an open cover of B .
- (iv) The collection (ϕ_α) is maximal with respect to the first 3 conditions.

The maps ϕ_α are called local trivialisations, The sets U_α are called trivialisising neighbourhoods and the map $\tau_{\alpha,\beta}$ is called the transition function for ϕ_α, ϕ_β . Furthermore, we call E the total space, B the base space and F the fiber. We write

$$F \hookrightarrow E \xrightarrow{p} B$$

to denote the fiber bundle.

Remark 2.8. We require surjectivity of p to exclude the case $F = \emptyset$. The requirement, that G acts effectively guarantees for the transition maps to be unique. However, this is not a strong requirement to impose, since if G does not act effectively, i.e.

$$\Phi: G \rightarrow \text{Aut}(F)$$

is not injective we get an effective group action by taking a quotient of G :

$$G/\ker\Phi \rightarrow \text{Aut}(F).$$

It follows from the universal property of the quotient topology, that this map is still continuous.

Remark 2.9. The maximality condition for the trivialisising atlas is well-defined: Given any trivialisising atlas $\mathcal{F} = (\phi_\alpha)_\alpha$ satisfying conditions (i) - (iii) of Definition 2.7 we can define

$$\hat{\mathcal{F}} := \{\phi: U \times F \rightarrow p^{-1}(U) \mid U \subseteq B \text{ open, } \phi \text{ homeomorphism, for any } \phi_\alpha \in \mathcal{F} \text{ exists a map } \tau_{\alpha,\phi}: U \cap U_\alpha \rightarrow G, \text{ with } (\phi_\alpha^{-1} \circ \phi)(b, f) = (b, \tau_{\alpha,\phi}(b) \cdot f)\}.$$

This is the maximal trivialisising atlas induced by $\mathcal{F}$ satisfying (i) - (iii). Note however, that there is no unique trivialisising atlas, since beginning with a different $\mathcal{F}$ may yield a different $\hat{\mathcal{F}}$ (as is the case for smooth atlases of smooth manifolds).

Proof of above remark. Given $\phi: U \times F \rightarrow p^{-1}(U), \phi': U' \times F \rightarrow p^{-1}(U') \in \hat{\mathcal{F}}$, we claim that there exists $\tau: U \cap U' \rightarrow G$, such that

$$(\phi^{-1} \circ \phi')(b, f) = (b, \tau(b) \cdot f).$$

For $b_0 \in B$ choose $(\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha)) \in \mathcal{F}$ with $b_0 \in U_\alpha$. Then there exists

$$\tau_{\alpha,\phi}: U \cap U_\alpha \rightarrow G, \tau_{\alpha,\phi'}: U' \cap U_\alpha \rightarrow G$$

with

$$(\phi_\alpha^{-1} \circ \phi)(b, f) = (b, \tau_{\alpha,\phi}(b) \cdot f), (\phi_\alpha^{-1} \circ \phi')(b, f) = (b, \tau_{\alpha,\phi'}(b) \cdot f).$$

Then

$$(\phi^{-1} \circ \phi_\alpha) \circ (\phi_\alpha^{-1} \circ \phi'): (U_\alpha \cap U \cap U') \times F \rightarrow (U_\alpha \cap U \cap U') \times F$$

is given by

$$(b, f) \mapsto (b, \tau_{\alpha, \phi}(b)^{-1} \tau_{\alpha, \phi'}(b) \cdot f).$$

We conclude that around $\{b_0\} \times F$ we have

$$(\phi^{-1} \circ \phi')(b, f) = (b, \tau(b) \cdot f)$$

for $\tau(b) := \tau_{\alpha, \phi}(b)^{-1} \tau_{\alpha, \phi'}(b)$. Since group inversion and multiplication are continuous, τ is continuous. As b_0 was chosen arbitrary, such a map τ can be defined on all of $U \cap U'$. Furthermore, the group action of G is effective, hence τ is unique and the above argument shows that it is continuous. $\square$

For the transition functions we have the following Lemma, where again the effectiveness of the group action is crucial.

Lemma 2.10. *Let $F \hookrightarrow E \xrightarrow{p} B$ be a fiber bundle with structure group G . Then (using the notation in Definition 2.7):*

- (i) $\tau_{\alpha, \alpha}(b) = e$
- (ii) $\tau_{\alpha, \beta}^{-1}(b) = \tau_{\beta, \alpha}(b)$
- (iii) $\tau_{\gamma, \alpha}(b) \tau_{\alpha, \beta}(b) = \tau_{\gamma, \beta}(b)$

Proof. Let us first proof (iii), then (i) and (ii) follow directly from the group axioms. We compute:

$$\begin{aligned} (b, \tau_{\gamma, \beta}(b) \cdot f) &= (\phi_{\gamma}^{-1} \circ \phi_{\beta})(b, f) \\ &= (\phi_{\gamma}^{-1} \circ \phi_{\alpha} \circ \phi_{\alpha}^{-1} \circ \phi_{\beta})(b, f) = (b, \tau_{\gamma, \alpha}(b) \tau_{\alpha, \beta}(b) \cdot f). \end{aligned}$$

Since the transition functions are uniquely determined, the statement follows. $\square$

Interestingly, the transition functions already uniquely determine the fiber bundle, in a sense made precise by the following theorem.

Theorem 2.11 (Fiber bundle construction A). *Let B, F be (non-empty) topological spaces and G a topological group acting effectively on F . Suppose we are given the following data:*

- (i) *An open cover $\mathcal{U} := (U_{\alpha})_{\alpha}$ of B (where α ranges over some index set A)*
- (ii) *A collection $(\tau_{\alpha, \beta})$ of continuous maps*

$$\tau_{\alpha, \beta}: U_{\alpha} \cap U_{\beta} \rightarrow G$$

for each pair $U_{\alpha}, U_{\beta} \in \mathcal{U}$

- (iii) *$\tau_{\gamma, \alpha}(b) \tau_{\alpha, \beta}(b) = \tau_{\gamma, \beta}(b)$ for all $b \in U_{\alpha} \cap U_{\beta} \cap U_{\gamma}$*

Then there exists a unique (up to isomorphism) fiber bundle $F \hookrightarrow E \xrightarrow{p} B$ with structure group G , such that the maximal trivialising atlas is induced by a trivialising atlas of the form

$$(\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha))_\alpha,$$

such that $(\phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\alpha, \beta}(b) \cdot f)$.

Proof. Consider the space

$$E := \left(\coprod_{\alpha \in A} U_\alpha \times F \right) / \sim$$

where the equivalence relation $\sim$ is defined by

$$(b, f, \beta) \sim (b, \tau_{\alpha, \beta}(b) \cdot f, \alpha)$$

and let $p: E \rightarrow B$, $[(b, f, \alpha)] \mapsto b$. Furthermore, let $\pi: \coprod_{\alpha} U_\alpha \times F \rightarrow E$ be the projection map.

Step 1. $\sim$ is an equivalence relation.

This follows directly from Lemma 2.10: (i) implies reflexivity, (ii) implies symmetry and (iii) implies transitivity.

Step 2. p is well-defined, continuous and surjective.

Clearly p is well-defined and surjective as $F \neq \emptyset$. Continuity then follows directly from the universal property of the quotient topology.

Step 3. Defining the trivialising atlas

Consider

$$\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha), (b, f) \mapsto [(b, f, \alpha)].$$

Clearly ϕ_α is well-defined ($[(b, f, \alpha)] \in p^{-1}(U_\alpha)$) and continuous (the projection map π is continuous). The inverse is given by

$$\phi_\alpha^{-1}([(b, f, \beta)]) = (b, \tau_{\alpha, \beta}(b) \cdot f).$$

To show that ϕ_α^{-1} is continuous, we show that ϕ_α is an open map. To this end, let $U \times V \subseteq U_\alpha \times F$ be open and consider

$$\theta_{\alpha, \beta}: (U_\alpha \cap U_\beta) \times F \rightarrow (U_\alpha \cap U_\beta) \times F, (b, f) \mapsto (b, \tau_{\alpha, \beta}(b) \cdot f).$$

Clearly $\theta_{\alpha, \beta}$ is continuous (in fact it is a homeomorphism), hence

$$V_\beta := \theta_{\alpha, \beta}^{-1}((U \cap U_\beta) \times V) \subseteq (U_\alpha \cap U_\beta) \times F$$

is open and thus

$$\pi^{-1}(\phi_\alpha(U \times V)) = \coprod_{\beta \in A} V_\beta \subseteq \coprod_{\beta \in A} U_\beta \times F$$

is open. Thus $\phi_\alpha(U \times V)$ is open in E and since $p^{-1}(U_\alpha)$ is also open, we conclude that ϕ_α is an open map. Furthermore, the diagram

$$\begin{array}{ccc}
U_\alpha \times F & \xrightarrow{\phi_\alpha} & p^{-1}(U_\alpha) \\
& \searrow & \swarrow p \\
& U_\alpha &
\end{array}$$

commutes, such that ϕ_α is a local trivialisation. Moreover,

$$(\phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\alpha, \beta}(b) \cdot f).$$

So the collection of maps $(\phi_\alpha)_\alpha$ is a trivialising atlas with the desired transition functions, making $F \hookrightarrow E \xrightarrow{p} B$ into a fiber bundle with structure group G .

Step 4. Uniqueness

Given two fiber bundles

$$\begin{aligned}
\delta &:= (F \hookrightarrow E \xrightarrow{p} B) \\
\epsilon &:= (F \hookrightarrow \tilde{E} \xrightarrow{\tilde{p}} B)
\end{aligned}$$

with structure group G and trivialising atlases

$$\begin{aligned}
(\phi_\alpha &: U_\alpha \times F \rightarrow p^{-1}(U_\alpha))_{\alpha \in A} \\
(\tilde{\phi}_\alpha &: U_\alpha \times F \rightarrow \tilde{p}^{-1}(U_\alpha))_{\alpha \in A}
\end{aligned}$$

respectively, such that

$$(\phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\alpha, \beta}(b) \cdot f) = (\tilde{\phi}_\alpha^{-1} \circ \tilde{\phi}_\beta)(b, f).$$

We can then construct an isomorphism $\Gamma: \delta \rightarrow \epsilon$ as follows: for $x \in E$ choose $U_\alpha \in \mathcal{U}$ with $p(x) \in U_\alpha$. Then define

$$\Gamma(x) := (\tilde{\phi}_\alpha \circ \phi_\alpha^{-1})(x).$$

Well-definedness of Γ follows from the commutativity of the following diagram:

$$\begin{array}{ccccc}
& & (U_\alpha \cap U_\beta) \times F & & \\
& \swarrow \phi_\beta & \downarrow \tau & \searrow \tilde{\phi}_\beta & \\
p^{-1}(U_\alpha \cap U_\beta) & & (U_\alpha \cap U_\beta) \times F & & \tilde{p}^{-1}(U_\alpha \cap U_\beta) \\
& \swarrow \phi_\alpha & \downarrow & \searrow \tilde{\phi}_\alpha & \\
& & (U_\alpha \cap U_\beta) \times F & &
\end{array}$$

where $\tau(b, f) := (b, \tau_{\alpha, \beta}(b) \cdot f)$. Locally Γ is given by a continuous map, so it is continuous. Analogously we define the inverse map. Furthermore,

$$\begin{array}{ccc}
E & \xrightarrow{\Gamma} & \tilde{E} \\
& \searrow p & \swarrow \tilde{p} \\
& B &
\end{array}$$

commutes, so Γ is a bundle isomorphism. This completes the proof. $\square$

We may also formulate Theorem 2.11 slightly different, if we are given the fibers $F_x = p^{-1}(x)$ as sets. This may be more practical in some situations, as the description of the bundle is more explicit:

Theorem 2.12 (Fiber bundle construction B). *Let B, F be (non-empty) topological spaces and G a topological group acting effectively on F . Given sets F_x for each $x \in B$ we define*

$$E := \coprod_{x \in B} F_x$$

and $p: E \rightarrow B$ by sending each element in F_x to x . Suppose we are given the following data:

- (i) *An open cover $\mathcal{U} := (U_\alpha)_\alpha$ of B (where α ranges over some index set A)*
- (ii) *A collection of bijections $\phi_\alpha: U_\alpha \times F \rightarrow p^{-1}(U_\alpha)$, that restrict to bijections $\{x\} \times F \rightarrow F_x$*
- (iii) *A collection $(\tau_{\alpha,\beta})$ of continuous maps*

$$\tau_{\alpha,\beta}: U_\alpha \cap U_\beta \rightarrow G$$

for each pair $U_\alpha, U_\beta \in \mathcal{U}$, such that $(\phi_\alpha^{-1} \circ \phi_\beta)(b, f) = (b, \tau_{\alpha,\beta}(b) \cdot f)$

Then E has a unique topology making $F \hookrightarrow E \xrightarrow{p} B$ into a fiber bundle with structure group G , where the maps ϕ_α are local trivialisations.

Proof. Define the topology on E as the final topology with respect to the maps ϕ_α , i.e.

$$V \subseteq E \text{ is open} \Leftrightarrow \phi_\alpha^{-1}(V) \subseteq U_\alpha \times F \text{ is open for each } \alpha$$

We show that the maps ϕ_α are open with this topology. To this end, let $W \subseteq U_\alpha \times F$ be open. Then

$$\phi_\beta^{-1}(\phi_\alpha(W)) = (\phi_\beta^{-1} \circ \phi_\alpha)(W \cap ((U_\alpha \cap U_\beta) \times F)) \subseteq U_\beta \times F$$

is open, since $\phi_\beta \circ \phi_\alpha$ is a homeomorphism by condition (iii). Thus ϕ_α is a homeomorphism. In particular, for any open set $U \subseteq U_\alpha$, we get that

$$p^{-1}(U) = \phi_\alpha(U \times F)$$

is open, since by (ii) the following diagram commutes:

$$\begin{array}{ccc} U_\alpha \times F & \xrightarrow{\phi_\alpha} & p^{-1}(U_\alpha) \\ & \searrow & \swarrow p \\ & U_\alpha & \end{array}$$

So p is also continuous. Moreover, as $F \neq \emptyset$, we get that p is surjective. It remains to show uniqueness. So let E be equipped with any topology, such that p is continuous and ϕ_α are homeomorphisms. Let $V \subseteq E$, such that

$$W_\alpha := \phi_\alpha^{-1}(V) \subseteq U_\alpha \times F$$

is open for all α . Then

$$V = \bigcup_{\alpha} W_\alpha,$$

since the U_α cover B (and hence $E = \bigcup_{\alpha} \text{im} \phi_\alpha$). As the ϕ_α are homeomorphisms and $p^{-1}(U_\alpha)$ is open, we conclude that V is open. Furthermore, the topology can not be finer than the final topology defined above, such that the two must coincide. $\square$

Given a fiber bundle, Theorem 2.11 allows to change the fiber:

Corollary 2.13 (Changing the fiber). *Let $F \hookrightarrow E \xrightarrow{p} B$ be a fiber bundle with structure group G . Given a space F' upon which G acts effectively, there exists a unique fiber bundle*

$$F' \hookrightarrow E' \xrightarrow{p'} B$$

which has the same trivialising neighbourhoods (U_α) and the same transition functions $(\tau_{\alpha,\beta})$ as $F \hookrightarrow E \xrightarrow{p} B$.

Theorem 2.11 tells us, that the new bundle still retains all of the information of the old bundle, as it is uniquely determined by the transition functions.

2.2.1 Smooth bundles

If we are in the smooth category, we can analogously define smooth fiber bundles by requiring the spaces E, B and F to be smooth manifolds (where either F or B has no boundary), the group G to be a Lie group and all maps to be smooth. In particular, the local trivialisation are diffeomorphisms. All of the above results also hold in the smooth category.

Theorem 2.14 (Fiber bundle construction A'). *The conclusions of Theorem 2.11 remain true in the smooth category.*

Proof. The only difficulty is in defining the smooth structure on

$$E := \left(\coprod_{\alpha \in A} U_\alpha \times F \right) / \sim$$

This smooth structure is defined via the local trivialisations (ϕ_α) as follows: Choose an open cover (V_α) of B , such that

- $V_\alpha \subseteq U_\alpha$
- there exist smooth charts $h_\alpha: W_\alpha \rightarrow V_\alpha$

Furthermore, choose a smooth atlas

$$(k_i: W_i^F \rightarrow V_i^F)_i$$

of F . Then

$$(\phi_\alpha \circ (h_\alpha \times k_i): W_\alpha \times W_i^F \rightarrow E)_{i,\alpha}$$

defines a smooth atlas for E , since the maps $\phi_\alpha^{-1} \circ \phi_\beta$ are diffeomorphisms. It is then easy to verify, that this atlas satisfies the first countability and Hausdorff axioms. Furthermore, this smooth structure is uniquely determined by requiring the local trivialisations to be diffeomorphisms. $\square$

Theorem 2.15 (Fiber bundle construction B'). *The conclusions of Theorem 2.12 remain true in the smooth category.*

Proof. As in the above theorem, the local trivialisations determine a unique smooth structure on E . $\square$

2.2.2 Whitney Sums

References

- [GP74] Victor Guillemin and Alan Pollack. *Differential Topology*. 1974.
- [Lee11] John M. Lee. *Introduction to Topological Manifolds*, volume 202. Springer, New York, 2011.
- [Mil63] John W. Milnor. *Morse Theory*. Princeton University Press, 1963.
- [Vin24] Johannes Vinnemeier. Homology and perfect morse functions for manifolds with boundary and their gluings. 2024.