

Laplace's Equations

Johnson Zhang

June 26, 2025

Content of this report

- Fundamental Solutions to Laplace's Equation and Poisson's Equations.

Content of this report

- Fundamental Solutions to Laplace's Equation and Poisson's Equations.
- Properties of harmonic functions.

Laplace's Equations

Laplace's Equations

Definition

Let $u(x_1, \dots, x_n) : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a unknown function, the Laplacian of the function is defined by

$$\Delta u = \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2}. \quad (0.1)$$

Laplace equation is of the form

$$\Delta u = 0 \quad (0.2)$$

for all $x \in U$, any function satisfying the above is also called harmonic.

Laplace's Equations

Lemma

Laplace's equation is invariant under rotation, i.e if P is an orthogonal matrix with entries p_{ij} , then $\Delta u(Px) = 0$ for all $x \in U$.

Proof.

By Chain rule, we have

$$D_{x_i}(u(Px)) = \sum_{k=1}^n D_{x_k}(u(Px)p_{ik}) \quad (0.3)$$

and similarly

$$D_{x_i x_j}(u(Px)) = \sum_{l=1}^n \sum_{k=1}^n D_{x_k x_l}(u(Px)p_{ik}p_{jl}). \quad (0.4)$$

Since P is orthogonal, thus $p_{ik}p_{il} = 1$ iff $k = l$ and hence

$$\Delta u(Px) = \sum_{i=1}^n \sum_{l=1}^n \sum_{k=1}^n D_{x_k x_l}(u(Px)p_{ik}p_{il}) = \Delta u = 0. \quad (0.5)$$



Fundamental Solutions

Given this, we seek for radial solutions which takes the form $u(x) = v(r)$ where $r = |x|$. Denote $r = (x_1^2 + \cdots + x_n^2)^{1/2}$, let $x \neq 0$, then we see that

$$\frac{\partial r}{\partial x_i} = (x_1^2 + \cdots + x_n^2)^{-1/2} x_i = \frac{x_i}{r} \quad (0.6)$$

Fundamental Solutions

Given this, we seek for radial solutions which takes the form $u(x) = v(r)$ where $r = |x|$. Denote $r = (x_1^2 + \cdots + x_n^2)^{1/2}$, let $x \neq 0$, then we see that

$$\frac{\partial r}{\partial x_i} = (x_1^2 + \cdots + x_n^2)^{-1/2} x_i = \frac{x_i}{r} \quad (0.6)$$

and by Chain rule we have

$$u_{x_i} = v'(r) \frac{x_i}{r} \quad (0.7)$$

$$u_{x_i x_i} = v''(r) \frac{x_i^2}{r^2} + v'(r) \frac{r - x_i \frac{x_i}{r}}{r^2}, v''(r) \frac{x_i^2}{r^2} + v'(r) \left(\frac{1}{r} - \frac{x_i^2}{r^3} \right) \quad (0.8)$$

Fundamental Solutions

using the fact that $x_1^2 + \cdots + x_n^2 = r^2$, we would have

$$\Delta u = \sum_{i=1}^n \left(v''(r) \frac{x_i^2}{r^2} + v'(r) \left(\frac{1}{r} - \frac{x_i^2}{r^3} \right) \right) = v''(r) + \frac{n-1}{r} v'(r). \quad (0.9)$$

Fundamental Solutions

using the fact that $x_1^2 + \cdots + x_n^2 = r^2$, we would have

$$\Delta u = \sum_{i=1}^n \left(v''(r) \frac{x_i^2}{r^2} + v'(r) \left(\frac{1}{r} - \frac{x_i^2}{r^3} \right) \right) = v''(r) + \frac{n-1}{r} v'(r). \quad (0.9)$$

by letting $\Delta u = 0$, we obtain the following ODE:

$$v''(r) + \frac{n-1}{r} v'(r) = 0. \quad (0.10)$$

Fundamental Solutions

using the fact that $x_1^2 + \cdots + x_n^2 = r^2$, we would have

$$\Delta u = \sum_{i=1}^n \left(v''(r) \frac{x_i^2}{r^2} + v'(r) \left(\frac{1}{r} - \frac{x_i^2}{r^3} \right) \right) = v''(r) + \frac{n-1}{r} v'(r). \quad (0.9)$$

by letting $\Delta u = 0$, we obtain the following ODE:

$$v''(r) + \frac{n-1}{r} v'(r) = 0. \quad (0.10)$$

Now if $v'(r) \neq 0$, we see that

$$\log(v'(r))' = \frac{v''(r)}{v'(r)} = \frac{1-n}{r} \quad (0.11)$$

and we integrate with respect to r , we have

$$\log(v'(r)) = (1 - n) \log r + C \quad (0.12)$$

Fundamental Solutions

and we integrate with respect to r , we have

$$\log(v'(r)) = (1 - n) \log r + C \quad (0.12)$$

which simplifies to $v'(r) = C_0/r^{n-1}$ for some constant C_0 , hence the general solution $v(r)$ takes the form

$$v(r) = \begin{cases} A \log r + B & n = 2 \\ \frac{C}{r^{n-2}} + D & n \geq 3 \end{cases} \quad (0.13)$$

where A, B, C, D are all constants. This motivates the fundamental solution to Laplace's equation.

Definition

The function defined by

$$\Phi(x) = \begin{cases} -\frac{1}{2\pi} \log |x| & n = 2 \\ \frac{1}{n(n-2)\alpha(n)} \cdot \frac{1}{|x|^{n-2}} & n \geq 3 \end{cases} \quad (0.14)$$

is the fundamental solution to Laplace's equation $\Delta u = 0$ for all $x \in \mathbb{R}^n / \{0\}$ where $\alpha(n)$ denotes the volume of the unit ball in \mathbb{R}^n .

Poisson's Equations

Laplace's equation is a special class of Poisson's equations, we define Poisson's equation as

Definition

For a given $f : \mathbb{R}^n \rightarrow \mathbb{R}$, and the unknown $u(x_1, \dots, x_n) : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$, the Poisson's equation takes the form

$$-\Delta u = f. \tag{0.15}$$

Fundamental Solutions

The following theorem will state the fundamental solution of Poisson's equations:

The following theorem will state the fundamental solution of Poisson's equations:

Theorem

Define $u(x) = \int_{\mathbb{R}^n} \Phi(x - y)f(y)dy$ and assume that $f \in C_c^2(\mathbb{R}^n)$, then:

- (i) $u \in C^2(\mathbb{R}^n)$;
- (ii) $-\Delta u = f$ in \mathbb{R}^n .

The following theorem will state the fundamental solution of Poisson's equations:

Theorem

Define $u(x) = \int_{\mathbb{R}^n} \Phi(x - y)f(y)dy$ and assume that $f \in C_c^2(\mathbb{R}^n)$, then:

- (i) $u \in C^2(\mathbb{R}^n)$;
- (ii) $-\Delta u = f$ in \mathbb{R}^n .

We will prove both results separately:

For (i) :Use the property of a convolution, we have

$$u(x) = \int_{\mathbb{R}^n} \Phi(y)f(x-y)dy \quad (0.16)$$

For (i) :Use the property of a convolution, we have

$$u(x) = \int_{\mathbb{R}^n} \Phi(y)f(x-y)dy \quad (0.16)$$

then denote e_i as the unit vector in i th position of $x = (x_1, \dots, x_n)$, we have

$$\begin{aligned} \frac{\partial u}{\partial x_i}(x) &= \lim_{h \rightarrow 0} \frac{u(x + he_i) - u(x)}{h} \\ &= \int_{\mathbb{R}^n} \Phi(y) \cdot \lim_{h \rightarrow 0} \left[\frac{f(x + he_i - y) - f(x - y)}{h} \right] dy \\ &= \int_{\mathbb{R}^n} \Phi(y) \cdot \frac{\partial f}{\partial x_i}(x - y) dy. \end{aligned}$$

For (i) :Use the property of a convolution, we have

$$u(x) = \int_{\mathbb{R}^n} \Phi(y)f(x-y)dy \quad (0.16)$$

then denote e_i as the unit vector in i th position of $x = (x_1, \dots, x_n)$, we have

$$\begin{aligned} \frac{\partial u}{\partial x_i}(x) &= \lim_{h \rightarrow 0} \frac{u(x + he_i) - u(x)}{h} \\ &= \int_{\mathbb{R}^n} \Phi(y) \cdot \lim_{h \rightarrow 0} \left[\frac{f(x + he_i - y) - f(x - y)}{h} \right] dy \\ &= \int_{\mathbb{R}^n} \Phi(y) \cdot \frac{\partial f}{\partial x_i}(x - y) dy. \end{aligned}$$

Similarly one can show

$$\frac{\partial u}{\partial x_i \partial x_j}(x) = \int_{\mathbb{R}^n} \Phi(y) \cdot \frac{\partial^2 f}{\partial x_i \partial x_j}(x - y) dy. \quad (0.17)$$

For (i) :Use the property of a convolution, we have

$$u(x) = \int_{\mathbb{R}^n} \Phi(y)f(x-y)dy \quad (0.16)$$

then denote e_i as the unit vector in i th position of $x = (x_1, \dots, x_n)$, we have

$$\begin{aligned} \frac{\partial u}{\partial x_i}(x) &= \lim_{h \rightarrow 0} \frac{u(x + he_i) - u(x)}{h} \\ &= \int_{\mathbb{R}^n} \Phi(y) \cdot \lim_{h \rightarrow 0} \left[\frac{f(x + he_i - y) - f(x - y)}{h} \right] dy \\ &= \int_{\mathbb{R}^n} \Phi(y) \cdot \frac{\partial f}{\partial x_i}(x - y) dy. \end{aligned}$$

Similarly one can show

$$\frac{\partial u}{\partial x_i \partial x_j}(x) = \int_{\mathbb{R}^n} \Phi(y) \cdot \frac{\partial^2 f}{\partial x_i \partial x_j}(x - y) dy. \quad (0.17)$$

Since $f \in C^2_C(\mathbb{R}^n)$, the second partial derivative is continuous w.r.t x , hence $u \in C^2(\mathbb{R}^n)$.

Since $f \in C^2_{\mathbb{C}}(\mathbb{R}^n)$, the second partial derivative is continuous w.r.t x , hence $u \in C^2(\mathbb{R}^n)$. For (ii): We split the integral into two parts to avoid the blow-up of Φ near 0, $\forall \epsilon > 0$ fixed, denote $B(0, \epsilon)$ as the ball centered at 0 with radius ϵ , write

$$u(x) = \int_{B(0, \epsilon)} \Phi(y) f(x - y) dy + \int_{\mathbb{R}^n / B(0, \epsilon)} \Phi(y) f(x - y) dy \quad (0.18)$$

Since $f \in C^2_C(\mathbb{R}^n)$, the second partial derivative is continuous w.r.t x , hence $u \in C^2(\mathbb{R}^n)$. For (ii): We split the integral into two parts to avoid the blow-up of Φ near 0, $\forall \epsilon > 0$ fixed, denote $B(0, \epsilon)$ as the ball centered at 0 with radius ϵ , write

$$u(x) = \int_{B(0, \epsilon)} \Phi(y) f(x-y) dy + \int_{\mathbb{R}^n / B(0, \epsilon)} \Phi(y) f(x-y) dy \quad (0.18)$$

hence

$$\Delta u(x) = \int_{B(0, \epsilon)} \Phi(y) \Delta_x f(x-y) dy + \int_{\mathbb{R}^n / B(0, \epsilon)} \Phi(y) \Delta_x f(x-y) dy. \quad (0.19)$$

Since $f \in C^2_C(\mathbb{R}^n)$, the second partial derivative is continuous w.r.t x , hence $u \in C^2(\mathbb{R}^n)$. For (ii): We split the integral into two parts to avoid the blow-up of Φ near 0, $\forall \epsilon > 0$ fixed, denote $B(0, \epsilon)$ as the ball centered at 0 with radius ϵ , write

$$u(x) = \int_{B(0, \epsilon)} \Phi(y) f(x-y) dy + \int_{\mathbb{R}^n / B(0, \epsilon)} \Phi(y) f(x-y) dy \quad (0.18)$$

hence

$$\Delta u(x) = \int_{B(0, \epsilon)} \Phi(y) \Delta_x f(x-y) dy + \int_{\mathbb{R}^n / B(0, \epsilon)} \Phi(y) \Delta_x f(x-y) dy. \quad (0.19)$$

Denote the first term as I_ϵ , second term as J_ϵ . Our first goal is to bound I_ϵ , proceed as follows:

When $n = 2$, we have

$$\begin{aligned} |I_\epsilon| &= \left| - \int_{B(0,\epsilon)} \frac{1}{2\pi} \log |y| \cdot \Delta_x f(x-y) dy \right| \\ &\leq C \|Df\|_{L^\infty(\mathbb{R}^n)} \cdot \left| \int_{B(0,\epsilon)} \log |y| dy \right| \end{aligned}$$

When $n = 2$, we have

$$\begin{aligned} |I_\epsilon| &= \left| - \int_{B(0,\epsilon)} \frac{1}{2\pi} \log |y| \cdot \Delta_x f(x-y) dy \right| \\ &\leq C \|Df\|_{L^\infty(\mathbb{R}^n)} \cdot \left| \int_{B(0,\epsilon)} \log |y| dy \right| \end{aligned}$$

for some constant C , and we further have

$$\int_{B(0,\epsilon)} \log |y| dy = \int_0^\epsilon \int_0^{2\pi} \log |r| r d\theta dr = \epsilon^2 \log |\epsilon| \quad (0.20)$$

When $n = 2$, we have

$$\begin{aligned} |I_\epsilon| &= \left| - \int_{B(0,\epsilon)} \frac{1}{2\pi} \log |y| \cdot \Delta_x f(x-y) dy \right| \\ &\leq C \|Df\|_{L^\infty(\mathbb{R}^n)} \cdot \left| \int_{B(0,\epsilon)} \log |y| dy \right| \end{aligned}$$

for some constant C , and we further have

$$\int_{B(0,\epsilon)} \log |y| dy = \int_0^\epsilon \int_0^{2\pi} \log |r| r d\theta dr = \epsilon^2 \log |\epsilon| \quad (0.20)$$

so we have $|I_\epsilon| < C\epsilon^2 |\log \epsilon|$ for some constant C . When $n \geq 3$, similarly we can show that $|I_\epsilon| < C\epsilon$, hence the term I_ϵ is bounded in terms of arbitrary ϵ .

Now for J_ϵ , first note that $\Delta_x f(x - y) = \Delta_y f(x - y)$, we have

$$J_\epsilon = \int_{\mathbb{R}^n/B(0,\epsilon)} \Phi(y) \Delta_y f(x - y) dy. \quad (0.21)$$

Now for J_ϵ , first note that $\Delta_x f(x - y) = \Delta_y f(x - y)$, we have

$$J_\epsilon = \int_{\mathbb{R}^n/B(0,\epsilon)} \Phi(y) \Delta_y f(x - y) dy. \quad (0.21)$$

To compute this integral, we need some technical lemmas:

Now for J_ϵ , first note that $\Delta_x f(x - y) = \Delta_y f(x - y)$, we have

$$J_\epsilon = \int_{\mathbb{R}^n/B(0,\epsilon)} \Phi(y) \Delta_y f(x - y) dy. \quad (0.21)$$

To compute this integral, we need some technical lemmas:

Lemma

(Gauss-Green Theorem) Let $u(x_1, \dots, x_n) \in C^1(\overline{U})$, let $\nu = (\nu_1, \dots, \nu_n)$ be the outward pointing unit normal vector of U defined on ∂U then

$$\int_U u_{x_i} dx = \int_{\partial U} u \nu_i dS. \quad (0.22)$$

We can use this lemma to further derive an integration by parts formula:

Lemma

(Integration by Parts) Let $u, v \in C^1(\overline{U})$, then

$$\int_U u_{x_i} v dx = - \int_U u v_{x_i} dx + \int_{\partial U} u v \nu^i dS. \quad (0.23)$$

We can use this lemma to further derive an integration by parts formula:

Lemma

(Integration by Parts) Let $u, v \in C^1(\overline{U})$, then

$$\int_U u_{x_i} v dx = - \int_U u v_{x_i} dx + \int_{\partial U} u v \nu^i dS. \quad (0.23)$$

Then we may rewrite J_ϵ as

$$J_\epsilon = - \int_{\mathbb{R}^n/B(0,\epsilon)} D\Phi(y) \cdot D_y f(x-y) dy + \int_{\partial B(0,\epsilon)} \Phi(y) \nu Df(x-y) dS(y) \quad (0.24)$$

We can use this lemma to further derive an integration by parts formula:

Lemma

(Integration by Parts) Let $u, v \in C^1(\overline{U})$, then

$$\int_U u_{x_i} v dx = - \int_U u v_{x_i} dx + \int_{\partial U} u v \nu^i dS. \quad (0.23)$$

Then we may rewrite J_ϵ as

$$J_\epsilon = - \int_{\mathbb{R}^n/B(0,\epsilon)} D\Phi(y) \cdot D_y f(x-y) dy + \int_{\partial B(0,\epsilon)} \Phi(y) \nu Df(x-y) dS(y) \quad (0.24)$$

where ν is the inward pointing unit normal vector. We now denote the first term by K_ϵ and the second term by L_ϵ , then using the same technique as we did for I_ϵ , we can check that

$$|L_\epsilon| \leq C \|Df\|_{L^\infty(\mathbb{R}^n)} \int_{\partial B(0,\epsilon)} |\Phi(y)| dS(y) \leq \begin{cases} C\epsilon |\log \epsilon| & n = 2 \\ C\epsilon & n \geq 3 \end{cases} \quad (0.25)$$

$$|L_\epsilon| \leq C \|Df\|_{L^\infty(\mathbb{R}^n)} \int_{\partial B(0,\epsilon)} |\Phi(y)| dS(y) \leq \begin{cases} C\epsilon |\log \epsilon| & n = 2 \\ C\epsilon & n \geq 3 \end{cases} \quad (0.25)$$

hence L_ϵ is also bounded in terms of ϵ . We again perform integration by parts in K_ϵ , and we have

$$\begin{aligned} K_\epsilon &= \int_{\mathbb{R}^n/B(0,\epsilon)} \Delta \Phi(y) f(x-y) dy - \int_{\partial B(0,\epsilon)} \nu D\Phi(y) f(x-y) dS(y) \\ &= - \int_{\partial B(0,\epsilon)} \nu D\Phi(y) f(x-y) dS(y). \end{aligned}$$

Since $D\Phi(y) = \frac{-1}{n\alpha(n)} \frac{y}{|y|^n}$ when $y \neq 0$ and $\nu = -y/|y| = -y/\epsilon$ on $\partial B(0, \epsilon)$, so we will have

$$\nu D\Phi(y) = \frac{1}{n\alpha(n)\epsilon^{n-1}} \quad (0.26)$$

Since $D\Phi(y) = \frac{-1}{n\alpha(n)} \frac{y}{|y|^n}$ when $y \neq 0$ and $\nu = -y/|y| = -y/\epsilon$ on $\partial B(0, \epsilon)$, so we will have

$$\nu D\Phi(y) = \frac{1}{n\alpha(n)\epsilon^{n-1}} \quad (0.26)$$

also $n\alpha(n)\epsilon^{n-1}$ is the surface area of the ball $\partial B(0, \epsilon)$, so

$$K_\epsilon = -\frac{1}{n\alpha(n)\epsilon^{n-1}} \int_{\partial B(0, \epsilon)} f(x-y) dS(y) = -\oint_{\partial B(0, \epsilon)} f(y) dS(y) \rightarrow -f(x) \quad (0.27)$$

as $\epsilon \rightarrow 0$.

Since $D\Phi(y) = \frac{-1}{n\alpha(n)} \frac{y}{|y|^n}$ when $y \neq 0$ and $\nu = -y/|y| = -y/\epsilon$ on $\partial B(0, \epsilon)$, so we will have

$$\nu D\Phi(y) = \frac{1}{n\alpha(n)\epsilon^{n-1}} \quad (0.26)$$

also $n\alpha(n)\epsilon^{n-1}$ is the surface area of the ball $\partial B(0, \epsilon)$, so

$$K_\epsilon = -\frac{1}{n\alpha(n)\epsilon^{n-1}} \int_{\partial B(0, \epsilon)} f(x-y) dS(y) = -\oint_{\partial B(0, \epsilon)} f(y) dS(y) \rightarrow -f(x) \quad (0.27)$$

as $\epsilon \rightarrow 0$. Since we have shown that I_ϵ, L_ϵ are all bounded by terms of ϵ so by setting $\epsilon \rightarrow 0$ they will vanish as well. Hence we have

$$-\Delta u(x) = f(x). \quad (0.28)$$

Properties of Harmonic functions

Now we shall introduce some properties of harmonic functions:

- Mean- Value Property

Properties of Harmonic functions

Now we shall introduce some properties of harmonic functions:

- Mean- Value Property
- Maximum Principle

Properties of Harmonic functions

Now we shall introduce some properties of harmonic functions:

- Mean- Value Property
- Maximum Principle
- Smoothness

Properties of Harmonic functions

Now we shall introduce some properties of harmonic functions:

- Mean- Value Property
- Maximum Principle
- Smoothness
- Liouville's Theorem

Properties of Harmonic functions

Now we shall introduce some properties of harmonic functions:

- Mean- Value Property
- Maximum Principle
- Smoothness
- Liouville's Theorem
- Harnack's Inequality

Properties of Harmonic functions

Now we shall introduce some properties of harmonic functions:

- Mean- Value Property
- Maximum Principle
- Smoothness
- Liouville's Theorem
- Harnack's Inequality
- Convergences

Mean-Value Property

Theorem

Let $U \subset \mathbb{R}^n$ open, if $u(x_1, \dots, x_n) \in C^2(U)$ is harmonic then for each ball $B(x, r) \subset U$,

$$u(x) = \oint_{\partial B(x,r)} u(y) dS(y) = \int_{B(x,r)} u(y) dy \quad (0.29)$$

where

$$\oint_{\partial B(x,r)} u(y) dS(y) = \frac{1}{n\alpha(n)r^{n-1}} \int_{\partial B(x,r)} u(y) dS(y), \quad (0.30)$$

$$\int_{B(x,r)} u(y) dy = \frac{1}{\alpha(n)r^n} \int_{B(x,r)} u(y) dy \quad (0.31)$$

$\alpha(n)$ is the volume of the unit ball in \mathbb{R}^n and $n\alpha(n)$ is the surface area of the unit ball in \mathbb{R}^n .

Let $u(x_1, \dots, x_n) \in C^2(U)$ be harmonic, and define

$$\phi(r) = \begin{cases} \int_{\partial B(x,r)} u(y) dS(y) & r > 0 \\ u(x) & r = 0 \end{cases}. \quad (0.32)$$

Let $u(x_1, \dots, x_n) \in C^2(U)$ be harmonic, and define

$$\phi(r) = \begin{cases} \int_{\partial B(x,r)} u(y) dS(y) & r > 0 \\ u(x) & r = 0 \end{cases}. \quad (0.32)$$

If u is a smooth function, then $\lim_{r \rightarrow 0} \phi(r) = u(x)$ and hence ϕ is continuous. Note that by change of variables, we have

$$\phi(r) = \int_{\partial B(0,1)} u(x + rz) dS(z) \quad (0.33)$$

Let $u(x_1, \dots, x_n) \in C^2(U)$ be harmonic, and define

$$\phi(r) = \begin{cases} \int_{\partial B(x,r)} u(y) dS(y) & r > 0 \\ u(x) & r = 0 \end{cases}. \quad (0.32)$$

If u is a smooth function, then $\lim_{r \rightarrow 0} \phi(r) = u(x)$ and hence ϕ is continuous. Note that by change of variables, we have

$$\phi(r) = \int_{\partial B(0,1)} u(x + rz) dS(z) \quad (0.33)$$

and we take the derivative of $\phi(r)$:

$$\begin{aligned}
\phi'(r) &= \oint_{\partial B(0,1)} \nabla u(x + rz) \cdot z dS(z) \\
&= \oint_{\partial B(x,r)} \nabla u(y) \cdot \frac{y - x}{r} dS(y) \\
&= \oint_{\partial B(x,r)} \frac{\partial u}{\partial \nu}(y) dS(y) \\
&= \frac{1}{n\alpha(n)r^{n-1}} \int_{\partial B(x,r)} \frac{\partial u}{\partial \nu}(y) dS(y) \\
&= \frac{1}{n\alpha(n)r^{n-1}} \int_{B(x,r)} \nabla(\nabla u) dy \\
&= \frac{1}{n\alpha(n)r^{n-1}} \int_{B(x,r)} \Delta u(y) dy \equiv 0.
\end{aligned}$$

which means ϕ is a constant function, and hence we have

$$u(x) = \oint_{\partial B(x,r)} u(y) dS(y). \quad (0.34)$$

which means ϕ is a constant function, and hence we have

$$u(x) = \oint_{\partial B(x,r)} u(y) dS(y). \quad (0.34)$$

Furthermore we have

$$\begin{aligned} \int_{B(x,r)} u(y) dy &= \int_0^r \left(\int_{\partial B(x,s)} u dS \right) ds \\ &= u(x) \int_0^r n\alpha(n)s^{n-1} ds \\ &= \alpha(n)r^n u(x). \end{aligned}$$

which means ϕ is a constant function, and hence we have

$$u(x) = \oint_{\partial B(x,r)} u(y) dS(y). \quad (0.34)$$

Furthermore we have

$$\begin{aligned} \int_{B(x,r)} u(y) dy &= \int_0^r \left(\int_{\partial B(x,s)} u dS \right) ds \\ &= u(x) \int_0^r n\alpha(n)s^{n-1} ds \\ &= \alpha(n)r^n u(x). \end{aligned}$$

and that finishes the proof.

Theorem

If $u \in C^2(U)$ satisfies

$$u(x) = \oint_{\partial B(x,r)} u dS \quad (0.35)$$

for each ball $B(x,r) \in U$, then u is harmonic.

Theorem

If $u \in C^2(U)$ satisfies

$$u(x) = \oint_{\partial B(x,r)} u dS \quad (0.35)$$

for each ball $B(x,r) \in U$, then u is harmonic.

Proof.

Suppose $\Delta u \neq 0$, then $\exists B(x,r) \in U$ such that $\Delta u > 0$ say, but then

$$\phi'(r) = \frac{r}{n} \oint_{B(x,r)} \Delta u(y) dy > 0 \quad (0.36)$$

which contradicts the fact that ϕ is a constant function. □

Maximum Principle

Theorem

(Strong Maximum Principle) Suppose $u \in C^2(U) \cap C(\bar{U})$ is harmonic within U , then

(i) $\max_{\bar{U}} u = \max_{\partial U} u$;

(ii) Furthermore, if U is connected and there exists a point $x_0 \in U$ such that $u(x_0) = \max_{\bar{U}} u$, then u is constant within U .

Maximum Principle

Theorem

(Strong Maximum Principle) Suppose $u \in C^2(U) \cap C(\bar{U})$ is harmonic within U , then

(i) $\max_{\bar{U}} u = \max_{\partial U} u$;

(ii) Furthermore, if U is connected and there exists a point $x_0 \in U$ such that $u(x_0) = \max_{\bar{U}} u$, then u is constant within U .

It suggests that if u is harmonic on a bounded domain U , then u attains its maximum value on the boundary of U

Proof.

We will prove the second statement since the first followed from the second.

Proof.

We will prove the second statement since the first followed from the second. Suppose $\exists x_0 \in U$ such that $u(x_0) = \max_{\bar{U}} u(x) \equiv M$. Then $\forall r \in (0, \text{dist}(x_0, \partial U))$, the mean-value property states that

$$M = u(x_0) = \oint_{B(x_0, r)} u(y) dy \quad (0.37)$$

Proof.

We will prove the second statement since the first followed from the second. Suppose $\exists x_0 \in U$ such that $u(x_0) = \max_{\bar{U}} u(x) \equiv M$. Then $\forall r \in (0, \text{dist}(x_0, \partial U))$, the mean-value property states that

$$M = u(x_0) = \oint_{B(x_0, r)} u(y) dy \quad (0.37)$$

by taking the "average" around the maximum x_0 , we would expect

$$\oint_{B(x_0, r)} u(y) dy \leq M \quad (0.38)$$

Proof.

We will prove the second statement since the first followed from the second. Suppose $\exists x_0 \in U$ such that $u(x_0) = \max_{\bar{U}} u(x) \equiv M$. Then $\forall r \in (0, \text{dist}(x_0, \partial U))$, the mean-value property states that

$$M = u(x_0) = \oint_{B(x_0, r)} u(y) dy \quad (0.37)$$

by taking the "average" around the maximum x_0 , we would expect

$$\oint_{B(x_0, r)} u(y) dy \leq M \quad (0.38)$$

and such will hold only if $u \equiv M$ within $B(x_0, r)$, and $u(y) = M$. □

An important application of maximum principle is establishing the uniqueness of solutions to certain boundary value problems for Poisson's equation.

An important application of maximum principle is establishing the uniqueness of solutions to certain boundary value problems for Poisson's equation.

Theorem

Let $g \in C(\partial U)$, $f \in C(U)$, then there exists at most one solution $u \in C^2(U) \cap C(\bar{U})$ of the boundary value problem

$$\begin{cases} -\Delta u = f & \text{in } U \\ u = g & \text{on } \partial U \end{cases} \quad (0.39)$$

Proof.

Assume u_1, u_2 both solves the boundary value problem, then let $w = u_1 - u_2$, then we have

$$\begin{cases} -\Delta w = 0 & \text{in } U \\ w = 0 & \text{in } \partial U \end{cases} \quad (0.40)$$

then by maximum principle, $\max_{\partial U} w = \max_{\partial \bar{U}} w = 0$, i.e $u_1 = u_2$. \square

A nice thing about harmonic functions is that they are smooth!

A nice thing about harmonic functions is that they are smooth!

Theorem

If $u \in C(U)$ satisfies the mean-value property for each ball $B(x, r) \in U$, then $u \in C^\infty(U)$.

A nice thing about harmonic functions is that they are smooth!

Theorem

If $u \in C(U)$ satisfies the mean-value property for each ball $B(x, r) \in U$, then $u \in C^\infty(U)$.

The proof of this theorem requires some knowledge on mollifiers, so before the proof we shall spend some time on mollifiers.

Definition

Define a smooth function $\eta \in C^\infty(\mathbb{R}^n)$ by

$$\eta(x) = \begin{cases} C \exp\left(\frac{1}{|x|^2-1}\right) & |x| < 1 \\ 0 & |x| \geq 1 \end{cases} \quad (0.41)$$

for some constant $C > 0$ so that $\int_{\mathbb{R}^n} \eta dx = 1$. Then for each $\epsilon > 0$, set

$$\eta_\epsilon(x) = \frac{1}{\epsilon^n} \eta\left(\frac{x}{\epsilon}\right). \quad (0.42)$$

We call η the standard mollifier, the functions η_ϵ are C^∞ as well.

Definition

Denote $U_\epsilon = \{x \in U \mid \text{dist}(x, \partial U) > \epsilon\}$, if a function $f : U \rightarrow \mathbb{R}$ is locally integrable, its mollification is defined by

$$f^\epsilon = \int_U \eta_\epsilon(x - y) f(y) dy \quad (0.43)$$

for $x \in U_\epsilon$.

Definition

Denote $U_\epsilon = \{x \in U \mid \text{dist}(x, \partial U) > \epsilon\}$, if a function $f : U \rightarrow \mathbb{R}$ is locally integrable, its mollification is defined by

$$f^\epsilon = \int_U \eta_\epsilon(x - y) f(y) dy \quad (0.43)$$

for $x \in U_\epsilon$.

Now we prove the theorem.

Let η be the standard mollifier, set $u^\epsilon = \eta_\epsilon * u$ in U_ϵ , we first show $u^\epsilon \in C^\infty$, which is left as an exercise to the reader. Then to show u is smooth, we will show in fact $u \equiv u^\epsilon$ on U_ϵ for each $\epsilon > 0$. Let $x \in U_\epsilon$, then we know the support of $\eta_\epsilon(x - y)$ as a function of y is given by $\{y : |x - y| < \epsilon\}$, and thus we have

Let η be the standard mollifier, set $u^\epsilon = \eta_\epsilon * u$ in U_ϵ , we first show $u^\epsilon \in C^\infty$, which is left as an exercise to the reader. Then to show u is smooth, we will show in fact $u \equiv u^\epsilon$ on U_ϵ for each $\epsilon > 0$. Let $x \in U_\epsilon$, then we know the support of $\eta_\epsilon(x - y)$ as a function of y is given by $\{y : |x - y| < \epsilon\}$, and thus we have

$$\begin{aligned}
 u^\epsilon(x) &= \frac{1}{\epsilon^n} \int_{B(x, \epsilon)} \eta\left(\frac{|x - y|}{\epsilon}\right) u(y) dy \\
 &= \frac{1}{\epsilon^n} \int_0^\epsilon \eta\left(\frac{r}{\epsilon}\right) \left(\int_{\partial B(x, r)} u dS \right) dr \\
 &= \frac{1}{\epsilon^n} u(x) \int_0^\epsilon \eta\left(\frac{r}{\epsilon}\right) n\alpha(n) r^{n-1} dr \\
 &= u(x) \int_{B(0, \epsilon)} \eta_\epsilon dy \\
 &= u(x).
 \end{aligned}$$

Let η be the standard mollifier, set $u^\epsilon = \eta_\epsilon * u$ in U_ϵ , we first show $u^\epsilon \in C^\infty$, which is left as an exercise to the reader. Then to show u is smooth, we will show in fact $u \equiv u^\epsilon$ on U_ϵ for each $\epsilon > 0$. Let $x \in U_\epsilon$, then we know the support of $\eta_\epsilon(x - y)$ as a function of y is given by $\{y : |x - y| < \epsilon\}$, and thus we have

$$\begin{aligned} u^\epsilon(x) &= \frac{1}{\epsilon^n} \int_{B(x, \epsilon)} \eta\left(\frac{|x - y|}{\epsilon}\right) u(y) dy \\ &= \frac{1}{\epsilon^n} \int_0^\epsilon \eta\left(\frac{r}{\epsilon}\right) \left(\int_{\partial B(x, r)} u dS \right) dr \\ &= \frac{1}{\epsilon^n} u(x) \int_0^\epsilon \eta\left(\frac{r}{\epsilon}\right) n\alpha(n) r^{n-1} dr \\ &= u(x) \int_{B(0, \epsilon)} \eta_\epsilon dy \\ &= u(x). \end{aligned}$$

so $u \in C^\infty(U_\epsilon)$ for all $\epsilon > 0$.

Liouville's Theorem

Theorem

If $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is harmonic and bounded (either from above or below), then u is constant.

Liouville's Theorem

Theorem

If $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is harmonic and bounded (either from above or below), then u is constant.

To prove Liouville's theorem, we first introduce a technical lemma for estimating the derivatives of harmonic functions:

Liouville's Theorem

Theorem

If $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is harmonic and bounded (either from above or below), then u is constant.

To prove Liouville's theorem, we first introduce a technical lemma for estimating the derivatives of harmonic functions:

Lemma

Assume u is harmonic in U , then

$$|D^\alpha u(x_0)| \leq \frac{C_k}{r^{n+k}} \|u\|_{L^1(B(x_0, r))}, \quad C_0 = \frac{1}{\alpha(n)}, \quad C_k = \frac{(2^{n+1}nk)^k}{\alpha(n)} \quad (0.44)$$

for each ball $B(x_0, r) \in U$ and each multiindex α of order $|\alpha| = k$

Liouville's Theorem

Theorem

If $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is harmonic and bounded (either from above or below), then u is constant.

To prove Liouville's theorem, we first introduce a technical lemma for estimating the derivatives of harmonic functions:

Lemma

Assume u is harmonic in U , then

$$|D^\alpha u(x_0)| \leq \frac{C_k}{r^{n+k}} \|u\|_{L^1(B(x_0, r))}, \quad C_0 = \frac{1}{\alpha(n)}, \quad C_k = \frac{(2^{n+1}nk)^k}{\alpha(n)} \quad (0.44)$$

for each ball $B(x_0, r) \in U$ and each multiindex α of order $|\alpha| = k$

The proof to the lemma is left as an exercise.

Liouville's theorem can be done when $\alpha = 1$:

Liouville's theorem can be done when $\alpha = 1$:

Proof.

Fix $x_0 \in \mathbb{R}^n$ and $r > 0$, we have

$$|Du(x_0)| \leq \frac{C_1}{r^{n+1}} \|u\|_{L^1(B(x_0, r))} \leq \frac{C_1 \alpha(n)}{r} \|u\|_{L^\infty(\mathbb{R}^n)} \xrightarrow{r \rightarrow \infty} 0 \quad (0.45)$$

which means u is a constant. □

Liouville's theorem can be done when $\alpha = 1$:

Proof.

Fix $x_0 \in \mathbb{R}^n$ and $r > 0$, we have

$$|Du(x_0)| \leq \frac{C_1}{r^{n+1}} \|u\|_{L^1(B(x_0, r))} \leq \frac{C_1 \alpha(n)}{r} \|u\|_{L^\infty(\mathbb{R}^n)} \xrightarrow{r \rightarrow \infty} 0 \quad (0.45)$$

which means u is a constant. □

Liouville's theorem can also be used to conclude the uniqueness of Poisson's equation:

Liouville's theorem can be done when $\alpha = 1$:

Proof.

Fix $x_0 \in \mathbb{R}^n$ and $r > 0$, we have

$$|Du(x_0)| \leq \frac{C_1}{r^{n+1}} \|u\|_{L^1(B(x_0, r))} \leq \frac{C_1 \alpha(n)}{r} \|u\|_{L^\infty(\mathbb{R}^n)} \xrightarrow{r \rightarrow \infty} 0 \quad (0.45)$$

which means u is a constant. □

Liouville's theorem can also be used to conclude the uniqueness of Poisson's equation:

Theorem

Let $f \in C_c^2(\mathbb{R}^n)$ and $n \geq 3$, then any bounded solution of $-\Delta u = f$ in \mathbb{R}^n has the form

$$u(x) = \int_{\mathbb{R}^n} \Phi(x - y) f(y) dy + C \quad (0.46)$$

for some constant C .

Proof.

If u' is another solution, then $u - u'$ is a constant by Liouville's theorem. □

Harnack's Inequality

We write $V \subset\subset U$ to denote $V \subset \bar{V} \subset U$ and \bar{V} is compact, we say V is compactly contained in U .

Harnack's Inequality

We write $V \subset\subset U$ to denote $V \subset \bar{V} \subset U$ and \bar{V} is compact, we say V is compactly contained in U .

Theorem

For each connected open set $V \subset\subset U$, there exists a positive constant C depending only on V such that

$$\sup_V u \leq C \inf_V u \quad (0.47)$$

for all non-negative harmonic functions u in U . In particular

$$\frac{1}{C}u(y) \leq u(x) \leq Cu(y) \quad (0.48)$$

for all $x, y \in V$.

Proof.

Consider a ball $B(a, r) \subset U$, and let $x, y \in B(a, \frac{1}{4}r)$, we first claim that $B(x, \frac{1}{4}r) \subset B(y, \frac{3}{4}r) \subset B(a, r)$.

Proof.

Consider a ball $B(a, r) \subset U$, and let $x, y \in B(a, \frac{1}{4}r)$, we first claim that $B(x, \frac{1}{4}r) \subset B(y, \frac{3}{4}r) \subset B(a, r)$. Then we use mean value property and we have

$$u(x) = \int_{B(x, \frac{1}{4}r)} u(z) dz \leq \int_{B(\frac{3}{4}r, y)} u(z) dz = 3^n u(y). \quad (0.49)$$

Proof.

Consider a ball $B(a, r) \subset U$, and let $x, y \in B(a, \frac{1}{4}r)$, we first claim that $B(x, \frac{1}{4}r) \subset B(y, \frac{3}{4}r) \subset B(a, r)$. Then we use mean value property and we have

$$u(x) = \int_{B(x, \frac{1}{4}r)} u(z) dz \leq \int_{B(\frac{3}{4}r, y)} u(z) dz = 3^n u(y). \quad (0.49)$$

Now since V is compact so there exists a finite cover of balls $\{B_i\}_{i=1}^M$ where $B_i \equiv B(a_i, r_i) \subset U$.

Proof.

Consider a ball $B(a, r) \subset U$, and let $x, y \in B(a, \frac{1}{4}r)$, we first claim that $B(x, \frac{1}{4}r) \subset B(y, \frac{3}{4}r) \subset B(a, r)$. Then we use mean value property and we have

$$u(x) = \int_{B(x, \frac{1}{4}r)} u(z) dz \leq \int_{B(y, \frac{3}{4}r)} u(z) dz = 3^n u(y). \quad (0.49)$$

Now since V is compact so there exists a finite cover of balls $\{B_i\}_{i=1}^M$ where $B_i \equiv B(a_i, r_i) \subset U$. Then for any $x, y \in V$, we may find a ordered subcover $B_{i_1}, \dots, B_{i_\ell}$ such that $x \in B_{i_1}$, $y \in B_{i_\ell}$ and $B_{ij} \cap B_{ij+1} \neq \emptyset$, and the previous conclusion holds.

Proof.

Consider a ball $B(a, r) \subset U$, and let $x, y \in B(a, \frac{1}{4}r)$, we first claim that $B(x, \frac{1}{4}r) \subset B(y, \frac{3}{4}r) \subset B(a, r)$. Then we use mean value property and we have

$$u(x) = \int_{B(x, \frac{1}{4}r)} u(z) dz \leq \int_{B(\frac{3}{4}r, y)} u(z) dz = 3^n u(y). \quad (0.49)$$

Now since V is compact so there exists a finite cover of balls $\{B_i\}_{i=1}^M$ where $B_i \equiv B(a_i, r_i) \subset U$. Then for any $x, y \in V$, we may find a ordered subcover $B_{i_1}, \dots, B_{i_\ell}$ such that $x \in B_{i_1}$, $y \in B_{i_\ell}$ and $B_{ij} \cap B_{ij+1} \neq \emptyset$, and the previous conclusion holds. Then we have a chain of inequalities:

$$u(x) \leq 3^n u(x_1) \leq \dots \leq 3^{n\ell} u(y) \leq 3^{nM} u(y) \quad (0.50)$$

for all $x, y \in V$,

Proof.

Consider a ball $B(a, r) \subset U$, and let $x, y \in B(a, \frac{1}{4}r)$, we first claim that $B(x, \frac{1}{4}r) \subset B(y, \frac{3}{4}r) \subset B(a, r)$. Then we use mean value property and we have

$$u(x) = \int_{B(x, \frac{1}{4}r)} u(z) dz \leq \int_{B(y, \frac{3}{4}r)} u(z) dz = 3^n u(y). \quad (0.49)$$

Now since V is compact so there exists a finite cover of balls $\{B_i\}_{i=1}^M$ where $B_i \equiv B(a_i, r_i) \subset U$. Then for any $x, y \in V$, we may find a ordered subcover $B_{i_1}, \dots, B_{i_\ell}$ such that $x \in B_{i_1}$, $y \in B_{i_\ell}$ and $B_{ij} \cap B_{ij+1} \neq \emptyset$, and the previous conclusion holds. Then we have a chain of inequalities:

$$u(x) \leq 3^n u(x_1) \leq \dots \leq 3^{n\ell} u(y) \leq 3^{nM} u(y) \quad (0.50)$$

for all $x, y \in V$, and hence we have

$$\sup_V u \leq C \inf_V u. \quad (0.51)$$

Theorem

(Weierstrass Convergence Theorem) If a sequence of harmonic functions $\{h_n\}$ converges locally uniformly on U , then the limit $h = \lim_{n \rightarrow \infty} h_n$ is also harmonic on U , furthermore the derivatives converge locally uniformly as well, i.e $\lim_{n \rightarrow \infty} D_{x_i} h_n = D_{x_i} h$.

Proof.

Let $K \subset U$ be compact, we choose a larger compact set \tilde{K} and $r := \text{dist}(K, \mathbb{R}^n / \tilde{K}) > 0$, we use the derivative estimate to get

$$|D^\alpha h_l(a) - D^\alpha h_k(a)| \leq \frac{C}{r^{|\alpha|}} \sup_{\tilde{K}} |h_l - h_k| \quad (0.52)$$

and from the uniform convergence of $\{h_n\}$ on \tilde{K} , and we see that $\{D^\alpha h_n\}$ is a Cauchy sequence on K , thus uniformly convergent on K , and locally uniformly convergent on U . We can also show that $\lim_{n \rightarrow \infty} D^\alpha h_n = D^\alpha h$ locally uniformly on U for all index α . In particular we have $\Delta h = \lim_{n \rightarrow \infty} \Delta h_n$ and hence h is harmonic. □

Theorem

(Harnack Convergence Theorem) Consider an increasing sequence of harmonic functions $\{h_n\}_{n=1}^{\infty}$ on $U \subset \mathbb{R}^n$, then either $\lim_{n \rightarrow \infty} h_n = \infty$ or it converge locally uniformly to a harmonic function. In particular, if $\exists x_0 \in U$ such that $\lim_{n \rightarrow \infty} h_n(x_0) \neq \infty$, then we may conclude locally uniform convergence.

Proof.

Assume $x_0 \in U$ such that $\lim_{n \rightarrow \infty} h(x_0) \neq \infty$. For all $k \leq l$, K compact, and $x_0 \in K \subset U$, we have $h_l - h_k$ as a non-negative harmonic function, and hence by Harnack's inequality,

$$\max_K (h_l - h_k) \leq C \min_K (h_l - h_k) \leq C(h_l(x_0) - h_k(x_0)) \quad (0.53)$$

which implies the uniform Cauchy property of $\{h_n\}$ on K , and hence locally uniform convergence is achieved. We now let $\lim_{n \rightarrow \infty} h_n = h$ and it remains to show h is harmonic, which follows from Weierstrass convergence theorem.



Thanks