

# INTRODUCTION TO HOMOLOGICAL ALGEBRA

ROBERT CARDONA, MASSY KHOSHBIN, AND SIAVASH MORTEZAVI

## 1. MATH 697 NOTES

**AM Proposition 1.1:** There is a one-to-one order-preserving correspondence between the ideals  $I$  of  $R$  which contain  $(r)$  and the ideals  $\bar{I}$  of  $R/(r)$ .

*Proof.* Define  $\varphi : \{I \text{ ideal of } R : (r) \subseteq I\} \rightarrow \{\bar{I} : \bar{I} \text{ is an ideal of } R/(r)\}$  by  $\varphi(I) = I + (r)$ . We must first show that this mapping is well defined, i.e., does  $\varphi$  in fact map into an ideal of  $R/(r)$ ? Say  $I$  is an ideal of  $R$  with  $(r) \subseteq I$ . Consider  $\varphi(I) = I + (r)$ . Choose  $t \in R$  arbitrary and conclude that  $[s + (r)][I + (r)] = sI + (r)$  but since  $I$  is an ideal of  $R$ ,  $sI = I$  so  $[s + (r)][I + (r)] = sI + (r) = [s + (r)]\bar{I} \subseteq \bar{I}$ . Hence  $\varphi$  is in fact well-defined.

Now we want to show that  $\varphi$  is surjective. Choose  $\bar{I}$  an ideal of  $R/(r)$ . Does there exist an ideal  $I$  of  $R$  containing  $(r)$  such that  $\varphi(I) = \bar{I}$ ? Since  $\bar{I}$  is an ideal of  $R/(r)$ , it follows that  $\bar{I} = \{s + (r) : \text{for some } s \in R\}$ . Define  $I = \{s\}$  where  $s$  comes from previous definition. Is this in fact an ideal of  $R$ ? Well we need to first show that it is an additive subgroup:

- (1) Observe that  $0 \in \bar{I}$  since  $(r) \in \bar{I}$ . so  $0 \in I$  so  $I \neq \emptyset$ .
- (2) Let  $s, t \in I$ , Then  $s + t + (r) = [s + (r)] + [t + (r)] \in \bar{I}$  since  $\bar{I}$  is closed under addition. So  $s + t \in I$ .
- (3) Let  $s \in I$ , then  $s + (r) \in \bar{I}$  and since  $\bar{I}$  is an additive subgroup it follows that  $-s + (r) \in \bar{I}$  so  $-s \in I$ .

Conclude by definition of subgroup that  $I$  is in fact an additive subgroup. Now we must show that  $I$  is closed under multiplication. Say  $t \in R$ , is  $ts \in I$ ? well observe that since  $s + (r) \in \bar{I}$  and  $\bar{I}$  is an ideal, it follows that  $t[s + (r)] = ts + t(r) = ts + (r) \in \bar{I}$  so it follows that  $ts \in I$ . Conclude that  $I$  is in fact an ideal. Now we must ask, is  $(r) \subseteq I$ ? Let  $x \in (r)$  then  $x = rt$  for some  $t \in R$ . So  $rs + (r) = 0 + (r) \in \bar{I}$ . So it follows that  $(r) \subseteq I$ . Now  $\varphi(I) = I + (r) = \bar{I}$  so  $\varphi$  is in fact surjective.

Now we must show that  $\varphi$  is injective. Suppose that  $\varphi(I) = \bar{0} = 0 + (r) = (r)$ . Then since  $\varphi(I) = I + (r) = 0 + (r)$  it follows that  $I \subseteq (r)$ . But since we know  $(r) \subseteq I$  it follows that  $I + (r)$  which is our *zero* in this case. Conclude that  $\varphi$  is in fact injective.

*Alternate Injective Proof:* Suppose that  $\varphi(I_1) = \varphi(I_2)$  then we have  $I_1 + (r) = I_2 + (r)$  but by property of cosets we have  $I_1 - I_2 = (r)$  but this is the same as  $I_1 + I_2 = (r)$  by way of just absorbing the negative sign (being an ideal). But since  $(r) \subseteq I_1$  and  $(r) \subseteq I_2$  it follows that  $I_1 = I_2$ . Conclude that  $\varphi$  is injective.

Now we must show that order is preserved. Suppose  $I_1 \subseteq I_2$ . We want to show that  $\varphi(I_1) \subseteq \varphi(I_2)$ . Say  $x + (r) \in \varphi(I_1)$  then it follows that  $x \in I_1 = I_2$  so immediately  $x + (r) \in I_2 + (r) = \varphi(I_2)$ . Conclude result. □

### R Proposition 2.18:

- (1) A sequence  $0 \rightarrow A \xrightarrow{f} B$  is exact if and only if  $f$  is injective.
- (2) A sequence  $B \xrightarrow{g} C \rightarrow 0$  is exact if and only if  $g$  is surjective.
- (3) A sequence  $0 \rightarrow A \xrightarrow{h} B \rightarrow 0$  is exact if and only if  $h$  is an isomorphism.

**DF §10.5 Proposition 24:** (*The Short Five Lemma*) Let  $\alpha, \beta, \gamma$  be homomorphisms of short exact sequences:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{f} & B & \xrightarrow{g} & C & \longrightarrow & 0 \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \\ 0 & \longrightarrow & A' & \xrightarrow{f'} & B' & \xrightarrow{g'} & C' & \longrightarrow & 0 \end{array}$$

- (1) If  $\alpha$  and  $\gamma$  are injective then so is  $\beta$ .

*Proof.* Let  $b \in B$  such that  $\beta(b) = 0$ . We want to show that  $b = 0$ . Observe that  $g'(\beta(b)) = g'(0) = 0$ . By commutativity we have  $\gamma(g(b)) = g'(\beta(b)) = g'(0) = 0$ . Since  $\gamma$  is injective we know  $g(b) = 0$  so  $b \in \ker g$  but since we are in an exact sequence we have  $\text{im } f = \ker g$  and hence  $b \in \text{im } f$ . By definition there exists  $a \in A$  with  $f(a) = b$ . Now  $f'(\alpha(a)) = \beta(f(a)) = \beta(b) = 0$ . Since  $f'$  is injective, it follows that  $\alpha(a) = (f')^{-1}(0) = 0$ . Now we have  $a = \alpha^{-1}(0)$  and so  $a = 0$ . So  $0 = f(a) = b$ . □

- (2) If  $\alpha$  and  $\gamma$  are surjective then so is  $\beta$ .

*Proof.* Let  $b' \in B'$  then  $g'(b') \in C'$ . Since  $\gamma$  is surjective there exists  $c \in C$  such that  $\gamma(c) = g'(b')$ . Since this is an exact sequence,  $g$  is surjective so there exists  $b \in B$  such that  $g(b) = c$ . By equality we have  $\gamma(c) = \gamma(g(b)) = g'(b')$ . Now observe that

$$g'(b' - \beta(b)) = g'(b') - g'(\beta(b)) = g'(b') - \gamma(g(b)) = g'(b') - g'(b') = 0$$

So in particular  $b' - \beta(b) \in \ker g'$  but by exactness  $\text{im } f' = \ker g'$  so there exists  $a' \in A'$  such that  $f'(a') = b' - \beta(b)$ . But since  $\alpha$  is surjective, there exists  $a \in A$  such that  $\alpha(a) = a'$ . Now  $f'(a') = f'(\alpha(a)) = \beta(f(a)) = b' - \beta(b)$ . By commutativity  $f'(\alpha(a)) = \beta(f(a)) = b' - \beta(b)$  so  $\beta(f(a)) + \beta(b) = b'$  and we have  $\beta(f(a) + b) = b'$ .  $\square$

- (3) If  $\alpha$  and  $\gamma$  are isomorphisms then so is  $\beta$  (and then the two sequences are isomorphism).

*Proof.* Follows from (1) and (2).  $\square$

**R Proposition 2.72:** (*Five Lemma*) Consider the commutative diagram with exact rows.

$$\begin{array}{ccccccccc} A_1 & \xrightarrow{f} & A_2 & \xrightarrow{g} & A_3 & \xrightarrow{h} & A_4 & \xrightarrow{k} & A_5 \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \delta & & \downarrow \varepsilon \\ B_1 & \xrightarrow{f'} & B_2 & \xrightarrow{g'} & B_3 & \xrightarrow{h'} & B_4 & \xrightarrow{k'} & B_5 \end{array}$$

- (1) If  $\beta$  and  $\delta$  are surjective and  $\varepsilon$  is injective, then  $\gamma$  is surjective.
- (2) If  $\beta$  and  $\delta$  are injective and  $\alpha$  is surjective, then  $\gamma$  is injective.
- (3) If  $\alpha, \beta, \delta$  and  $\varepsilon$  are isomorphisms, then  $\gamma$  is an isomorphism.

**R Exercise 2.14:** Let  $A \xrightarrow{f} B \xrightarrow{g} C$  be a sequence of module maps. Prove that  $gf = 0$  if and only if  $\text{im } f \subseteq \ker g$ . Give an example of such a sequence that is not exact.

*Proof.* Suppose  $gf = 0$ , that is,  $f(g(a)) = 0$  for all  $a \in A$ . Let  $b \in \text{im } f$  then by definition there exists  $a \in A$  such that  $f(a) = b$ . But we know by hypothesis that  $0 = g(f(a)) = g(b)$  so  $b \in \ker g$ . Conclude that  $\text{im } f \subseteq \ker g$ . Conversely, suppose that  $\text{im } f \subseteq \ker g$ . Let  $a \in A$  and observe that  $f(a) \in \text{im } f$ . By hypothesis  $f(a) \in \ker g$  so  $g(f(a)) = 0$ . Since  $a$  was arbitrary conclude  $gf = 0$ .  $\square$

Consider the sequence of module maps  $\mathbb{Z}/4\mathbb{Z} \xrightarrow{f} \mathbb{Z}/3\mathbb{Z} \xrightarrow{g} \mathbb{Z}/2\mathbb{Z}$  where  $f(\bar{x}) = 2\bar{x}$  and  $g(\bar{y}) = \bar{y}$ . Observe that  $\text{im } f = \mathbb{Z}/3\mathbb{Z}$  and  $\ker g = \{0\}$ . Since  $\text{im } f \neq \ker g$  it follows that this is **not** an exact sequence.

**R Exercise 2.15:**

- (1) Prove that  $f : M \rightarrow N$  is surjective if and only if  $\text{coker } f = \{0\}$ .

*Proof.* Suppose  $f : M \rightarrow N$  is surjective then for  $n \in N$ , there exists  $m \in M$  such that  $f(m) = n$ . By definition  $\text{coker } f = N/\text{im } f = N/M = 0$ . Conversely, suppose that  $\text{coker } f = 0$ , i.e.,  $N/\text{im } f = 0$  implying that if  $m + \text{im } f \in N/\text{im } f$  then  $m + \text{im } f = 0$  or equivalently  $m \in \text{im } f$ . Since  $m$  is arbitrary, conclude  $N = \text{im } f$  and hence  $f$  is surjective by definition.  $\square$

- (2) If  $f : M \rightarrow N$  is a map, prove that there is an exact sequence

$$0 \rightarrow \ker f \rightarrow M \xrightarrow{f} N \rightarrow \text{coker } f \rightarrow 0.$$

*Proof.* Define  $h : \ker f \rightarrow M$  by  $h(m) = m$ , that is, map each element to itself. It follows immediately that  $\text{im } h = \ker f$ . Define  $g : N \rightarrow \text{coker } f = N/\text{im } f$  by  $g(n) = n + \text{im } f$ , that is, the canonical/projection mapping. Observe that  $\ker g = \text{im } f$ . Conclude that the following sequence is in fact exact:

$$0 \xrightarrow{\text{identity}} \ker f \xrightarrow{h} M \xrightarrow{f} N \xrightarrow{g} \text{coker } f \xrightarrow{\text{zero}} 0. \quad \square$$

**R Exercise 2.16:**

- (1) If  $0 \rightarrow M \rightarrow 0$  is an exact sequence, prove that  $M = \{0\}$ .

*Proof.* Consider  $0 \xrightarrow{f} M \xrightarrow{g} 0$ . Since  $f$  is surjective then for  $m \in M$  there exists  $x \in 0$  such that  $f(x) = m$  but  $x$  must be  $0$  so  $m = 0$ .  $\square$

- (2) If  $A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} D$  is an exact sequence, prove that  $f$  is surjective if and only if  $h$  is injective.

*Proof.* Suppose  $f$  is surjective. Then  $\text{im } f = B = \ker g$  but this immediately implies that  $\text{im } g = 0 = \ker h$  so  $h$  is injective by definition. Conversely, suppose  $h$  is injective. Then  $\ker h = 0 = \text{im } g$  which immediately implies  $\ker g = B = \text{im } f$ . Conclude by definition  $f$  is surjective.  $\square$

- (3) Let  $A \xrightarrow{\alpha} B \xrightarrow{\beta} C \xrightarrow{\gamma} D \xrightarrow{\delta} E$  be exact. If  $\alpha$  and  $\delta$  are isomorphisms, prove that  $C = \{0\}$ .

*Proof.* Observe that, by previous exercise,  $\beta$  is surjective and  $\gamma$  is injective so we have  $\text{im } \beta = C$  and  $\ker \gamma = 0$ . Result follows by exactness:  $C = \text{im } \beta = \ker \gamma = 0$ . Conclude  $C = \{0\}$ .  $\square$

**R Exercise 2.17:** If  $A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} D \xrightarrow{k} E$  is exact, prove that there is an exact sequence  $0 \rightarrow \text{coker } f \xrightarrow{\alpha} C \xrightarrow{\beta} \ker k \rightarrow 0$ , where  $\alpha : b + \text{im } f \mapsto g(b)$  and  $\beta : c \mapsto h(c)$ .

*Proof.* Observe that  $\ker \beta = \{c \in C : \beta(c) = h(c) = 0\} = \ker h = \text{im } g$  by exactness and since  $\alpha$  can run through any  $b \in B$  conclude  $\text{im } \alpha = \text{im } g = \ker \beta$   $\square$

**R Exercise 2.32:** ( $3 \times 3$  Lemma) Consider the following commutative diagram in  ${}_R \mathbf{Mod}$  having exact columns.

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & A' & \xrightarrow{\alpha_1} & A & \xrightarrow{\alpha_2} & A'' \longrightarrow 0 \\
 & & \downarrow f' & & \downarrow f & & \downarrow f'' \\
 0 & \longrightarrow & B' & \xrightarrow{\beta_1} & B & \xrightarrow{\beta_2} & B'' \longrightarrow 0 \\
 & & \downarrow g' & & \downarrow g & & \downarrow g'' \\
 0 & \longrightarrow & C' & \xrightarrow{\gamma_1} & C & \xrightarrow{\gamma_2} & C'' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

If the bottom two rows are exact, prove that the top row is exact; if the top two rows are exact, prove that the bottom row is exact.

*Proof.* Suppose that the bottom two rows are exact. We must show four things:

- (1)  $\alpha_1$  is injective: Suppose  $a' \in A'$  with  $\alpha_1(a') = 0$ . Observe that by commutativity  $f(\alpha_1(a')) = \beta_1(f'(a')) = 0$  so since  $\beta_1$  is injective, by exactness of the second row,  $f'(a') = 0$ . But since the first column is exact by hypothesis, we have  $a' = 0$ . Conclude that  $\alpha_1$  is in fact injective.
- (2)  $\text{im } \alpha_1 \subseteq \ker \alpha_2$ : Choose  $a \in \text{im } \alpha_1$  then by definition there exists  $a' \in A'$  such that  $\alpha_1(a') = a$ . Observe that  $f(a) = f(\alpha_1(a')) = \beta_1(f'(a'))$  by commutativity. Moreover  $\beta_2(f(a)) = f''(\alpha_2(a))$  again by commutativity. Hence we have

$$\beta_2(f(a)) = \beta_2(\beta_1(f'(a'))) = 0 = f''(\alpha_2(a)).$$

Since  $f''$  is injective by exactness of the third column,  $\alpha_2(a) = 0$  and so  $a \in \ker \alpha_2$ . Conclude that  $\text{im } \alpha_1 \subseteq \ker \alpha_2$  as desired.

- (3)  $\ker \alpha_2 \subseteq \text{im } \alpha_1$ : Let  $a \in \ker \alpha_2$  then by definition  $\alpha_2(a) = 0$  and so  $f''(\alpha_2(a)) = \beta_2(f(a)) = 0$ . So  $f(a) \in \ker \beta_2 = \text{im } \beta_1$  by exactness of the second column, so there exists  $b' \in B'$  such that  $\beta_1(b') = f(a)$ . Now

$$\gamma_1(g'(b')) = g(\beta_1(b')) = g(f(a)) = 0$$

by commutativity and exactness so there exists  $a' \in A'$  such that  $f'(a') = b'$ . Now by commutativity

$$f(\alpha_1(a')) = \beta_1(f'(a')) = \beta_1(b') = f(a)$$

and so  $f(a - \alpha_1(a')) = 0$  since  $f$  is a homomorphism.  $f$  is injective since the second column is exact so it follows that  $\alpha_1(a') = a$  and so  $a \in \text{im } \alpha_1$ . Conclude that  $\ker \alpha_2 \subseteq \text{im } \alpha_1$  is in fact true.

- (4)  $\alpha_2$  is surjective: Choose  $a'' \in A''$ . Consider  $f''(a'') \in B''$ . Since the second row is exact  $\beta_2$  is surjective so there exists  $b \in B$  such that  $\beta_2(b) = f''(a'')$ . By commutativity  $g''(\beta_1(b)) = \gamma_2(g(b))$  but by exactness of the third column  $g''(\beta_2(b)) = g''(f''(a'')) = 0$ . So  $\gamma_2(g(b)) = 0$  which implies that  $g(b) \in \ker \gamma_2 = \text{im } \gamma_1$  by exactness of the third row. So there exists  $c' \in C'$  such that  $\gamma_1(c') = g(b)$ . Since the first column is exact  $g'$  is surjective so there exists  $b' \in B'$  such that  $g'(b') = c'$ . So  $\gamma_1(c') = \gamma_1(g'(b')) = g(\beta_1(b'))$ . Observe that

$$g(b - \beta_1(b')) = g(b) - g(\beta_1(b')) = 0$$

since  $g(b) = \gamma_1(c)$  and  $g(\beta_1(b')) = \gamma_1(c)$  also. So  $b - \beta_1(b') \in \ker g = \text{im } f$  by exactness of the second column. So there exists  $a \in A$  such that  $f(a) = b - \beta_1(b')$ . Now since  $f''(\alpha_2(a)) = \beta_2(f(a)) = \beta_2(b - \beta_1(b'))$  we get

$$f''(\alpha_2(a)) = \beta_2(b) - \beta_2(\beta_1(b')) = \beta_2(b) = f''(a'').$$

In particular we get  $f''(\alpha_2(a) - a'') = 0$  but  $f''$  is injective by exactness of the third column so  $\alpha_2(a) = a''$ . Conclude that  $\alpha_2$  is in fact surjective.

We have shown that if the bottom two rows are exact, then the top row is exact.

Suppose that the top two rows are exact. We must show four things:

- (1)  $\gamma_1$  is injective: Suppose  $c' \in C'$  with  $\gamma_1(c') = 0$ . Since the first column is exact,  $g'$  is surjective so there exists  $b' \in B'$  such that  $g'(b') = c'$ . Now by commutativity

$$0 = \gamma_1(c') = \gamma_1(g'(b')) = g(\beta_1(b')).$$

So  $\beta_1(b') \in \ker g = \operatorname{im} f$  since the second row is exact. So there exists  $a \in A$  such that  $f(a) = \beta_1(b')$ . But by commutativity

$$f''(\alpha_2(a)) = \beta_2(f(a)) = \beta_2(\beta_1(b')) = 0$$

so  $\alpha_2(a) = 0$ . Since  $f''$  is injective, because the third column is exact. Now  $a \in \ker \alpha_2 = \operatorname{im} \alpha_1$  by exactness of the first row so there exists  $a' \in A'$  such that  $\alpha_2(a') = a$ . Now

$$f(a) = f(\alpha_1(a')) = \beta_1(f'(a')) = \beta_1(b')$$

So  $\beta_1(f'(a') - b') = 0$ . Since the second row is exact,  $\beta_1$  is injective so  $f'(a') - b = 0$ . Hence  $f'(a') = b'$ . Taking  $g'$  of both sides we get

$$0 = g'(f'(a')) = g'(b') = c'.$$

Hence  $\gamma_1$  is injective.

- (2)  $\operatorname{im} \gamma_1 \subseteq \ker \gamma_2$ : Let  $c \in \operatorname{im} \gamma_1$  then by definition there exists  $c' \in C'$  such that  $\gamma_1(c') = c$ . Since the first column is exact  $g'$  is surjective so there exists  $b' \in B'$  such that  $g'(b') = c'$ . Observe that

$$g(\beta_1(b')) = \gamma_1(g'(b')) = \gamma_1(c') = c$$

by commutativity. Again by commutativity we observe

$$\gamma_2(c) = \gamma_1(g(\beta_1(b'))) = g''(\beta_1(\beta_1(b'))) = g''(0) = 0$$

and so  $c \in \ker \gamma_2$ . Conclude that  $\operatorname{im} \gamma_1 \subseteq \ker \gamma_2$ .

- (3)  $\ker \gamma_2 \subseteq \operatorname{im} \gamma_1$ : Choose  $c \in \ker \gamma_2$  then by definition  $\gamma_2(c) = 0$ . Since the second column is exact,  $g$  is surjective. So there exists  $b \in B$  such that  $g(b) = c$ . By commutativity

$$g''(\beta_2(b)) = \gamma_2(g(b)) = \gamma_2(c) = 0.$$

So  $\beta_2(b) \in \ker g'' = \operatorname{im} f''$  by exactness of the third column so there exists  $a'' \in A''$  such that  $f''(a'') = \beta_2(b)$ . Since the first row is exact  $\alpha_2$  is surjective so there exists  $a \in A$  such that  $\alpha_2(a) = a''$ . Now

$$\beta_2(f(a)) = f''(\alpha_2(a)) = f''(a'') = \beta_2(b)$$

so  $\beta_2(f(a) - b) = 0$  which implies that  $f(a) - b \in \ker \beta_2 = \operatorname{im} \beta_1$  by exactness of the second row. So there exists  $b' \in B'$  such that  $\beta_1(b') = f(a) - b$ . But now

$$\gamma_1(g'(b')) = g(\beta_1(b')) = g(f(a) - b) = g(f(a)) - g(b) = -c$$

yielding  $\gamma_1(-g'(b')) = c$ . So  $c \in \operatorname{im} \gamma_1$  and we can conclude that  $\ker \gamma_2 \subseteq \operatorname{im} \gamma_1$ .

- (4)  $\gamma_2$  is surjective: Choose  $c'' \in C''$ . Since the third column is exact  $g''$  is surjective so there exists  $b'' \in B''$  such that  $g''(b'') = c''$ . But since the second row is exact  $\beta_2$  is surjective so there exists  $b \in B$  such that  $\beta_2(b) = b''$ . Now by commutativity

$$c'' = g''(b'') = g''(\beta_2(b)) = \gamma_2(g(b)).$$

Conclude that  $\gamma_2$  is in fact surjective.

We have shown that if the top two rows are exact, then the bottom row is exact. □

**AM Proposition 2.10:** (*Snake Lemma*) Let

$$\begin{array}{ccccccc} 0 & \longrightarrow & M' & \xrightarrow{g} & M & \xrightarrow{h} & M'' \longrightarrow 0 \\ & & \downarrow f' & & \downarrow f & & \downarrow f'' \\ 0 & \longrightarrow & N' & \xrightarrow{g'} & N & \xrightarrow{h'} & N'' \longrightarrow 0 \end{array}$$

be a commutative diagram of  $R$ -modules and homomorphisms, with the rows exact. Then there exists a sequence

$$0 \rightarrow \ker(f') \xrightarrow{\bar{u}} \ker(f) \xrightarrow{\bar{v}} \ker(f'') \xrightarrow{\bar{d}} \operatorname{coker}(f') \xrightarrow{\bar{u}'} \operatorname{coker}(f) \xrightarrow{\bar{v}'} \operatorname{coker}(f'') \rightarrow 0$$

in which  $\bar{u}, \bar{v}$  are restrictions of  $u, v$ , and  $\bar{u}', \bar{v}'$  are induced by  $u', v'$ .

*Proof.* We consider:

$$\begin{array}{ccccccc}
& 0 & & 0 & & 0 & \\
& \downarrow & & \downarrow & & \downarrow & \\
0 & \longrightarrow & \ker f' & \xrightarrow{\bar{u}} & \ker f & \xrightarrow{\bar{v}} & \ker f'' \\
& & \downarrow g' & & \downarrow g & & \downarrow g'' \\
0 & \longrightarrow & M' & \xrightarrow{u} & M & \xrightarrow{v} & M'' \longrightarrow 0 \\
& & \downarrow f' & & \downarrow f & & \downarrow f'' \\
0 & \longrightarrow & N' & \xrightarrow{u'} & N & \xrightarrow{v'} & N'' \longrightarrow 0 \\
& & \downarrow h' & & \downarrow h' & & \downarrow h' \\
& & \operatorname{coker} f' & \xrightarrow{\bar{u}'} & \operatorname{coker} f & \xrightarrow{\bar{v}'} & \operatorname{coker} f'' \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array}$$

□