

INTRODUCTION TO HOMOLOGICAL ALGEBRA

ROBERT CARDONA, MASSY KHOSHBIN, AND SIAVASH MORTEZAVI

0. MATH 697 HOMEWORK ZERO.TWO

AM 2.1: Show that $(\mathbb{Z}/m\mathbb{Z}) \otimes (\mathbb{Z}/n\mathbb{Z}) = 0$ if m and n are coprime.

Proof. Choose $a \otimes b \in \mathbb{Z}/m\mathbb{Z} \otimes \mathbb{Z}/n\mathbb{Z}$. Since m and n are coprime, there exist $s, t \in \mathbb{Z}$ such that $ms + nt = 1$. Observe that

$$a = a \cdot 1 = a(ms + nt) = ams + ant \equiv ant \pmod{m}.$$

Now observe that

$$a \otimes b = atn \otimes b = a \otimes nb = at \otimes 0 = 0.$$

We have shown that any simple tensor is zero, so any finite linear combination of simple tensors is zero. Conclude $(\mathbb{Z}/m\mathbb{Z}) \otimes (\mathbb{Z}/n\mathbb{Z}) = 0$. □

AM 2.2: Let R be a ring, I an ideal of R , M an R -module. Show that $(R/I) \otimes_R M$ is isomorphic to M/IM .

Proof. Define $\varphi : R/I \times M \rightarrow M/IM$ by $\varphi(r + I, m) = rm + IM$, which we shall henceforth write as $\varphi(\bar{r}, m) = \overline{rm}$. Let $(\bar{r}, m) = (\bar{s}, m)$. Then $\bar{r} = \bar{s} \implies r \in \bar{s} \implies r = s + i$, some $i \in I$. Then $\varphi(\bar{r}, m) = \overline{rm} = \overline{(s + i)m} = \overline{sm + im} = \overline{sm} + \overline{im} = \overline{sm} + \bar{0} = \overline{sm} = \varphi(\bar{s}, m)$. Thus φ is well-defined.

Observe $\varphi(\bar{r} + \bar{s}, m) = \overline{(r + s)m} = \overline{rm + sm} = \overline{rm} + \overline{sm} = \overline{rm} + \overline{sm} = \varphi(\bar{r}, m) + \varphi(\bar{s}, m)$. Similarly, $\varphi(\bar{r}, m + n) = \varphi(\bar{r}, m) + \varphi(\bar{r}, n)$. Lastly, $\varphi(\bar{r}\bar{s}, m) = \overline{(rs)m} = \overline{r(sm)} = \varphi(\bar{r}, sm)$. Thus φ is R -biadditive (In fact, φ is R -bilinear).

Now we are guaranteed a unique R -homomorphism $\phi : R/I \otimes_R M \rightarrow M/IM$ given by $\phi(\bar{r} \otimes m) = \overline{rm}$. Notice if we define $f : M/IM \rightarrow R/I \otimes_R M$ via $f(\bar{m}) = \bar{1} \otimes m$ then f is a \mathbb{Z} -homomorphism which makes $f \circ \phi$ and $\phi \circ f$ the identity map in $R/I \otimes_R M$ and M/IM , respectively. So ϕ has a two-sided inverse, hence a bijective function, and accordingly is an isomorphism when considered as an R -map. □

R 2.28: Let R be a domain with $Q = \text{Frac}(R)$, its field of fractions. If A is an R -module, prove that every element of $Q \otimes_R A$ has the form $q \otimes a$ for $q \in Q$ and $a \in A$ (i.e. every element is a simple tensor).

Proof. Let $\sum_1^n q_i \otimes a_i \in Q \otimes_R A$. We can write $\sum_1^n q_i \otimes a_i = \sum_1^n \frac{r_i}{s_i} \otimes a_i$ for $r_i, s_i \in R, s_i \neq 0$. Write $s = s_1 s_2 \cdots s_n$ and $\hat{s}_i = \frac{s}{s_i}$. Then $\sum_1^n \frac{r_i}{s_i} \otimes a_i = \sum_1^n (1 \cdot \frac{r_i}{s_i}) \otimes a_i = \sum_1^n (\frac{\hat{s}_i}{\hat{s}_i} \cdot \frac{r_i}{s_i}) \otimes a_i = \sum_1^n \frac{\hat{s}_i r_i}{\hat{s}_i} \otimes a_i = \sum_1^n (\frac{1}{\hat{s}}) \hat{s}_i r_i \otimes a_i = \sum_1^n \frac{1}{\hat{s}} \otimes (\hat{s}_i r_i) a_i = \frac{1}{\hat{s}} \otimes (\sum_1^n \hat{s}_i r_i a_i)$. □

R 2.32: Consider the following commutative diagram in $R\mathbf{Mod}$ having exact columns.

If the bottom two rows are exact, prove that the top row is exact; if the top two rows are exact, prove that the bottom row is exact.

Proof. α_1 is injective: Let $a' \in \ker \alpha_1$. Then $\alpha_1(a') = 0$. So $f(\alpha_1(a')) = 0$. Now $0 = f(\alpha_1(a')) = \beta_1(f'(a'))$ by commutativity. The injectivity of β_1 implies $f'(a') = 0$ and the injectivity of f' gives us $a' = 0$. Thus $\ker \alpha_1 = 0$ and α_1 is injective.

α_2 is surjective:

$\text{im } \alpha_1 \subseteq \ker \alpha_2$: Let $a \in \text{im } \alpha_1$. Then there exists $a' \in A'$ with $a = \alpha_1(a')$. Observe $f(a) = f(\alpha_1(a')) = \beta_1(f'(a'))$ by commutativity. Thus $\beta_2(f(a)) = \beta_2(\beta_1(f'(a')))$ by exactness. Now $0 = \beta_2(f(a)) = f''(\alpha_2(a))$ by commutativity. The injectivity of f'' gives us $\alpha_2(a) = 0$. Hence $a \in \ker \alpha_2$.

$\ker \alpha_2 \subseteq \text{im } \alpha_1$: Let $a \in \ker \alpha_2$. Then $\alpha_2(a) = 0$. So $f''(\alpha_2(a)) = 0$. By commutativity, $\beta_2(f(a)) = 0$. Now $f(a) \in \ker \beta_2 = \text{im } \beta_1$, so there exists $b' \in B'$ such that $f(a) = \beta_1(b')$. Now $g(f(a)) = 0$ by exactness, so $g(\beta_1(b')) = 0$. By commutativity, $\gamma_1(g'(b')) = 0$. Since γ_1 is injective, $g'(b') = 0$. Now $b' \in \ker g' = \text{im } f'$ so there exists $a' \in A'$ such that $b' = f'(a')$. Thus $f(a) = \beta_1(b') = \beta_1(f'(a'))$. By commutativity, $\beta_1(f'(a')) = f(\alpha_1(a'))$. So $f(a) = f(\alpha_1(a'))$. Since f is injective, we have $a = \alpha_1(a')$, and therefore $a \in \text{im } \alpha_1$. □

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY LONG BEACH

E-mail address: `mrrobertcardona@gmail.com` and `massy255@gmail.com` and `siavash.mortezavi@gmail.com`